Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra
MD (Pathology)
CEO & Consultant Pathologist

TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT.

NAME	**: Mrs. NEELAM**		
AGE/ GENDER	: 39 YRS/FEMALE	**PATIENT ID**	: 1738759
COLLECTED BY	:	**REG. NO./LAB NO.**	**: 012501290039**
REFERRED BY	:	**REGISTRATION DATE**	: 29/Jan/2025 12:21 PM
BARCODE NO.	: 01524618	**COLLECTION DATE**	: 29/Jan/2025 12:22PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	**REPORTING DATE**	: 29/Jan/2025 01:57PM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval

ENDOCRINOLOGY

LUTEINISING HORMONE (LH)

LUTEINISING HORMONE (LH): SERUM *by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)*	6.74	mIU/mL	MALES: 0.57 - 12.07 FOLLICULAR PHASE: 1.80 - 11.78 MID-CYCLE PEAK: 7.59 - 89.08 LUTEAL PHASE: 0.56 - 14.0 POST MENOPAUSAL WITHOUT HRT: 5.16 - 61.99

INTERPRETATION:

1. Luteinizing hormone (LH) is a glycoprotein hormone consisting of 2 non covalently bound subunits (alpha and beta). Gonadotropin-releasing hormone from the hypothalamus controls the secretion of the gonadotropins, FSH and LH, from the anterior pituitary.
2. In both males and females, LH is essential for reproduction. In females, the menstrual cycle is divided by a mid cycle surge of both LH and FSH into a follicular phase and a luteal phase.
3. This "LH surge" triggers ovulation thereby not only releasing the egg, but also initiating the conversion of the residual follicle into a corpus luteum that, in turn, produces progesterone to prepare the endometrium for a possiblei mplantation.
4. LH supports thecal cells in the ovary that provide androgens and hormonal precursors for estradiol production. LH in males acts on testicular interstitial cells of Leydig to cause increased synthesis of testosterone.

The test is useful in the following situations:

1. An adjunctin the evaluation of menstrual irregularities.
2. Evaluating patients with suspected hypogonadism
3. Predicting ovulation & Evaluating infertility
4. Diagnosing pituitary disorders
5. In both males and females, primary hypogonadism results in an elevation of basal follicle-stimulating hormone and luteinizing hormone levels.

FSH AND LH ELEVTED IN:

1. Primary gonadal failure
2. Complete testicular feminization syndrome
3. Precocious puberty (either idiopathic or secondary to a central nervous system lesion)
4. Menopause
5. Primary ovarian hypo dysfunction in females
6. Polycystic ovary disease in females
7. Primary hypogonadism in males

LH IS DECREASED IN:

1 .Primary ovarian hyper function in females
2. Primary hypergonadism in males

NOTE

1 .FSH and LH are both decreased in failure of the pituitary or hypothalamus.

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)

Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra
MD (Pathology)
CEO & Consultant Pathologist

TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT.

NAME	**: Mrs. NEELAM**		
AGE/ GENDER	: 39 YRS/FEMALE	**PATIENT ID**	: 1738759
COLLECTED BY	:	**REG. NO./LAB NO.**	**: 012501290039**
REFERRED BY	:	**REGISTRATION DATE**	: 29/Jan/2025 12:21 PM
BARCODE NO.	: 01524618	**COLLECTION DATE**	: 29/Jan/2025 12:22PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	**REPORTING DATE**	: 29/Jan/2025 01:57PM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval

FOLLICLE STIMULATING HORMONE (FSH)

FOLLICLE STIMULATING HORMONE (FSH): SERUM *by CLIA (CHEMILUMINESCENCE IMMUNOASSAY)*	7.55	mIU/mL	FEMALE FOLLICULAR PHASE: 2.5 - 11.4 FEMALE MID-CYCLE PEAK: 3.3 - 21.7 FEAMLE LUTEAL PHASE: 1.2 - 7.0 FEMALE POST-MENOPAUSAL: 18.8 - 132 MALE: 1.0 - 12.1

INTERPRETATION:

1. Gonadotropin-releasing hormone from the hypothalamus controls the secretion of the gonadotropins, follicle-stimulating hormone (FSH) and luteinizing hormone (LH) from the anterior pituitary.
2. The menstrual cycle is divided by a midcycle surge of both FSH and LH into a follicular phase and a luteal phase.
3. FSH appears to control gametogenesis in both males and females.

The test is useful in the following settings:

1. An adjunct in the evaluation of menstrual irregularities.
2. Evaluating patients with suspected hypogonadism.
3. Predicting ovulation
4. Evaluating infertility
5. Diagnosing pituitary disorders
6. In both males and females, primary hypogonadism results in an elevation of basal follicle-stimulating hormone (FSH) and luteinizing hormone (LH) levels.

FSH and LH LEVELS ELEVATED IN:

1. Primary gonadal failure
2. Complete testicular feminization syndrome.
3. Precocious puberty (either idiopathic or secondary to a central nervous system lesion)
4. Menopause (postmenopausal FSH levels are generally >40 IU/L)
5. Primary ovarian hypofunction in females
6. Primary hypogonadism in males

NOTE:

1. Normal or decreased FSH is seen in polycystic ovarian disease in females
2. FSH and LH are both decreased in failure of the pituitary or hypothalamus.

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)

Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra
MD (Pathology)
CEO & Consultant Pathologist

TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT.

NAME	**: Mrs. NEELAM**		
AGE/ GENDER	: 39 YRS/FEMALE	**PATIENT ID**	: 1738759
COLLECTED BY	:	**REG. NO./LAB NO.**	**: 012501290039**
REFERRED BY	:	**REGISTRATION DATE**	: 29/Jan/2025 12:21 PM
BARCODE NO.	: 01524618	**COLLECTION DATE**	: 29/Jan/2025 12:22PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	**REPORTING DATE**	: 29/Jan/2025 01:57PM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval

ANTI MULLERIAN HORMONE (AMH) GEN II

ANTI MULLERIAN HORMONE (AMH) GEN II: SERUM *by ECLIA (ELECTROCHEMILUMINESCENCE IMMUNOASSAY)*	3.229	ng/mL	0.05 - 11.00

INTERPRETATION:-

A Correlation of FERTILITY POTENTIAL and AMH levels are :

OVARIAN FERTILITY POTENTIAL	AMH VALUES IN (ng/mL)
OPTIMAL FERTILITY:	4.00 – 6.80 ng/mL
SATISFACTORY FERTILITY:	2.20 – 4.00 ng/mL
LOW FERTILITY:	0.30 – 2.20 ng/mL
VERY LOW/UNDETECTABLE:	0.00 – 0.30 ng/mL
HIGH LEVEL:	>6.8 ng/mL (PCOD/GRANULOSA CELL TUMOUR)

Anti Mullerian Hormone (AMH) is also known as Mullerian Inhibiting Substance provided by sertoli cells of the testis in males and by ovarian granulose cells in females upto antral stage in females.

IN MALES:
1.It is used to evaluate testicular presence and function in infants with intersex conditions or ambiguous genitalia, and to distinguish between cryptorchidism and anorchia in males

IN FEMALES:
1.During reproductive age, follicular AMH productionbegins during the primary stage, peaks in preantral stage & has influence on follicular sensitivity to FSH which is impoetant in selection for follicular dominance. AMH levels thus represents the pool or number of primordial follicles but not thequality of oocytes.AMH does not vary significantly during menstrual cycle & hence can be measured independently of day of cycle.
2.Polycystic ovarian syndrome can elevate AMH 2 to 5 fold higher than age specific reference range & predict anovulatory, irregular cycles, ovarian tumours like Granulosa cell tumour are often associated with higher AMH levels.
3.Obese women are often associated with diminished ovarian reserve and can have 65% lower mean AMH levels than non-obese women.
4.In females , AMH levels do not change significantly throughout the menstrual cycle and decrease with age.
5.Assess Ovarian Reserve - correlates with the number of antral follicies in the ovaries.
6.Evaluate fertility potential and ovarian response in IVF- Women with low AMG levels are more likely to the poor ovarian responders.
7.Assess the condition of Polycystic Ovary and premature ovarian failure.

A combination of Age, Ultrasound markers-Ovarian Volume and Antral Follicle Count, AMH and FSH levels are useful for optimal assessment of ovarian reserve. Studies in various fertility clinics are ongoing to establish optimal AMH concentretaion for predicting response to invitro fertilization, however,given below is suggested interpretative reference.

AMH levels (ng/mL)	Suggested patient	Anticipated Antral	Anticipated FSH levels	Anticipated Response

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)

Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra
MD (Pathology)
CEO & Consultant Pathologist

NAME	**: Mrs. NEELAM**		
AGE/ GENDER	: 39 YRS/FEMALE	**PATIENT ID**	: 1738759
COLLECTED BY	:	**REG. NO./LAB NO.**	**: 012501290039**
REFERRED BY	:	**REGISTRATION DATE**	: 29/Jan/2025 12:21 PM
BARCODE NO.	: 01524618	**COLLECTION DATE**	: 29/Jan/2025 12:22PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	**REPORTING DATE**	: 29/Jan/2025 01:57PM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	**Value**	**Unit**	**Biological Reference interval**

	Categorization for fertility based on AMH for age group (20 to 45 yrs)	**Follicle counts**	**(day 3)**	**to IVF/COH cycle**
Below 0.3	Very low	Below 4	Above 20	Negligible/Poor
0.3 to 2.19	Low	4 - 10	Usually 16 - 20	Reduced
2.19 t0 4.00	Satisfactory	11 - 25	Within reference range or between 11 - 15	Safe/Normal
Above 4.00	Optimal	Upto 30 and Above	Within reference range or between 11 – 15 or Above 15	Possibly Excessive

INCREASED:
1.Polycystic ovarian syndrome (most common)
2. Ovarian Tumour: Granulosa cell tumour

DECREASED:
1.Anorchia , Abnormal or absence of testis in males
2.Pseudohermaphroditism
3.Post Menopause

NOTE:
1.AMH measurement alone is seldom suffcient for diagnosis and results should be interpreted in the light of clinical finding and other relevant test such as ovarian ultrasonography(In fertility applications); abdominal or testicular ultrasound(intersex or testicular function applications); measurement of sex steroids (estradiol,Progesterone,Testosterone),FSH, Inhibin B (For fertility), and Inhibin A and B (for tumour work up).
2.Conversion of AMH grom ng/mL to pmol/L can be performed by using equation 1 ng/mL = 7.14 pmol/L

*** End Of Report ***

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)