

## **KOS Diagnostic Lab**

(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

**NAME** : Mr. NARESH

**AGE/ GENDER** : 64 YRS/MALE **PATIENT ID** : 1747462

**COLLECTED BY** REG. NO./LAB NO. :012502060028

REFERRED BY : P.G.I. (CHANDIGARH) **REGISTRATION DATE** : 06/Feb/2025 12:08 PM BARCODE NO. :01525052 **COLLECTION DATE** :06/Feb/2025 12:10PM CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 06/Feb/2025 01:02PM

**CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

**Value** Unit **Biological Reference interval Test Name** 

## IMMUNOPATHOLOGY/SEROLOGY **C-REACTIVE PROTEIN (CRP)**

C-REACTIVE PROTEIN (CRP) QUANTITATIVE: 1.36 0.0 - 6.0mg/L

by NEPHLOMETRY

## **INTERPRETATION:**

C-reactive protein (CRP) is one of the most sensitive acute-phase reactants for inflammation.

- 2. CRP levels can increase dramatically (100-fold or more) after severe trauma, bacterial infection, inflammation, surgery, or neoplastic
- 3. CRP levels (Quantitative) has been used to assess activity of inflammatory disease, to detect infections after surgery, to detect transplant rejection, and to monitor these inflammatory processes.

  4. As compared to ESR, CRP shows an earlier rise in inflammatory disorders which begins in 4-6 hrs, the intensity of the rise being higher than ESR and the recovery being earlier than ESR. Unlike ESR, CRP levels are not influenced by hematologic conditions like Anemia, Polycythemia etc., 5. Elevated values are consistent with an acute inflammatory process.

NOTE:

- 1. Elevated C-reactive protein (CRP) values are nonspecific and should not be interpreted without a complete clinical history.
- 2. Oral contraceptives may increase CRP levels.

End Of Report \*\*\*



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST

