

TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT.



NAME : Mrs. RANI BHATTI AGE / GENDER : 50 YRS/FEMALE PATIENT ID : 1755208 COLLECTED BY :			Chopra / & Microbiology) onsultant Pathologist	Dr. Yugam MD CEO & Consultant	(Pathology)
CULLECTED BY E. REG. NO./LAB NO. : 112502130004 REFERRED BY :: REGISTRATION DATE : 137Feb/2025 08:34 AM SARCODE NO. : 01525410 COLLECTION DATE : 137Feb/2025 08:34 AM SARCODE NO. : 01525410 COLLECTION DATE : 137Feb/2025 08:34 AM SILENT CODE : KOS DIAGNOSTIC LAB REPORTING DATE : 137Feb/2025 11:16AM SILENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT Biological Reference interva CLINICAL CHEMISTRY BIOCHEMISTRY BUCOSE FASTING (F): PLASMA 278.14 ^H mg/dL NORMAL: < 100.0 by GLUCOSE OXIDASE - PEROXIDASE (GOL-POD) 278.14 ^H mg/dL NORMAL: < 100.0 MACORDANCE WITH AMERICAN DIABETES ASSOCIATION GUIDELINES: . A fasting plasma glucose level below 100 mg/dl is considered normal. A fasting plasma glucose level Delow 125 mg/dl is seroindrened for all such patients. A fasting plasma glucose level 125 mg/dl is considered as glucose intolerant or prediabetic. A fasting and post-prandial blo est (after consumption of 75 gms of glucose) is recommended for all such patients. A fasting plasma glucose level 125 mg/dl is considered as glucose intoler	JAME	: Mrs. RANI BHATTI			
REFERENCE BY: E. E	AGE/ GENDER	: 56 YRS/FEMALE	PATIE	INT ID	: 1755268
ARCODE NO. : 19525412 CDLECTION DATE : 13/Feb/2025 08:34AM CLENT CODE : KOS DIAGNOSTIC LAB REPORTING DATE : 13/Feb/2025 11:16AM LIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT Test Name Value Unit Biological Reference interva CLENICAL CHEMISTRY/BIOCHEMISTRY GLUCOSE FASTING (F): PLASMA 278.14 ^H mg/dL NORMAL: < 100.0 PREDIABETIC: 100.0 - 125.0 DIABETIC: > 0 R = 126.0 NACCORDANCE VITH AMERICAN DIABETES ASSOCIATION GUIDELINES: . A fasting plasma glucose level below 100 mg/dl is considered normal. . A fasting plasma glucose level below 100 mg/dl is considered normal. . A fasting plasma glucose level below 100 mg/dl is considered normal. . A fasting plasma glucose level below 100 mg/dl is considered normal. . A fasting plasma glucose level of above 125 mg/dl is nighly suggestive of diabetic state. A repeat post-prandial is strongly recommended for the patients. A fasting plasma glucose level of above 125 mg/dl on both occasions is confirmatory for diabetic state. *** End Of Report ***	COLLECTED BY	:	REG. N	IO./LAB NO.	:012502130004
LIENT CODE : KOS DIAGNOSTIC LAB REPORTING DATE : 13/Feb/2025 11:16AM LIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT Test Name Value Unit Biological Reference interval CLINICAL CHEMISTRY/BIOCHEMISTRY GLUCOSE FASTING (F): CLINICAL CHEMISTRY/BIOCHEMISTRY SULCOSE FASTING (F): PLASMA 278,14 ^H mg/dL NORMAL: < 100.0	REFERRED BY	:	REGIS	TRATION DATE	: 13/Feb/2025 08:34 AM
CLENT ADRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT Test Name Value Unit Biological Reference interval CLENICAL CHEMISTRY/BIOCHEMISTRY CLENICAL CHEMISTRY/BIOCHEMISTRY CLENICAL CHEMISTRY/BIOCHEMISTRY CLUCOSE FASTING (F): PLASMA Transition NORMAL: < 100.0 by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD) Transition NORMAL: < 100.0 Machine Transition Mg/dL NORMAL: < 100.0 Display Organization Mg/dL NORMAL: < 100.0 Machine Transition plasma glucose (cooperotion) Mg/dL NORMAL: < 100.0 Machine Transition plasma glucose level below 100 mg/dl is considered normal. Ansting plasma glucose level below 125 mg/dl is considered normal. A fasting plasma glucose level below 100 mg/dl is considered normal. A fasting plasma glucose level of above 125 mg/dl is considered as glucose intolerant or prediabetic. A fasting nad post-prandial bloc est (after consumption of 75 gms of glucose) is recommended for all such patients. A fasting plasma glucose level of above 125 mg/dl is considered normal. A fasting plasma glucose level of above 125 mg/dl is nightly suggestive of diabetic state. A repeat post-prandial is strongly recommended for all such patients. A fasting plasma glucose level of above 125 mg/dl is nightly suggestive of diabetic state. A repeat post-prandial is strongly recommended for all such patients. <td>BARCODE NO.</td> <td>:01525410</td> <td>COLLE</td> <td>CTION DATE</td> <td>: 13/Feb/2025 08:34AM</td>	BARCODE NO.	:01525410	COLLE	CTION DATE	: 13/Feb/2025 08:34AM
Test Name Value Unit Biological Reference interval CLINICAL CHEMISTRY/BIOCHEMISTRY CLINICAL CHEMISTRY/BIOCHEMISTRY CLINICAL CHEMISTRY/BIOCHEMISTRY CLINICAL CHEMISTRY/BIOCHEMISTRY CLINICAL CHEMISTRY/BIOCHEMISTRY CLINICAL CHEMISTRY/BIOCHEMISTRY GLUCOSE FASTING (F): PLASMA by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD) 278.14 ^H mg/dL NORMAL: < 100.0		: KOS DIAGNOSTIC LAB	REPOI	RTING DATE	: 13/Feb/2025 11:16AM
CLINICAL CHEMISTRY/BIOCHEMISTRY GLUCOSE FASTING (F): SLUCOSE FASTING (F): PLASMA by GLUCOSE FASTING (F): SLUCOSE OXIDASE (GOD-POD) TREDIABETIC: 100.0 - 125.00 DIABETIC: 100.0 - 125.00 DIABETIC: 100.0 - 125.00 DIABETIC: > OR = 126.0 NTERPRETATION NACCORDANCE WITH AMERICAN DIABETES ASSOCIATION GUIDELINES: A fasting plasma glucose level belowen 100 mg/dl is considered normal. A fasting plasma glucose level between 100 - 125 mg/dl is considered as glucose intolerant or prediabetic. A fasting and post-prandial blo est (after consumption of 75 gms of glucose) is recommended for all such patients. A fasting plasma glucose level between 100 - 125 mg/dl is bighly suggestive of diabetic state. A repeat post-prandial is strongly recommended for all such patients. A fasting plasma glucose level evel in excess of 125 mg/dl on both occasions is confirmatory for diabetic state. *** End Of Report ***	LIENT ADDRESS	: 6349/1, NICHOLSON ROA	D, AMBALA CANTT		
GLUCOSE FASTING (F): PLASMA by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD) 278.14 ^H mg/dL NORMAL: < 100.0 PREDIABETIC: 100.0 - 125.0 DIABETIC: > 0R = 126.0 MACCORDANCE WITH AMERICAN DIABETES ASSOCIATION GUIDELINES: 1. A fasting plasma glucose level below 100 mg/dl is considered normal. 2. A fasting plasma glucose level between 100 - 125 mg/dl is considered as glucose intolerant or prediabetic. A fasting and post-prandial blo rest (after consumption of 75 gms of glucose) is recommended for all such patients. 3. A fasting plasma glucose level of above 125 mg/dl is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for such patients. A fasting plasma glucose level in excess of 125 mg/dl on both occasions is confirmatory for diabetic state. **** End Of Report ***	Fest Name		Value	Unit	Biological Reference interval
by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD) PREDIABETIC: 100.0 - 125.0 DIABETIC: > 0R = 126.0 WIERPRETATION VACCORDANCE WITH AMERICAN DIABETES ASSOCIATION GUIDELINES: . A fasting plasma glucose level below 100 mg/dl is considered normal. . A fasting plasma glucose level between 100 - 125 mg/dl is considered as glucose intolerant or prediabetic. A fasting and post-prandial blo est (after consumption of 75 gms of glucose) is recommended for all such patients. . A fasting plasma glucose level of above 125 mg/dl is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for uch patients. A fasting plasma glucose level in excess of 125 mg/dl on both occasions is confirmatory for diabetic state. *** End Of Report ***	LUCOSE FASTING	G (F): PLASMA			NORMAL: < 100.0
GLUCOSE FASTING (F): PLASMA by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD) 278.14 ^H mg/dL NORMAL: < 100.0 PREDIABETIC: 100.0 - 125.0 DIABETIC: > 0R = 126.0 MACCORDANCE WITH AMERICAN DIABETES ASSOCIATION GUIDELINES: 1. A fasting plasma glucose level below 100 mg/dl is considered normal. 2. A fasting plasma glucose level between 100 - 125 mg/dl is considered as glucose intolerant or prediabetic. A fasting and post-prandial blo rest (after consumption of 75 gms of glucose) is recommended for all such patients. 3. A fasting plasma glucose level of above 125 mg/dl is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for such patients. A fasting plasma glucose level in excess of 125 mg/dl on both occasions is confirmatory for diabetic state. **** End Of Report ***		CLIN	ICAL CHEMISTRY/	BIOCHEMIST	RY
DIABETTIC: > OR = 126.0 NACCORDANCE WITH AMERICAN DIABETES ASSOCIATION GUIDELINES: . A fasting plasma glucose level below 100 mg/dl is considered normal. . A fasting plasma glucose level between 100 - 125 mg/dl is considered as glucose intolerant or prediabetic. A fasting and post-prandial blo est (after consumption of 75 gms of glucose) is recommended for all such patients. . A fasting plasma glucose level of above 125 mg/dl is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for uch patients. A fasting plasma glucose level in excess of 125 mg/dl on both occasions is confirmatory for diabetic state. *** End Of Report ***			278.14 ^H	mg/dL	
	N ACCORDANCE WIT	H AMERICAN DIABETES ASSOCI	ATION GUIDELINES: is considered normal.		
	N ACCORDANCE WIT 1. A fasting plasma g 2. A fasting plasma g rest (after consumpti	lucose level below 100 mg/dl lucose level between 100 - 12 ion of 75 gms of glucose) is rec	s considered normal. 5 mg/dl is considered as glu ommended for all such pat	ients	prediabetic. A fasting and post-prandial blood
	N ACCORDANCE WIT . A fasting plasma g 2. A fasting plasma g est (after consumpti	lucose level below 100 mg/dl lucose level between 100 - 12 ion of 75 gms of glucose) is rec	is considered normal. 5 mg/dl is considered as glu ommended for all such pat dl is highly suggestive of dia ess of 125 mg/dl on both c	ients. abetic state. A repe occasions is confirm	prediabetic. A fasting and post-prandial blood
	N ACCORDANCE WIT . A fasting plasma g 2. A fasting plasma g est (after consumpti	lucose level below 100 mg/dl lucose level between 100 - 12 ion of 75 gms of glucose) is rec	is considered normal. 5 mg/dl is considered as glu ommended for all such pat dl is highly suggestive of dia ess of 125 mg/dl on both c	ients. abetic state. A repe occasions is confirm	prediabetic. A fasting and post-prandial blood
	N ACCORDANCE WIT A fasting plasma g A fasting plasma g est (after consumpti	lucose level below 100 mg/dl lucose level between 100 - 12 ion of 75 gms of glucose) is rec	is considered normal. 5 mg/dl is considered as glu ommended for all such pat dl is highly suggestive of dia ess of 125 mg/dl on both c	ients. abetic state. A repe occasions is confirm	prediabetic. A fasting and post-prandial blood
	N ACCORDANCE WIT . A fasting plasma g 2. A fasting plasma g est (after consumpti	lucose level below 100 mg/dl lucose level between 100 - 12 ion of 75 gms of glucose) is rec	is considered normal. 5 mg/dl is considered as glu ommended for all such pat dl is highly suggestive of dia ess of 125 mg/dl on both c	ients. abetic state. A repe occasions is confirm	prediabetic. A fasting and post-prandial blood
	N ACCORDANCE WIT . A fasting plasma g 2. A fasting plasma g est (after consumpti 3. A fasting plasma g uch patients. A fasti	lucose level below 100 mg/dl lucose level between 100 - 12 ion of 75 gms of glucose) is rec lucose level of above 125 mg/ ing plasma glucose level in exc	is considered normal. 5 mg/dl is considered as glu ommended for all such pat dl is highly suggestive of dia ess of 125 mg/dl on both c	ients. abetic state. A repe occasions is confirm	prediabetic. A fasting and post-prandial blood
	N ACCORDANCE WIT 1. A fasting plasma g 2. A fasting plasma g est (after consumpti 3. A fasting plasma g such patients. A fasti	lucose level below 100 mg/dl lucose level between 100 - 12 ion of 75 gms of glucose) is rec lucose level of above 125 mg/ ing plasma glucose level in exc	is considered normal. 5 mg/dl is considered as glu ommended for all such pat dl is highly suggestive of dia ess of 125 mg/dl on both c	ients. abetic state. A repe occasions is confirm	prediabetic. A fasting and post-prandial blood
	N ACCORDANCE WIT 1. A fasting plasma g 2. A fasting plasma g est (after consumpti 3. A fasting plasma g such patients. A fasti	lucose level below 100 mg/dl lucose level between 100 - 12 ion of 75 gms of glucose) is rec lucose level of above 125 mg/ ing plasma glucose level in exc	is considered normal. 5 mg/dl is considered as glu ommended for all such pat dl is highly suggestive of dia ess of 125 mg/dl on both c	ients. abetic state. A repe occasions is confirm	prediabetic. A fasting and post-prandial blood
	N ACCORDANCE WIT . A fasting plasma g 2. A fasting plasma g est (after consumpti 3. A fasting plasma g such patients. A fasti	lucose level below 100 mg/dl lucose level between 100 - 12 ion of 75 gms of glucose) is rec lucose level of above 125 mg/ ing plasma glucose level in exc	is considered normal. 5 mg/dl is considered as glu ommended for all such pat dl is highly suggestive of dia ess of 125 mg/dl on both c	ients. abetic state. A repe occasions is confirm	prediabetic. A fasting and post-prandial blood
	N ACCORDANCE WIT . A fasting plasma g 2. A fasting plasma g est (after consumpti 3. A fasting plasma g uch patients. A fasti	lucose level below 100 mg/dl lucose level between 100 - 12 ion of 75 gms of glucose) is rec lucose level of above 125 mg/ ing plasma glucose level in exc	is considered normal. 5 mg/dl is considered as glu ommended for all such pat dl is highly suggestive of dia ess of 125 mg/dl on both c	ients. abetic state. A repe occasions is confirm	prediabetic. A fasting and post-prandial blood

KOS Diagnostic Lab (A Unit of KOS Healthcare)





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)

 KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana

 KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt -133 001, Haryana

 0171-2643898, +91 99910 43898
 care@koshealthcare.com
 www.koshealthcare.com

