

Dr. Vinay Chopra  
MD (Pathology & Microbiology)  
Chairman & Consultant Pathologist

Dr. Yugam Chopra  
MD (Pathology)  
CEO & Consultant Pathologist

<b>NAME</b>	: Mr. AJIT BHALLA	<b>PATIENT ID</b>	: 508003
<b>AGE/ GENDER</b>	: 62 YRS/MALE	<b>REG. NO./LAB NO.</b>	: 012502240032
<b>COLLECTED BY</b>	:	<b>REGISTRATION DATE</b>	: 24/Feb/2025 10:21 AM
<b>REFERRED BY</b>	:	<b>COLLECTION DATE</b>	: 24/Feb/2025 10:23AM
<b>BARCODE NO.</b>	: 01526075	<b>REPORTING DATE</b>	: 26/Feb/2025 09:57AM
<b>CLIENT CODE.</b>	: KOS DIAGNOSTIC LAB		
<b>CLIENT ADDRESS</b>	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

### MICROBIOLOGY

#### CULTURE AEROBIC BACTERIA AND ANTIBIOTIC SENSITIVITY: URINE

##### CULTURE AND SUSCEPTIBILITY: URINE

DATE OF SAMPLE	24-02-2025
SPECIMEN SOURCE	URINE
INCUBATION PERIOD	48 HOURS
<i>by AUTOMATED BROTH CULTURE</i>	
CULTURE	STERILE
<i>by AUTOMATED BROTH CULTURE</i>	
ORGANISM	NO AEROBIC PYOGENIC ORGANISM GROWN AFTER 48 HOURS OF INCUBATION AT 37°C
<i>by AUTOMATED BROTH CULTURE</i>	

##### AEROBIC SUSCEPTIBILITY: URINE

###### INTERPRETATION:

1. In urine culture and sensitivity, presence of more than 100,000 organism per mL in midstream sample of urine is considered clinically significant. However in symptomatic patients, a smaller number of bacteria (100 to 10000/mL) may signify infection.
2. Colony count of 100 to 10000/ mL indicate infection, if isolate from specimen obtained by suprapubic aspiration or "in-and-out" catheterization or from patients with indwelling catheters.

###### SUSCEPTIBILITY:

1. A test interpreted as **SENSITIVE** implies that infection due to isolate may be appropriately treated with the dosage of an antimicrobial agent recommended for that type of infection and infecting species, unless otherwise indicated.
2. A test interpreted as **INTERMEDIATE** implies that the "infection due to the isolate may be appropriately treated in body sites where the drugs are physiologically concentrated or when a high dosage of drug can be used".
3. A test interpreted as **RESISTANT** implies that the "isolates are not inhibited by the usually achievable concentration of the agents with normal dosage, schedule and/or fall in the range where specific microbial resistance mechanism are likely (e.g. beta-lactamases), and clinical efficacy has not been reliable in treatment studies.

###### CAUTION:

Conditions which can cause a false Negative culture:

1. Patient is on antibiotics. Please repeat culture post therapy.
2. Anaerobic bacterial infection.
3. Fastidious aerobic bacteria which are not able to grow on routine culture media.
4. Besides all these factors, at least in 25-40 % of cases there is no direct correlation between in vivo clinical picture.
5. Renal tuberculosis to be confirmed by AFB studies.

\*\*\* End Of Report \*\*\*



DR. VINAY CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR. YUGAM CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY)



TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT.