

(A Unit of KOS Healthcare)



Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mr. SONA OBEROI

AGE/ GENDER : 35 YRS/MALE PATIENT ID : 1769376

COLLECTED BY : REG. NO./LAB NO. : 012502250006

 REFERRED BY
 : 25/Feb/2025 08:07 AM

 BARCODE NO.
 : 01526100
 COLLECTION DATE
 : 25/Feb/2025 08:08AM

 CLIENT CODE.
 : KOS DIAGNOSTIC LAB
 REPORTING DATE
 : 25/Feb/2025 09:55AM

**CLIENT ADDRESS**: 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit Biological Reference interval

### SWASTHYA WELLNESS PANEL: 1.5 COMPLETE BLOOD COUNT (CBC)

#### **RED BLOOD CELLS (RBCS) COUNT AND INDICES**

HAEMOGLOBIN (HB) by CALORIMETRIC	13.5	gm/dL	12.0 - 17.0
RED BLOOD CELL (RBC) COUNT by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	4.81	Millions/cmm	3.50 - 5.00
PACKED CELL VOLUME (PCV) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	41.3	%	40.0 - 54.0
MEAN CORPUSCULAR VOLUME (MCV) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	85.8	fL	80.0 - 100.0
MEAN CORPUSCULAR HAEMOGLOBIN (MCH) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	28	pg	27.0 - 34.0
MEAN CORPUSCULAR HEMOGLOBIN CONC. (MCHC) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	32.6	g/dL	32.0 - 36.0
RED CELL DISTRIBUTION WIDTH (RDW-CV) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	13.5	%	11.00 - 16.00
RED CELL DISTRIBUTION WIDTH (RDW-SD) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	43.6	fL	35.0 - 56.0
MENTZERS INDEX by CALCULATED	17.84	RATIO	BETA THALASSEMIA TRAIT: < 13.0 IRON DEFICIENCY ANEMIA: >13.0
GREEN & KING INDEX by CALCULATED	24.02	RATIO	BETA THALASSEMIA TRAIT:<= 65.0 IRON DEFICIENCY ANEMIA: > 65.0
WHITE BLOOD CELLS (WBCS)			
TOTAL LEUCOCYTE COUNT (TLC) by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	6830	/cmm	4000 - 11000
NUCLEATED RED BLOOD CELLS (nRBCS) by AUTOMATED 6 PART HEMATOLOGY ANALYZER	NIL		0.00 - 20.00
NUCLEATED RED BLOOD CELLS (nRBCS) %	NIL	%	< 10 %



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY)



by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER



(A Unit of KOS Healthcare)



Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra
MD (Pathology)
CEO & Consultant Pathologist

NAME : Mr. SONA OBEROI

**AGE/ GENDER** : 35 YRS/MALE **PATIENT ID** : 1769376

COLLECTED BY : REG. NO./LAB NO. : 012502250006

 REFERRED BY
 : 25/Feb/2025 08:07 AM

 BARCODE NO.
 : 01526100
 COLLECTION DATE
 : 25/Feb/2025 08:08 AM

 CLIENT CODE.
 : KOS DIAGNOSTIC LAB
 REPORTING DATE
 : 25/Feb/2025 09:55 AM

**CLIENT ADDRESS**: 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name	Value	Unit	Biological Reference interval		
DIFFERENTIAL LEUCOCYTE COUNT (DLC)					
NEUTROPHILS	46 <sup>L</sup>	%	50 - 70		
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY LYMPHOCYTES by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	43 <sup>H</sup>	%	20 - 40		
EOSINOPHILS	5	%	1 - 6		
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY  MONOCYTES by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	6	%	2 - 12		
BASOPHILS	0	%	0 - 1		
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY  ABSOLUTE LEUKOCYTES (WBC) COUNT					
ABSOLUTE NEUTROPHIL COUNT by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	3142	/cmm	2000 - 7500		
ABSOLUTE LYMPHOCYTE COUNT by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	2937	/cmm	800 - 4900		
ABSOLUTE EOSINOPHIL COUNT by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	342	/cmm	40 - 440		
ABSOLUTE MONOCYTE COUNT by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	410	/cmm	80 - 880		
ABSOLUTE BASOPHIL COUNT by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	0	/cmm	0 - 110		
PLATELETS AND OTHER PLATELET PREDICTIVE	MARKERS.				
PLATELET COUNT (PLT) by hydro dynamic focusing, electrical impedence	162000	/cmm	150000 - 450000		
PLATELETCRIT (PCT) by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	0.22	%	0.10 - 0.36		
MEAN PLATELET VOLUME (MPV) by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	14 <sup>H</sup>	fL	6.50 - 12.0		
PLATELET LARGE CELL COUNT (P-LCC) by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	88000	/cmm	30000 - 90000		
PLATELET LARGE CELL RATIO (P-LCR) by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	54.5 <sup>H</sup>	%	11.0 - 45.0		
PLATELET DISTRIBUTION WIDTH (PDW) by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE NOTE: TEST CONDUCTED ON EDTA WHOLE BLOOD	16.3	%	15.0 - 17.0		



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)





# KOS Diagnostic Lab (A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

**NAME** : Mr. SONA OBEROI

**AGE/ GENDER PATIENT ID** :1769376 : 35 YRS/MALE

**COLLECTED BY** : 012502250006 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 25/Feb/2025 08:07 AM **COLLECTION DATE** BARCODE NO. :01526100 : 25/Feb/2025 08:08AM REPORTING DATE CLIENT CODE. : KOS DIAGNOSTIC LAB : 25/Feb/2025 09:55AM

**CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

**Test Name Value** Unit **Biological Reference interval** 



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt -133 001, Haryana 0171-2643898, +91 99910 43898 | care@koshealthcare.com | www.koshealthcare.com







Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mr. SONA OBEROI

AGE/ GENDER : 35 YRS/MALE PATIENT ID : 1769376

COLLECTED BY : REG. NO./LAB NO. : 012502250006

 REFERRED BY
 : 25/Feb/2025 08:07 AM

 BARCODE NO.
 : 01526100
 COLLECTION DATE
 : 25/Feb/2025 08:08AM

 CLIENT CODE.
 : KOS DIAGNOSTIC LAB
 REPORTING DATE
 : 25/Feb/2025 04:41PM

CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit Biological Reference interval

#### **GLYCOSYLATED HAEMOGLOBIN (HBA1C)**

GLYCOSYLATED HAEMOGLOBIN (HbA1c): 5.2 % 4.0 - 6.4

WHOLE BLOOD

by HPLC (HIGH PERFORMANCE LIQUID CHROMATOGRAPHY)

ESTIMATED AVERAGE PLASMA GLUCOSE 102.54 mg/dL 60.00 - 140.00

by HPLC (HIGH PERFORMANCE LIQUID CHROMATOGRAPHY)

#### **INTERPRETATION:**

AS PER AMERICAN DIABETES ASSOCIATION (ADA):				
REFERENCE GROUP	GLYCOSYLATED HEMOGL	OGIB (HBAIC) in %		
Non diabetic Adults >= 18 years	<5.7			
At Risk (Prediabetes)	5.7 – 6.	4		
Diagnosing Diabetes	>= 6.5			
	Age > 19 Years			
	Goals of Therapy:	< 7.0		
Therapeutic goals for glycemic control	Actions Suggested:	>8.0		
	Age < 19 Y	ears		
	Goal of therapy:	<7.5		

#### COMMENTS:

- 1.Glycosylated hemoglobin (HbA1c) test is three monthly monitoring done to assess compliace with therapeutic regimen in diabetic patients. 2.Since Hb1c reflects long term fluctuations in blood glucose concentration, a diabetic patient who has recently under good control may still have high concentration of HbAlc. Converse is true for a diabetic previously under good control but now poorly controlled.
- 3. Target goals of < 7.0 % may be beneficial in patients with short duration of diabetes, long life expectancy and no significant cardiovascular disease. In patients with significant complications of diabetes, limited life expectancy or extensive co-morbid conditions, targetting a goal of < 7.0% may not be appropriate.
- 4.High HbA1c (>9.0 -9.5 %) is strongly associated with risk of development and rapid progression of microvascular and nerve complications 5.Any condition that shorten RBC life span like acute blood loss, hemolytic anemia falsely lower HbA1c results.
- 6.HbA1c results from patients with HbSS,HbSC and HbD must be interpreted with caution, given the pathological processes including anemia,increased red cell turnover, and transfusion requirement that adversely impact HbA1c as a marker of long-term gycemic control.

7. Specimens from patients with polycythemia or post-splenctomy may exhibit increse in HbA1c values due to a somewhat longer life span of the red cells.



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)





### **KOS Diagnostic Lab** (A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

mm/1st hr

**NAME** : Mr. SONA OBEROI

**AGE/ GENDER** : 35 YRS/MALE **PATIENT ID** : 1769376

**COLLECTED BY** REG. NO./LAB NO. :012502250006

REFERRED BY **REGISTRATION DATE** : 25/Feb/2025 08:07 AM BARCODE NO. :01526100 **COLLECTION DATE** : 25/Feb/2025 08:08AM CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 25/Feb/2025 11:25AM

**CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

**Value** Unit **Biological Reference interval Test Name** 

#### **ERYTHROCYTE SEDIMENTATION RATE (ESR)**

ERYTHROCYTE SEDIMENTATION RATE (ESR)

by RED CELL AGGREGATION BY CAPILLARY PHOTOMETRY

#### INTERPRETATION:

- 1. ESR is a non-specific test because an elevated result often indicates the presence of inflammation associated with infection, cancer and auto-immune disease, but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it.

  2. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other test such
- as C-reactive protein
- 3. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as some others, such as systemic lupus erythematosus
  CONDITION WITH LOW ESR

A low ESR can be seen with conditions that inhibit the normal sedimentation of red blood cells, such as a high red blood cell count (polycythaemia), significantly high white blood cell count (leucocytosis), and some protein abnormalities. Some changes in red cell shape (such as sickle cells in sickle cell anaemia) also lower the ESR.

NOTE:

- ESR and C reactive protein (C-RP) are both markers of inflammation.
   Generally, ESR does not change as rapidly as does CRP, either at the start of inflammation or as it resolves.
   CRP is not affected by as many other factors as is ESR, making it a better marker of inflammation.
   If the ESR is elevated, it is typically a result of two types of proteins, globulins or fibrinogen.
   Women tend to have a higher ESR, and menstruation and pregnancy can cause temporary elevations.
   Progs such as doubtern mathyldona, oral contracentives, popicillamino procesingmide, the only viling, and vitality in the orange of the contracentives.

- 6. Drugs such as dextran, methyldopa, oral contraceptives, penicillamine procainamide, theophylline, and vitamin A can increase ESR, while aspirin, cortisone, and quinine may decrease it



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)





(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

**NAME** : Mr. SONA OBEROI

**AGE/ GENDER** : 35 YRS/MALE **PATIENT ID** : 1769376

**COLLECTED BY** : 012502250006 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 25/Feb/2025 08:07 AM BARCODE NO. :01526100 **COLLECTION DATE** : 25/Feb/2025 08:08AM CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 25/Feb/2025 10:28AM

**CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

**Value** Unit **Biological Reference interval Test Name** 

### **CLINICAL CHEMISTRY/BIOCHEMISTRY GLUCOSE FASTING (F)**

GLUCOSE FASTING (F): PLASMA 87.9 NORMAL: < 100.0 mg/dL

by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD) PREDIABETIC: 100.0 - 125.0

DIABETIC: > 0R = 126.0

INTERPRETATION
IN ACCORDANCE WITH AMERICAN DIABETES ASSOCIATION GUIDELINES:

1. A fasting plasma glucose level below 100 mg/dl is considered normal.

2. A fasting plasma glucose level between 100 - 125 mg/dl is considered as glucose intolerant or prediabetic. A fasting and post-prandial blood

test (after consumption of 75 gms of glucose) is recommended for all such patients.

3. A fasting plasma glucose level of above 125 mg/dl is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for all such patients. A fasting plasma glucose level in excess of 125 mg/dl on both occasions is confirmatory for diabetic state.



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST





(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist Dr. Yugam Chopra
MD (Pathology)
CEO & Consultant Pathologist

NAME : Mr. SONA OBEROI

**AGE/ GENDER** : 35 YRS/MALE **PATIENT ID** : 1769376

COLLECTED BY : REG. NO./LAB NO. : 012502250006

 REFERRED BY
 : 25/Feb/2025 08:07 AM

 BARCODE NO.
 : 01526100
 COLLECTION DATE
 : 25/Feb/2025 08:08AM

 CLIENT CODE.
 : KOS DIAGNOSTIC LAB
 REPORTING DATE
 : 25/Feb/2025 10:49AM

**CLIENT ADDRESS**: 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name	Value	Unit	Biological Reference interval
	LIPID PROFILE	: BASIC	
CHOLESTEROL TOTAL: SERUM by CHOLESTEROL OXIDASE PAP	213.78 <sup>H</sup>	mg/dL	OPTIMAL: < 200.0 BORDERLINE HIGH: 200.0 - 239.0 HIGH CHOLESTEROL: > OR = 240.0
TRIGLYCERIDES: SERUM by GLYCEROL PHOSPHATE OXIDASE (ENZYMATIC)	323.24 <sup>H</sup>	mg/dL	OPTIMAL: < 150.0 BORDERLINE HIGH: 150.0 - 199.0 HIGH: 200.0 - 499.0 VERY HIGH: > OR = 500.0
HDL CHOLESTEROL (DIRECT): SERUM by SELECTIVE INHIBITION	44.34	mg/dL	LOW HDL: < 30.0 BORDERLINE HIGH HDL: 30.0 - 60.0 HIGH HDL: > OR = 60.0
LDL CHOLESTEROL: SERUM by CALCULATED, SPECTROPHOTOMETRY	104.79	mg/dL	OPTIMAL: < 100.0 ABOVE OPTIMAL: 100.0 - 129.0 BORDERLINE HIGH: 130.0 - 159.0 HIGH: 160.0 - 189.0 VERY HIGH: > OR = 190.0
NON HDL CHOLESTEROL: SERUM by CALCULATED, SPECTROPHOTOMETRY	169.44 <sup>H</sup>	mg/dL	OPTIMAL: < 130.0 ABOVE OPTIMAL: 130.0 - 159.0 BORDERLINE HIGH: 160.0 - 189.0 HIGH: 190.0 - 219.0 VERY HIGH: > OR = 220.0
VLDL CHOLESTEROL: SERUM by CALCULATED, SPECTROPHOTOMETRY	64.65 <sup>H</sup>	mg/dL	0.00 - 45.00
TOTAL LIPIDS: SERUM by CALCULATED, SPECTROPHOTOMETRY	750.8 <sup>H</sup>	mg/dL	350.00 - 700.00
CHOLESTEROL/HDL RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	4.82 <sup>H</sup>	RATIO	LOW RISK: 3.30 - 4.40 AVERAGE RISK: 4.50 - 7.0 MODERATE RISK: 7.10 - 11.0 HIGH RISK: > 11.0



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)





(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

**NAME** : Mr. SONA OBEROI

AGE/ GENDER : 35 YRS/MALE **PATIENT ID** : 1769376

**COLLECTED BY** : 012502250006 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 25/Feb/2025 08:07 AM BARCODE NO. :01526100 **COLLECTION DATE** : 25/Feb/2025 08:08AM CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 25/Feb/2025 10:49AM

**CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name	Value	Unit	Biological Reference interval
LDL/HDL RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	2.36	RATIO	LOW RISK: 0.50 - 3.0 MODERATE RISK: 3.10 - 6.0 HIGH RISK: > 6.0
TRIGLYCERIDES/HDL RATIO: SERUM by CALCULATED. SPECTROPHOTOMETRY	7.29 <sup>H</sup>	RATIO	3.00 - 5.00

#### **INTERPRETATION:**

1. Measurements in the same patient can show physiological analytical variations. Three serial samples 1 week apart are recommended for Total Cholesterol, Triglycerides, HDL & LDL Cholesterol.

2. As per NLA-2014 guidelines, all adults above the age of 20 years should be screened for lipid status. Selective screening of children above the age of 2 years with a family history of premature cardiovascular disease or those with at least one parent with high total cholesterol is recommended.

3. Low HDL levels are associated with increased risk for Atherosclerotic Cardiovascular disease (ASCVD) due to insufficient HDL being available

to participate in reverse cholesterol transport, the process by which cholesterol is eliminated from peripheral tissues.

4. NLA-2014 identifies Non HDL Cholesterol (an indicator of all atherogeniclipoproteins such as LDL, VLDL, IDL, Lpa, Chylomicron remnants) along with LDL-cholesterol as co- primary target for cholesterol lowering therapy. Note that major risk factors can modify treatment goals for LDL &Non

5. Additional testing for Apolipoprotein B, hsCRP,Lp(a) & LP-PLA2 should be considered among patients with moderate risk for ASCVD for risk refinement



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)





(A Unit of KOS Healthcare)



Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mr. SONA OBEROI

AGE/ GENDER : 35 YRS/MALE PATIENT ID : 1769376

COLLECTED BY : REG. NO./LAB NO. : 012502250006

 REFERRED BY
 : 25/Feb/2025 08:07 AM

 BARCODE NO.
 : 01526100
 COLLECTION DATE
 : 25/Feb/2025 08:08AM

 CLIENT CODE.
 : KOS DIAGNOSTIC LAB
 REPORTING DATE
 : 25/Feb/2025 10:49AM

**CLIENT ADDRESS**: 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit Biological Reference interval

#### **LIVER FUNCTION TEST (COMPLETE)**

BILIRUBIN TOTAL: SERUM by DIAZOTIZATION, SPECTROPHOTOMETRY	1.12	mg/dL	INFANT: 0.20 - 8.00 ADULT: 0.00 - 1.20
BILIRUBIN DIRECT (CONJUGATED): SERUM by DIAZO MODIFIED, SPECTROPHOTOMETRY	0.17	mg/dL	0.00 - 0.40
BILIRUBIN INDIRECT (UNCONJUGATED): SERUM by CALCULATED, SPECTROPHOTOMETRY	0.95	mg/dL	0.10 - 1.00
SGOT/AST: SERUM by IFCC, WITHOUT PYRIDOXAL PHOSPHATE	21.8	U/L	7.00 - 45.00
SGPT/ALT: SERUM by IFCC, WITHOUT PYRIDOXAL PHOSPHATE	22.2	U/L	0.00 - 49.00
AST/ALT RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	0.98	RATIO	0.00 - 46.00
ALKALINE PHOSPHATASE: SERUM by Para nitrophenyl phosphatase by amino methyl propanol	73.58	U/L	40.0 - 130.0
GAMMA GLUTAMYL TRANSFERASE (GGT): SERUN by SZASZ, SPECTROPHTOMETRY	1 20.86	U/L	0.00 - 55.0
TOTAL PROTEINS: SERUM by BIURET, SPECTROPHOTOMETRY	6.86	gm/dL	6.20 - 8.00
ALBUMIN: SERUM by BROMOCRESOL GREEN	4.38	gm/dL	3.50 - 5.50
GLOBULIN: SERUM by CALCULATED, SPECTROPHOTOMETRY	2.48	gm/dL	2.30 - 3.50
A: GRATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	1.77	RATIO	1.00 - 2.00

#### INTERPRETATION

*NOTE*:- To be correlated in individuals having SGOT and SGPT values higher than Normal Referance Range.

**USE**:- Differential diagnosis of diseases of hepatobiliary system and pancreas.

#### INCREASED:

DRUG HEPATOTOXICITY	> 2
ALCOHOLIC HEPATITIS	> 2 (Highly Suggestive)
CIRRHOSIS	1.4 - 2.0
INTRAHEPATIC CHOLESTATIS	> 1.5
HEPATOCELLULAR CARCINOMA & CHRONIC HEPATITIS	> 1.3 (Slightly Increased)



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)





(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mr. SONA OBEROI

**AGE/ GENDER** : 35 YRS/MALE **PATIENT ID** : 1769376

COLLECTED BY : REG. NO./LAB NO. : 012502250006

 REFERRED BY
 : 25/Feb/2025 08:07 AM

 BARCODE NO.
 : 01526100
 COLLECTION DATE
 : 25/Feb/2025 08:08AM

 CLIENT CODE.
 : KOS DIAGNOSTIC LAB
 REPORTING DATE
 : 25/Feb/2025 10:49AM

**CLIENT ADDRESS**: 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit Biological Reference interval

#### **DECREASED:**

1. Acute Hepatitis due to virus, drugs, toxins (with AST increased 3 to 10 times upper limit of normal)

2. Extra Hepatic cholestatis: 0.8 (normal or slightly decreased).

#### PROGNOSTIC SIGNIFICANCE:

NORMAL	< 0.65	
GOOD PROGNOSTIC SIGN	0.3 - 0.6	
POOR PROGNOSTIC SIGN	1.2 - 1.6	



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)





(A Unit of KOS Healthcare)



Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mr. SONA OBEROI

AGE/ GENDER : 35 YRS/MALE PATIENT ID : 1769376

COLLECTED BY : REG. NO./LAB NO. : 012502250006

 REFERRED BY
 : 25/Feb/2025 08:07 AM

 BARCODE NO.
 : 01526100
 COLLECTION DATE
 : 25/Feb/2025 08:08AM

 CLIENT CODE.
 : KOS DIAGNOSTIC LAB
 REPORTING DATE
 : 25/Feb/2025 10:49AM

**CLIENT ADDRESS**: 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name	Value	Unit	Biological Reference interval
KIDNI	EY FUNCTION T	EST (COMPLETE)	
UREA: SERUM by urease - glutamate dehydrogenase (gldh)	18.16	mg/dL	10.00 - 50.00
CREATININE: SERUM by ENZYMATIC, SPECTROPHOTOMETERY	0.95	mg/dL	0.40 - 1.40
BLOOD UREA NITROGEN (BUN): SERUM by CALCULATED, SPECTROPHOTOMETRY	8.49	mg/dL	7.0 - 25.0
BLOOD UREA NITROGEN (BUN)/CREATININE RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	8.94 <sup>L</sup>	RATIO	10.0 - 20.0
UREA/CREATININE RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	19.12	RATIO	
URIC ACID: SERUM by URICASE - OXIDASE PEROXIDASE	3.98	mg/dL	3.60 - 7.70
CALCIUM: SERUM by ARSENAZO III, SPECTROPHOTOMETRY	8.44 <sup>L</sup>	mg/dL	8.50 - 10.60
PHOSPHOROUS: SERUM by PHOSPHOMOLYBDATE, SPECTROPHOTOMETRY	4.24	mg/dL	2.30 - 4.70
ELECTROLYTES			
SODIUM: SERUM by ISE (ION SELECTIVE ELECTRODE)	139.7	mmol/L	135.0 - 150.0
POTASSIUM: SERUM by ISE (ION SELECTIVE ELECTRODE)	4.21	mmol/L	3.50 - 5.00
CHLORIDE: SERUM by ISE (ION SELECTIVE ELECTRODE)	104.78	mmol/L	90.0 - 110.0
ESTIMATED GLOMERULAR FILTERATION RATE			

ESTIMATED GLOMERULAR FILTERATION RATE 107

(eGFR): SERUM by CALCULATED INTERPRETATION:

To differentiate between pre- and post renal azotemia.

### INCREASED RATIO (>20:1) WITH NORMAL CREATININE:

- 1. Prerenal azotemia (BUN rises without increase in creatinine) e.g. heart failure, salt depletion, dehydration, blood loss) due to decreased glomerular filtration rate.
- 2. Catabolic states with increased tissue breakdown.
- 3. GI haemorrhage.



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)









Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

**NAME** : Mr. SONA OBEROI

AGE/ GENDER : 35 YRS/MALE **PATIENT ID** : 1769376

**COLLECTED BY** : 012502250006 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 25/Feb/2025 08:07 AM BARCODE NO. :01526100 **COLLECTION DATE** : 25/Feb/2025 08:08AM CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 25/Feb/2025 10:49AM

**CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

**Test Name** Value Unit **Biological Reference interval** 

4. High protein intake.

5. Impaired renal function plus

6. Excess protein intake or production or tissue breakdown (e.g. infection, GI bleeding, thyrotoxicosis, Cushing's syndrome, high protein diet, burns, surgery, cachexia, high fever).

7. Urine reabsorption (e.g. ureter colostomy)

8. Reduced muscle mass (subnormal creatinine production)

9. Certain drugs (e.g. tetracycline, glucocorticoids)

#### INCREASED RATIO (>20:1) WITH ELEVATED CREATININE LEVELS:

- 1. Postrenal azotemia (BUN rises disproportionately more than creatinine) (e.g. obstructive uropathy).
- 2. Prerenal azotemia superimposed on renal disease.

#### DECREASED RATIO (<10:1) WITH DECREASED BUN:

- 1. Acute tubular necrosis.
- 2. Low protein diet and starvation.
- 3. Severe liver disease.
- 4. Other causes of decreased urea synthesis.
- 5. Repeated dialysis (urea rather than creatinine diffuses out of extracellular fluid).
- 6. Inherited hyperammonemias (urea is virtually absent in blood).
- 7. SIADH (syndrome of inappropiate antidiuretic harmone) due to tubular secretion of urea.
- 8. Pregnancy.

#### **DECREASED RATIO (<10:1) WITH INCREASED CREATININE:**

- 1. Phenacimide therapy (accelerates conversion of creatine to creatinine).
- 2. Rhabdomyolysis (releases muscle creatinine).
- 3. Muscular patients who develop renal failure.

#### **INAPPROPIATE RATIO:**

1. Diabetic ketoacidosis (acetoacetate causes false increase in creatinine with certain methodologies, resulting in normal ratio when dehydration should produce an increased BUN/creatinine ratio).

2. Cephalosporin therapy (interferes with creatinine measurement). **ESTIMATED GLOMERULAR FILTERATION RATE**:

CONTROL OF CONTROL OF TAXABLE CO				
CKD STAGE	DESCRIPTION	GFR ( mL/min/1.73m2 )	ASSOCIATED FINDINGS	
G1	Normal kidney function	>90	No proteinuria	
G2	Kidney damage with normal or high GFR	>90	Presence of Protein , Albumin or cast in urine	
G3a	Mild decrease in GFR	60 -89		
G3b	Moderate decrease in GFR	30-59		
G4	Severe decrease in GFR	15-29		
G5	Kidney failure	<15		



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)

KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana



(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

**NAME** : Mr. SONA OBEROI

AGE/ GENDER : 35 YRS/MALE **PATIENT ID** : 1769376

COLLECTED BY REG. NO./LAB NO. :012502250006

REFERRED BY **REGISTRATION DATE** : 25/Feb/2025 08:07 AM BARCODE NO. :01526100 **COLLECTION DATE** : 25/Feb/2025 08:08AM CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 25/Feb/2025 10:49AM

**CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

**Test Name** Value Unit **Biological Reference interval** 

#### COMMENTS:

1. Estimated Glomerular filtration rate (eGFR) is the sum of filtration rates in all functioning nephrons and so an estimation of the GFR provides a measure of functioning nephrons of the kidney.

2. eGFR calculated using the 2009 CKD-EPI creatinine equation and GFR category reported as per KDIGO guideline 2012

3. In patients, with eGFR creating between 45-59 ml/min/1.73 m2 (G3) and without any marker of Kidney damage, It is recommended to measure

4. eGFR category G1 OR G2 does not fullfill the criteria for CKD, in the absence of evidence of Kidney Damage
5. In a suspected case of Acute Kidney Injury (AKI), measurement of eGFR should be done after 48-96 hours of any Intervention or procedure
6. eGFR calculated by Serum Creatinine may be less accurate due to certain factors like Race, Muscle Mass, Diet, Certain Drugs. In such cases, eGFR should be calculated using Serum Cystatin C
7. A decrease in eGFR implies either progressive renal disease, or a reversible process causing decreased nephron function (eg, severe dehydration).

KDIGO guideline, 2012 recommends Chronic Kidney Disease (CKD) should be classified based on cause, eGFR category and Albuminuria (ACR) category. GFR & ACR category combined together reflect risk of progression and helps Clinician to identify the individual who are progressing at more rapid rate than anticipated



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt - 133 001, Haryana



(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

**NAME** : Mr. SONA OBEROI

**AGE/ GENDER** : 35 YRS/MALE **PATIENT ID** : 1769376

**COLLECTED BY** : 012502250006 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 25/Feb/2025 08:07 AM BARCODE NO. :01526100 **COLLECTION DATE** : 25/Feb/2025 08:08AM CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 25/Feb/2025 10:49AM

**CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name	Value	Unit	Biological Reference interval
	IRON PRO	FILE	
IRON: SERUM by FERROZINE, SPECTROPHOTOMETRY	56.82 <sup>L</sup>	μg/dL	59.0 - 158.0
UNSATURATED IRON BINDING CAPACITY (UIBC) :SERUM by FERROZINE, SPECTROPHOTOMETERY	268.32	μg/dL	150.0 - 336.0
TOTAL IRON BINDING CAPACITY (TIBC) :SERUM by SPECTROPHOTOMETERY	325.14	μg/dL	230 - 430
%TRANSFERRIN SATURATION: SERUM by CALCULATED, SPECTROPHOTOMETERY (FERENE)	17.48	%	15.0 - 50.0
TRANSFERRIN: SERUM by SPECTROPHOTOMETERY (FERENE)	230.85	mg/dL	200.0 - 350.0

#### INTERDRETATION.

WIERFRETATION.				
VARIABLES	ANEMIA OF CHRONIC DISEASE	IRON DEFICIENCY ANEMIA	THALASSEMIA α/β TRAIT	
SERUM IRON:	Normal to Reduced	Reduced	Normal	
TOTAL IRON BINDING CAPACITY:	Decreased	Increased	Normal	
% TRANSFERRIN SATURATION:	Decreased	Decreased < 12-15 %	Normal	
SERUM FERRITIN:	Normal to Increased	Decreased	Normal or Increased	

#### IRON:

1. Serum iron studies is recommended for differential diagnosis of microcytic hypochromic anemia.i.e iron deficiency anemia, zinc deficiency anemia, anemia of chronic disease and thalassemia syndromes.

2. It is essential to isolate iron deficiency anemia from Beta thalassemia syndromes because during iron replacement which is therapeutic for iron deficiency anemia, is severely contra-indicated in Thalassemia.

TOTAL IRON BINDING CAPACITY (TIBC):

1. It is a direct meaning in the bone marrow.

% TRANSFERRIN SATURATION:

1.Occurs in idiopathic hemochromatosis and transfusional hemosiderosis where no unsaturated iron binding capacity is available for iron mobilization. Similar condition is seen in congenital deficiency of transferrin.



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana



(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

**NAME** : Mr. SONA OBEROI

**AGE/ GENDER** : 35 YRS/MALE **PATIENT ID** : 1769376

**COLLECTED BY** REG. NO./LAB NO. :012502250006

REFERRED BY **REGISTRATION DATE** : 25/Feb/2025 08:07 AM BARCODE NO. :01526100 **COLLECTION DATE** : 25/Feb/2025 08:08AM CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 25/Feb/2025 10:49AM

**CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

**Value** Unit **Biological Reference interval Test Name** 

#### **FERRITIN**

FERRITIN: SERUM 183.18 ng/mL 21.81 - 274.66

by CLIA (CHEMILUMINESCENCE IMMUNOASSAY)

Serum ferritin appears to be in equilibrium with tissue ferritin and is a good indicator of storage iron in normal subjects and in most disorders. In patients with some hepatocellular diseases, malignancies and inflammatory diseases, serum ferritin is a disproportionately high estimate of storage iron because serum ferritin is an acute phase reactant. In such disorders iron deficiency anemia may exist with a normal serum ferritin concentration. In the presence of inflammation, persons with low serum ferritin are likely to respond to iron therapy.

#### **DECREASED:**

- 1. Iron depletion appears to be the only condition associated with reduced serum ferritin concentrations.
- 2. Hypothyroidism.3. Vitamin-C deficiency

### INCREASED FERRITIN DUE TO IRON OVERLOAD (PRIMARY):

- 1. Hemochromatosis or hemosiderosis.
- Wilson Disease

### INCREASED FERRITIN DUE TO IRON OVERLOAD (SECONDARY):

- 1. Transfusion overload
- Excess dietary Iron
   Porphyria Cutanea tada

### 4. Ineffective erythropolesis. INCREASED FERRITIN WITHOUT IRON OVERLOAD:

- 1. Liver disorders (NASH) or viral hepatitis (B/C)
- 2. Inflammatory conditions (Ferritin is a acute phase reactant) both acute and chronic.
- 3. Leukaemia, hodgkin's disease.
- 4. Alcohol excess.
- 5. Other malignancies in which increases probably reflect the escape of ferritin from damaged liver cells, impaired clearance from the plasma, synthesis of ferritin by tumour cells.
- 6. Ferritin levels below 10 ng/ml have been reported as indicative of iron deficiency anemia.

#### NOTE:

1. As Ferritin is an acute phase reactant, it is often raised in both acute and chronic inflammatory condition of the body such as infections leading to false positive results. It can thererfore mask a diagnostically low result. In such Cases serum ferritin levels should always be correlated with C-Reactive proteins to rule out any inflammatory conditions.

2. Patients with iron deficiency anaemia may occasionally have elevated or normal ferritin levels. This is usually seen in patients already receiving iron therapy or in patients with concomitant hepatocellular injury.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana



(A Unit of KOS Healthcare)



Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mr. SONA OBEROI

AGE/ GENDER : 35 YRS/MALE PATIENT ID : 1769376

COLLECTED BY : REG. NO./LAB NO. : 012502250006

 REFERRED BY
 : 25/Feb/2025 08:07 AM

 BARCODE NO.
 : 01526100
 COLLECTION DATE
 : 25/Feb/2025 08:08AM

 CLIENT CODE.
 : KOS DIAGNOSTIC LAB
 REPORTING DATE
 : 25/Feb/2025 10:49AM

CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit Biological Reference interval

### **ENDOCRINOLOGY**

### THYROID FUNCTION TEST: TOTAL

TRIIODOTHYRONINE (T3): SERUM 0.706 ng/mL 0.35 - 1.93

by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

THYROXINE (T4): SERUM 6.11 μgm/dL 4.87 - 12.60

by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

THYROID STIMULATING HORMONE (TSH): SERUM 3.668  $\mu$ IU/mL 0.35 - 5.50

by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

3rd GENERATION, ULTRASENSITIVE

#### **INTERPRETATION:**

TSH levels are subject to circadian variation, reaching peak levels between 2-4 a.m and at a minimum between 6-10 pm. The variation is of the order of 50%. Hence time of the day has influence on the measured serum TSH concentrations. TSH stimulates the production and secretion of the metabolically active hormones, thyroxine (T4) and triiodothyronine (T3). Failure at any level of regulation of the hypothalamic-pituitary-thyroid axis will result in either underproduction (hypothyroidism) or overproduction (hyperthyroidism) of T4 and/or T3.

CLINICAL CONDITION	Т3	T4	TSH
Primary Hypothyroidism:	Reduced	Reduced	Increased (Significantly)
Subclinical Hypothyroidism:	Normal or Low Normal	Normal or Low Normal	High
Primary Hyperthyroidism:	Increased	Increased	Reduced (at times undetectable)
Subclinical Hyperthyroidism:	Normal or High Normal	Normal or High Normal	Reduced

#### LIMITATIONS:-

- 1. T3 and T4 circulates in reversibly bound form with Thyroid binding globulins (TBG), and to a lesser extent albumin and Thyroid binding Pre Albumin so conditions in which TBG and protein levels alter such as pregnancy, excess estrogens, anabolic steroids and glucocorticoids may falsely affect the T3 and T4 levels and may cause false thyroid values for thyroid function tests.
- 2. Normal levels of T4 can also be seen in Hyperthyroid patients with :T3 Thyrotoxicosis, Decreased binding capacity due to hypoproteinemia or ingestion of certain drugs (e.g.: phenytoin, salicylates)
- 3. Serum T4 levels in neonates and infants are higher than values in the normal adult, due to the increased concentration of TBG in neonate serum.
- 4. TSH may be normal in central hypothyroidism, recent rapid correction of hyperthyroidism or hypothyroidism, pregnancy, phenytoin therapy.

TRIIODOTHYRONINE (T3)		THYROXINE (T4)		THYROID STIMULATING HORMONE (TSH)	
Age	Refferance Range (ng/mL)	Age	Refferance Range (µg/dL)	Age	Reference Range ( μIU/mL)
0 - 7 Days	0.20 - 2.65	0 - 7 Days	5.90 - 18.58	0 - 7 Days	2.43 - 24.3
7 Days - 3 Months	0.36 - 2.59	7 Days - 3 Months	6.39 - 17.66	7 Days - 3 Months	0.58 - 11.00
3 - 6 Months	0.51 - 2.52	3 - 6 Months	6.75 – 17.04	3 Days – 6 Months	0.70 - 8.40
6 - 12 Months	0.74 - 2.40	6 - 12 Months	7.10 - 16.16	6 – 12 Months	0.70 - 7.00



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana



(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mr. SONA OBEROI

AGE/ GENDER : 35 YRS/MALE PATIENT ID : 1769376

COLLECTED BY : REG. NO./LAB NO. : 012502250006

 REFERRED BY
 : 25/Feb/2025 08:07 AM

 BARCODE NO.
 : 01526100
 COLLECTION DATE
 : 25/Feb/2025 08:08AM

 CLIENT CODE.
 : KOS DIAGNOSTIC LAB
 REPORTING DATE
 : 25/Feb/2025 10:49AM

**CLIENT ADDRESS**: 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name			Value	Unit		Biologi	cal Reference interval
1 - 10 Years	0.92 - 2.28	1 - 10 Years	6.00 - 13.80	1 – 10 Years	0.60 - 5.50		
11- 19 Years	0.35 - 1.93	11 - 19 Years	4.87- 13.20	11 – 19 Years	0.50 - 5.50		
> 20 years (Adults)	0.35 - 1.93	> 20 Years (Adults)	4.87 - 12.60	> 20 Years (Adults)	0.35-5.50		
	RECOMMENDATIONS OF TSH LEVELS DURING PREGNANCY ( µIU/mL)						
	1st Trimester		0.10 - 2.50				
	2nd Trimester		0.20 - 3.00				
	3rd Trimester		0.30 – 4.10				

#### **INCREASED TSH LEVELS:**

- 1. Primary or untreated hypothyroidism may vary from 3 times to more than 100 times normal depending upon degree of hypofunction.
- 2. Hypothyroid patients receiving insufficient thyroid replacement therapy.
- 3. Hashimotos thyroiditis
- 4.DRUGS: Amphetamines, iodine containing agents & dopamine antagonist.
- 5. Neonatal period, increase in 1st 2-3 days of life due to post-natal surge

#### **DECREASED TSH LEVELS:**

- 1.Toxic multi-nodular goiter & Thyroiditis.
- 2. Over replacement of thyroid hormone in treatment of hypothyroidism.
- 3. Autonomously functioning Thyroid adenoma
- 4. Secondary pituitary or hypothalamic hypothyroidism
- 5. Acute psychiatric illness
- 6. Severe dehydration.
- 7.DRUGS: Glucocorticoids, Dopamine, Levodopa, T4 replacement therapy, Anti-thyroid drugs for thyrotoxicosis.

8. Pregnancy: 1st and 2nd Trimester



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana



(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

**NAME** : Mr. SONA OBEROI

**AGE/ GENDER** : 35 YRS/MALE **PATIENT ID** : 1769376

**COLLECTED BY** REG. NO./LAB NO. :012502250006

REFERRED BY **REGISTRATION DATE** : 25/Feb/2025 08:07 AM BARCODE NO. :01526100 **COLLECTION DATE** : 25/Feb/2025 08:08AM CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 25/Feb/2025 10:49AM

**CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

**Value** Unit **Biological Reference interval Test Name** 

### IMMUNOPATHOLOGY/SEROLOGY **C-REACTIVE PROTEIN (CRP)**

C-REACTIVE PROTEIN (CRP) QUANTITATIVE: 0.37 0.0 - 6.0mg/L

by NEPHLOMETRY

**INTERPRETATION:** 

C-reactive protein (CRP) is one of the most sensitive acute-phase reactants for inflammation.

2. CRP levels can increase dramatically (100-fold or more) after severe trauma, bacterial infection, inflammation, surgery, or neoplastic

3. CRP levels (Quantitative) has been used to assess activity of inflammatory disease, to detect infections after surgery, to detect transplant rejection, and to monitor these inflammatory processes.

4. As compared to ESR, CRP shows an earlier rise in inflammatory disorders which begins in 4-6 hrs, the intensity of the rise being higher than ESR and the recovery being earlier than ESR. Unlike ESR, CRP levels are not influenced by hematologic conditions like Anemia, Polycythemia etc., 5. Elevated values are consistent with an acute inflammatory process.

NOTE:

1. Elevated C-reactive protein (CRP) values are nonspecific and should not be interpreted without a complete clinical history.

2. Oral contraceptives may increase CRP levels.



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST





### **KOS Diagnostic Lab** (A Unit of KOS Healthcare)





Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

**NAME** : Mr. SONA OBEROI

**AGE/ GENDER** : 35 YRS/MALE **PATIENT ID** : 1769376

**COLLECTED BY** : 012502250006 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 25/Feb/2025 08:07 AM BARCODE NO. :01526100 **COLLECTION DATE** : 25/Feb/2025 08:08AM CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 25/Feb/2025 11:46AM

**CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

**Value** Unit **Biological Reference interval Test Name** 

### **VITAMINS**

### VITAMIN D/25 HYDROXY VITAMIN D3

VITAMIN D (25-HYDROXY VITAMIN D3): SERUM ng/mL DEFICIENCY: < 20.0  $29.2^{L}$ 

by CLIA (CHEMILUMINESCENCE IMMUNOASSAY) INSUFFICIENCY: 20.0 - 30.0 SUFFICIENCY: 30.0 - 100.0

**TOXICITY:** > 100.0

**INTERPRETATION:** 

DEFICIENT:	< 20	ng/mL
INSUFFICIENT:	21 - 29	ng/mL
PREFFERED RANGE:	30 - 100	ng/mL
INTOXICATION:	> 100	ng/mL

- 1. Vitamin D compounds are derived from dietary ergocalciferol (from plants, Vitamin D2), or cholecalciferol (from animals, Vitamin D3), or by conversion of 7- dihydrocholecalciferol to Vitamin D3 in the skin upon Ultraviolet exposure.

  2.25-OH--Vitamin D represents the main body resevoir and transport form of Vitamin D and transport form of Vitamin D, being stored in adipose
- tissue and tightly bound by a transport protein while in circulation.
- 3. Vitamin D plays a primary role in the maintenance of calcium homeostatis. It promotes calcium absorption, renal calcium absorption and phosphate reabsorption, skeletal calcium deposition, calcium mobilization, mainly regulated by parathyroid harmone (PTH).

  4. Severe deficiency may lead to failure to mineralize newly formed osteoid in bone, resulting in rickets in children and osteomalacia in adults.
- DECREASED:
- 1.Lack of sunshine exposure.
- 2.Inadequate intake, malabsorption (celiac disease)
- 3. Depressed Hepatic Vitamin D 25- hydroxylase activity
- 4. Secondary to advanced Liver disease
- 5. Osteoporosis and Secondary Hyperparathroidism (Mild to Moderate deficiency)
- 6.Enzyme Inducing drugs: anti-epileptic drugs like phenytoin, phenobarbital and carbamazepine, that increases Vitamin D metabolism. INCREASED:
- 1. Hypervitaminosis D is Rare, and is seen only after prolonged exposure to extremely high doses of Vitamin D. When it occurs, it can result in severe hypercalcemia and hyperphophatemia.

CAUTION: Replacement therapy in deficient individuals must be monitored by periodic assessment of Vitamin D levels in order to prevent hypervitaminosis D

NOTE:-Dark coloured individuals as compare to whites, is at higher risk of developing Vitamin D deficiency due to excess of melanin pigment which interefere with Vitamin D absorption.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST





(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mr. SONA OBEROI

AGE/ GENDER : 35 YRS/MALE PATIENT ID : 1769376

COLLECTED BY : REG. NO./LAB NO. : 012502250006

 REFERRED BY
 : 25/Feb/2025 08:07 AM

 BARCODE NO.
 : 01526100
 COLLECTION DATE
 : 25/Feb/2025 08:08AM

 CLIENT CODE.
 : KOS DIAGNOSTIC LAB
 REPORTING DATE
 : 25/Feb/2025 10:49AM

CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit Biological Reference interval

### **VITAMIN B12/COBALAMIN**

VITAMIN B12/COBALAMIN: SERUM 227 pg/mL 190.0 - 890.0

by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

#### **INTERPRETATION:-**

INCREASED VITAMIN B12	DECREASED VITAMIN B12		
1.Ingestion of Vitamin C	1.Pregnancy		
2.Ingestion of Estrogen	2.DRUGS:Aspirin, Anti-convulsants, Colchicine		
3.Ingestion of Vitamin A	3.Ethanol Igestion		
4.Hepatocellular injury	4. Contraceptive Harmones		
5.Myeloproliferative disorder	5.Haemodialysis		
6.Uremia	6. Multiple Myeloma		

- 1. Vitamin B12 (cobalamin) is necessary for hematopoiesis and normal neuronal function.
- 2.In humans, it is obtained only from animal proteins and requires intrinsic factor (IF) for absorption.
- 3. The body uses its vitamin B12 stores very economically, reabsorbing vitamin B12 from the ileum and returning it to the liver; very little is excreted.
- 4.Vitamin B12 deficiency may be due to lack of IF secretion by gastric mucosa (eg. gastrectomy, gastric atrophy) or intestinal malabsorption (eg, ileal resection, small intestinal diseases).
- 5.Vitamin B12 deficiency frequently causes macrocytic anemia, glossitis, peripheral neuropathy, weakness, hyperreflexia, ataxia, loss of proprioception, poor coordination, and affective behavioral changes. These manifestations may occur in any combination; many patients have the neurologic defects without macrocytic anemia.
- 6.Serum methylmalonic acid and homocysteine levels are also elevated in vitamin B12 deficiency states.
- 7.Follow-up testing for antibodies to intrinsic factor (IF) is recommended to identify this potential cause of vitamin B12 malabsorption.

  NOTE:A normal serum concentration of vitamin B12 does not rule out tissue deficiency of vitamin B12. The most sensitive test for vitamin B12 deficiency at the cellular level is the assay for MMA. If clinical symptoms suggest deficiency, measurement of MMA and homocysteine should be considered, even if serum vitamin B12 concentrations are normal.



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)





(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

**NAME** : Mr. SONA OBEROI

**AGE/ GENDER** : 35 YRS/MALE **PATIENT ID** : 1769376

**COLLECTED BY** : 012502250006 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 25/Feb/2025 08:07 AM BARCODE NO. :01526100 **COLLECTION DATE** : 25/Feb/2025 08:08AM CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE : 25/Feb/2025 10:08AM

**CLIENT ADDRESS** : 6349/1, NICHOLSON ROAD, AMBALA CANTT

**Value** Unit **Biological Reference interval Test Name** 

### **CLINICAL PATHOLOGY URINE ROUTINE & MICROSCOPIC EXAMINATION**

#### PHYSICAL EXAMINATION

QUANTITY RECIEVED 10 ml by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

PALE YELLOW COLOUR PALE YELLOW

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

TRANSPARANCY **CLEAR CLEAR** 

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

SPECIFIC GRAVITY 1.02 1.002 - 1.030by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

**CHEMICAL EXAMINATION** 

**ACIDIC** REACTION by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

Negative NEGATIVE (-ve)

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY **SUGAR** 

NEGATIVE (-ve) Negative by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

рН 5.5 5.0 - 7.5

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

BILIRUBIN NEGATIVE (-ve) Negative by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

**NITRITE** Negative **NEGATIVE** (-ve)

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY. EU/dL

Normal by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

NEGATIVE (-ve) KETONE BODIES Negative by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

NEGATIVE (-ve) BLOOD Negative

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

ASCORBIC ACID NEGATIVE (-ve) NEGATIVE (-ve) by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

**MICROSCOPIC EXAMINATION** 

RED BLOOD CELLS (RBCs) NEGATIVE (-ve) /HPF 0 - 3



UROBILINOGEN

CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST



0.2 - 1.0



(A Unit of KOS Healthcare)



Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mr. SONA OBEROI

**AGE/ GENDER** : 35 YRS/MALE **PATIENT ID** : 1769376

COLLECTED BY : REG. NO./LAB NO. : 012502250006

 REFERRED BY
 : 25/Feb/2025 08:07 AM

 BARCODE NO.
 : 01526100
 COLLECTION DATE
 : 25/Feb/2025 08:08 AM

 CLIENT CODE.
 : KOS DIAGNOSTIC LAB
 REPORTING DATE
 : 25/Feb/2025 10:08 AM

**CLIENT ADDRESS**: 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name	Value	Unit	Biological Reference interval
by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT			
PUS CELLS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	2-3	/HPF	0 - 5
EPITHELIAL CELLS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	1-2	/HPF	ABSENT
CRYSTALS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	NEGATIVE (-ve)		NEGATIVE (-ve)
CASTS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	NEGATIVE (-ve)		NEGATIVE (-ve)
BACTERIA by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	NEGATIVE (-ve)		NEGATIVE (-ve)
OTHERS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	NEGATIVE (-ve)		NEGATIVE (-ve)
TRICHOMONAS VAGINALIS (PROTOZOA) by microscopy on centrifuged urinary sediment	ABSENT		ABSENT

\*\*\* End Of Report \*\*\*



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana
KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt -133 001, Haryana
0171-2643898, +91 99910 43898 | care@koshealthcare.com | www.koshealthcare.com