

(A Unit of KOS Healthcare)



Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mrs. KRISHNA DEVI

AGE/ GENDER : 47 YRS/FEMALE PATIENT ID : 1779136

COLLECTED BY : REG. NO./LAB NO. : 012503050006

 REFERRED BY
 : 05/Mar/2025 07:58 AM

 BARCODE NO.
 : 01526480
 COLLECTION DATE
 : 05/Mar/2025 08:06AM

 CLIENT CODE.
 : KOS DIAGNOSTIC LAB
 REPORTING DATE
 : 05/Mar/2025 08:59AM

CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit Biological Reference interval

VIRAL MARKERS COMBO PANEL: 1.0 HEPATITIS C VIRUS (HCV) ANTIBODIES SCREENING

HEPATITIS C ANTIBODY (HCV) TOTAL

NON - REACTIVE

RESULT

by IMMUNOCHROMATOGRAPHY

INTERPRETATION:

1. Anti HCV total antibody assay identifies presence IgG antibodies in the serum. It is a useful screening test with a specificity of nearly 99%. 2. It becomes positive approximately 24 weeks after exposure. The test can not isolate an active ongoing HCV infection from an old infection that has been cleared. All positive results must be confirmed for active disease by an HCV PCR test.

FALSE NEGATIVE RESULTS SEEN IN:

1. Window period

2.Immunocompromised states.



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)





(A Unit of KOS Healthcare)



Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mrs. KRISHNA DEVI

AGE/ GENDER : 47 YRS/FEMALE PATIENT ID : 1779136

COLLECTED BY : REG. NO./LAB NO. : 012503050006

 REFERRED BY
 : 05/Mar/2025 07:58 AM

 BARCODE NO.
 : 01526480
 COLLECTION DATE
 : 05/Mar/2025 08:06AM

 CLIENT CODE.
 : KOS DIAGNOSTIC LAB
 REPORTING DATE
 : 05/Mar/2025 08:59AM

CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit Biological Reference interval

ANTI HUMAN IMMUNODEFICIENCY VIRUS (HIV) ANTIBODIES HIV (1 & 2) SCREENING

HIV 1/2 AND P24 ANTIGEN RESULT

NON - REACTIVE

by IMMUNOCHROMATOGRAPHY

INTERPRETATION:-

1.AIDS is caused by at least 2 known types of HIV viruses, HIV-1 and HIV HIV-2.

- 2. This NACO approved immuno-chromatographic solid phase ELISA assay detects antibodies against both HIV-1 and HIV-2 viruses.
- 3. The test is used for routine serologic screening of patients at risk for HIV-1 or HIV-2 infection.
- 4.All screening ELISA assays for HIV antibody detection have high sensitivity but have low specificity.
- 5.At this laboratory, all positive samples are cross checked for positivity with two alternate assays prior to reporting.

NOTE:-

- 1. Confirmatory testing by Western blot is recommended for patients who are reactive for HIV by this assay.
- 2.Antibodies against HIV-1 and HIV-2 are usually not detectable until 6 to 12 weeks following exposure (window period) and are almost always detectable by 12 months.
- 3. The test is not recommended for children born to HIV infected mothers till the child turns two years old (as HIV antibodies may be transmitted passively to the child trans-placentally).

FALSE NÉGATIVE RESULT SEEN IN:

- 1. Window period
- 2. Severe immuno-suppression including advanced AIDS.



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)



KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana



(A Unit of KOS Healthcare)



Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mrs. KRISHNA DEVI

AGE/ GENDER : 47 YRS/FEMALE **PATIENT ID** : 1779136

COLLECTED BY : REG. NO./LAB NO. : 012503050006

 REFERRED BY
 : 05/Mar/2025 07:58 AM

 BARCODE NO.
 : 01526480
 COLLECTION DATE
 : 05/Mar/2025 08:06AM

 CLIENT CODE.
 : KOS DIAGNOSTIC LAB
 REPORTING DATE
 : 05/Mar/2025 10:29AM

CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit Biological Reference interval

HEPATITIS B SURFACE ANTIGEN (HBsAg) SCREENING

REACTIVE

HEPATITIS B SURFACE ANTIGEN (HBsAg)

BESULI
by IMMUNOCHROMATOGRAPHY

INTERPRETATION:-

1.HBsAG is the first serological marker of HBV infection to appear in the blood (approximately 30-60 days after infection and prior to the onset of clinical disease). It is also the last viral protein to disappear from blood and usually disappears by three months after infection in self limiting acute Hepatitis B viral infection.

2.Persistence of HBsAg in blood for more than six months implies chronic infection. It is the most common marker used for diagnosis of an acute Hepatitis B infection but has very limited role in assessing patients suffering from chronic hepatitis.

FALSE NEGATIVE RESULT SEEN IN:

- 1. Window period.
- 2.Infection with HBsAg mutant strains
- 3. Hepatitis B Surface antigen (HBsAg) is the earliest indicator of HBV infection. Usually it appears in 27 41 days (as early as 14 days).
- 4.Appears 7 26 days before biochemical abnormalities. Peaks as ALT rises. Persists during the acute illness. Usually disappears 12- 20 weeks after the onset of symptoms / laboratory abnormalities in 90% of cases.
- 5.ls the most reliable serologic marker of HBV infection. Persistence > 6 months defines carrier state. May also be found in chronic infection. Hepatitis B vaccination does not cause a positive HBsAg. Titers are not of clinical value.

NOTE:-

1.All reactive HBsAG Should be reconfirmed with neutralization test(HBsAg confirmatory test).

2.Anti - HAV IgM appears at the same time as symptoms in > 99% of cases, peaks within the first month, becomes nondetectable in 12 months (usually 6 months). Presence confirms diagnosis of recent acute infection.



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUĞAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)





(A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mrs. KRISHNA DEVI

AGE/ GENDER : 47 YRS/FEMALE **PATIENT ID** : 1779136

COLLECTED BY : REG. NO./LAB NO. : 012503050006

 REFERRED BY
 : 05/Mar/2025 07:58 AM

 BARCODE NO.
 : 01526480
 COLLECTION DATE
 : 05/Mar/2025 08:06AM

 CLIENT CODE.
 : KOS DIAGNOSTIC LAB
 REPORTING DATE
 : 05/Mar/2025 09:00AM

CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Test Name Value Unit Biological Reference interval

VDRL

VDRL NON REACTIVE NON REACTIVE

by IMMUNOCHROMATOGRAPHY

INTERPRETATION:

1. Does not become positive until 7 - 10 days after appearance of chancre.

- 2. High titer (>1:16) active disease.
- 3.Low titer (<1:8) biological falsepositive test in 90% cases or due to late or late latent syphillis.
- 4.Treatment of primary syphillis causes progressive decline tonegative VDRL within 2 years.
- 5. Rising titer (4X) indicates relapse, reinfection, or treatment failure and need for retreatment.
- 6. May benonreactive in early primary, late latent, and late syphillis (approx. 25% ofcases).
- 7. Reactive and weakly reactive tests should always be confirmed with FTA-ABS (fluorescent treponemal antibody absorption test).

SHORTTERM FALSE POSITIVE TEST RESULTS (<6 MONTHS DURATION) MAY OCCURIN:

- 1. Acute viral illnesses (e.g., hepatitis, measles, infectious mononucleosis)
- 2.M. pneumoniae; Chlamydia; Malaria infection.
- 3. Some immunizations
- 4.Pregnancy (rare)

LONGTERM FALSE POSITIVE TEST RESULTS (>6 MONTHS DURATION) MAY OCCUR IN:

- 1. Serious underlying disease e.g., collagen vascular diseases, leprosy, malignancy.
- 2.Intravenous drug users.
- 3. Rheumatoid arthritis, thyroiditis, AIDS, Sjogren's syndrome.
- 4.< 10 % of patients older thanage 70 years.
- 5. Patients taking some anti-hypertensive drugs.

*** End Of Report ***



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUĞAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)

