

**Dr. Vinay Chopra**  
 MD (Pathology & Microbiology)  
 Chairman & Consultant Pathologist

**Dr. Yugam Chopra**  
 MD (Pathology)  
 CEO & Consultant Pathologist

<b>NAME</b>	: Mr. SAHIL	<b>PATIENT ID</b>	: 1780475
<b>AGE/ GENDER</b>	: 20 YRS/MALE	<b>REG. NO./LAB NO.</b>	: 012503060019
<b>COLLECTED BY</b>	:	<b>REGISTRATION DATE</b>	: 06/Mar/2025 09:55 AM
<b>REFERRED BY</b>	:	<b>COLLECTION DATE</b>	: 06/Mar/2025 09:58AM
<b>BARCODE NO.</b>	: 01526549	<b>REPORTING DATE</b>	: 06/Mar/2025 12:07PM
<b>CLIENT CODE.</b>	: KOS DIAGNOSTIC LAB		
<b>CLIENT ADDRESS</b>	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

### IMMUNOPATHOLOGY/SEROLOGY

#### ANTI HUMAN IMMUNODEFICIENCY VIRUS (HIV) DUO ULTRA WITH (P-24 ANTIGEN DETECTION)

HIV 1/2 AND P24 ANTIGEN: SERUM **1120.33<sup>H</sup>** S/CO **NEGATIVE: < 1.00**  
 by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY) **POSITIVE: > 1.00**

**HIV 1/2 AND P24 ANTIGEN RESULT** **REACTIVE**  
 by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

#### INTERPRETATION:-

RESULT (INDEX)	REMARKS
< 1.00	NON - REACTIVE
> = 1.00	PROVISIONALLY REACTIVE

Non-Reactive result implies that antibodies to HIV 1/ 2 have not been detected in the sample . This means that patient has either not been exposed to HIV 1/ 2 infection or the sample has been tested during the "window phase" i.e. before the development of detectable levels of antibodies. Hence a Non Reactive result does not exclude the possibility of exposure or infection with HIV 1/ 2.

#### RECOMMENDATIONS:

1. Results to be clinically correlated
2. Rarely falsenegativity/positivity may occur.

\*\*\* End Of Report \*\*\*





**DR.VINAY CHOPRA**  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)



**DR.YUGAM CHOPRA**  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY)

