

TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT.



|   | MD (Patho   | y Chopra<br>ology & Microbiology)<br>& Consultant Patholog   |  | (Pathology)  |
|---|---|--|--|--|
| AME   | : Mrs. BOBBY  |  |  |  |
| GE/ GENDER  | : 49 YRS/FEMALE   |  | PATIENT ID   | : 1786936  |
| OLLECTED BY   | :   |  | REG. NO./LAB NO.   | : 012503110024   |
| EFERRED BY  | :   |  | <b>REGISTRATION DATE</b>   | : 11/Mar/2025 09:42 AM   |
| ARCODE NO.  | :01526920   |  | COLLECTION DATE  | : 11/Mar/2025 09:46AM  |
| LIENT CODE.   | : KOS DIAGNOSTIC LAB  |  | REPORTING DATE   | : 11/Mar/2025 11:36AM  |
| LIENT ADDRESS   | : 6349/1, NICHOLSON R   | OAD, AMBALA CANT   | Г  |  |
| Fest Name   |   | Value  | Unit   | Biological Reference interval  |
|   | CI  | INICAL CHEMI   | STRY/BIOCHEMIST  | 'RV  |
|   |   |  | E FASTING (F)  | N  |
| LUCOSE FASTING  |   | 320.2 <sup>H</sup>   | mg/dL  | NORMAL: < 100.0  |
| NTERPRETATION<br>N ACCORDANCE WIT<br>. A fasting plasma g<br>. A fasting plasma d   | SE - PEROXIDASE (GOD-POD)<br>TH AMERICAN DIABETES ASS<br>plucose level below 100 mg<br>plucose level between 100  | /dl is considered norn<br>- 125 mg/dl is conside   | nal.<br>red as glucose intolerant or   | PREDIABETIC: 100.0 - 125.0<br>DIABETIC: > 0R = 126.0<br>prediabetic. A fasting and post-prandial blood |
| <b><u>VTERPRETATION</u></b><br>• A <b>CCORDANCE WIT</b><br>• A fasting plasma g<br>• A fasting plasma g<br>• st (after consumpt | TH AMERICAN DIABETES ASS<br>glucose level below 100 mg<br>jlucose level between 100<br>ion of 75 ams of alucose) is   | /dl is considered norn<br>- 125 mg/dl is conside<br>recommended for all<br>ng/dl is highly sugges<br>excess of 125 mg/dl o | nal.<br>red as glucose intolerant or<br>such patients.<br>tive of diabetic state. A repe<br>on both occasions is confirm | DIABETIC: > 0R = 126.0<br>prediabetic. A fasting and post-prandial blood                               |
| NTERPRETATION<br>N ACCORDANCE WIT<br>A fasting plasma g<br>A fasting plasma g<br>est (after consumpt                            | TH AMERICAN DIABETES ASS<br>glucose level below 100 mg<br>jlucose level between 100<br>ion of 75 ams of alucose) is   | /dl is considered norn<br>125 mg/dl is conside<br>recommended for all  | nal.<br>red as glucose intolerant or<br>such patients.<br>tive of diabetic state. A repe<br>on both occasions is confirm | DIABETIC: $> 0R = 126.0$   |
| <b><u>VTERPRETATION</u></b><br>• A <b>CCORDANCE WIT</b><br>• A fasting plasma g<br>• A fasting plasma g<br>• st (after consumpt | TH AMERICAN DIABETES ASS<br>glucose level below 100 mg<br>jlucose level between 100<br>ion of 75 ams of alucose) is   | /dl is considered norn<br>- 125 mg/dl is conside<br>recommended for all<br>ng/dl is highly sugges<br>excess of 125 mg/dl o | nal.<br>red as glucose intolerant or<br>such patients.<br>tive of diabetic state. A repe<br>on both occasions is confirm | DIABETIC: > 0R = 126.0<br>prediabetic. A fasting and post-prandial blood                               |
| <b><u>VTERPRETATION</u></b><br>• A <b>CCORDANCE WIT</b><br>• A fasting plasma g<br>• A fasting plasma g<br>• st (after consumpt | TH AMERICAN DIABETES ASS<br>glucose level below 100 mg<br>jlucose level between 100<br>ion of 75 ams of alucose) is   | /dl is considered norn<br>- 125 mg/dl is conside<br>recommended for all<br>ng/dl is highly sugges<br>excess of 125 mg/dl o | nal.<br>red as glucose intolerant or<br>such patients.<br>tive of diabetic state. A repe<br>on both occasions is confirm | DIABETIC: > 0R = 126.0<br>prediabetic. A fasting and post-prandial blood                               |
| VTERPRETATION<br>N ACCORDANCE WIT<br>. A fasting plasma g<br>. A fasting plasma g<br>est (after consumpt                        | TH AMERICAN DIABETES ASS<br>glucose level below 100 mg<br>jlucose level between 100<br>ion of 75 ams of alucose) is   | /dl is considered norn<br>- 125 mg/dl is conside<br>recommended for all<br>ng/dl is highly sugges<br>excess of 125 mg/dl o | nal.<br>red as glucose intolerant or<br>such patients.<br>tive of diabetic state. A repe<br>on both occasions is confirm | DIABETIC: > 0R = 126.0<br>prediabetic. A fasting and post-prandial blood                               |
| VTERPRETATION<br>A ACCORDANCE WIT<br>A fasting plasma g<br>A fasting plasma g<br>est (after consumpt                            | TH AMERICAN DIABETES ASS<br>glucose level below 100 mg<br>jlucose level between 100<br>ion of 75 ams of alucose) is   | /dl is considered norn<br>- 125 mg/dl is conside<br>recommended for all<br>ng/dl is highly sugges<br>excess of 125 mg/dl o | nal.<br>red as glucose intolerant or<br>such patients.<br>tive of diabetic state. A repe<br>on both occasions is confirm | DIABETIC: > 0R = 126.0<br>prediabetic. A fasting and post-prandial blood                               |
| VTERPRETATION<br>I ACCORDANCE WIT<br>A fasting plasma g<br>A fasting plasma g<br>st (after consumpt                             | TH AMERICAN DIABETES ASS<br>glucose level below 100 mg<br>glucose level between 100-<br>ion of 75 gms of glucose) is<br>glucose level of above 125 r<br>ing plasma glucose level in | /dl is considered norn<br>- 125 mg/dl is conside<br>recommended for all<br>ng/dl is highly sugges<br>excess of 125 mg/dl o | nal.<br>red as glucose intolerant or<br>such patients.<br>tive of diabetic state. A repe<br>on both occasions is confirm | DIABETIC: > 0R = 126.0<br>prediabetic. A fasting and post-prandial blood                               |
| VTERPRETATION<br>N ACCORDANCE WIT<br>. A fasting plasma g<br>. A fasting plasma g<br>est (after consumpt                        | TH AMERICAN DIABETES ASS<br>glucose level below 100 mg<br>glucose level between 100-<br>ion of 75 gms of glucose) is<br>glucose level of above 125 r<br>ing plasma glucose level in | /dl is considered norn<br>- 125 mg/dl is conside<br>recommended for all<br>ng/dl is highly sugges<br>excess of 125 mg/dl o | nal.<br>red as glucose intolerant or<br>such patients.<br>tive of diabetic state. A repe<br>on both occasions is confirm | DIABETIC: > 0R = 126.0<br>prediabetic. A fasting and post-prandial blood                               |
| NTERPRETATION<br>N ACCORDANCE WIT<br>A fasting plasma g<br>A fasting plasma g<br>est (after consumpt                            | TH AMERICAN DIABETES ASS<br>glucose level below 100 mg<br>glucose level between 100-<br>ion of 75 gms of glucose) is<br>glucose level of above 125 r<br>ing plasma glucose level in | /dl is considered norn<br>- 125 mg/dl is conside<br>recommended for all<br>ng/dl is highly sugges<br>excess of 125 mg/dl o | nal.<br>red as glucose intolerant or<br>such patients.<br>tive of diabetic state. A repe<br>on both occasions is confirm | DIABETIC: > 0R = 126.0<br>prediabetic. A fasting and post-prandial blood                               |
| NTERPRETATION<br>N ACCORDANCE WIT<br>A fasting plasma g<br>A fasting plasma g<br>est (after consumpt                            | TH AMERICAN DIABETES ASS<br>glucose level below 100 mg<br>glucose level between 100-<br>ion of 75 gms of glucose) is<br>glucose level of above 125 r<br>ing plasma glucose level in | /dl is considered norn<br>- 125 mg/dl is conside<br>recommended for all<br>ng/dl is highly sugges<br>excess of 125 mg/dl o | nal.<br>red as glucose intolerant or<br>such patients.<br>tive of diabetic state. A repe<br>on both occasions is confirm | DIABETIC: > 0R = 126.0<br>prediabetic. A fasting and post-prandial blood                               |
| VTERPRETATION<br>A ACCORDANCE WIT<br>A fasting plasma g<br>A fasting plasma g<br>est (after consumpt                            | TH AMERICAN DIABETES ASS<br>glucose level below 100 mg<br>glucose level between 100-<br>ion of 75 gms of glucose) is<br>glucose level of above 125 r<br>ing plasma glucose level in | /dl is considered norn<br>- 125 mg/dl is conside<br>recommended for all<br>ng/dl is highly sugges<br>excess of 125 mg/dl o | nal.<br>red as glucose intolerant or<br>such patients.<br>tive of diabetic state. A repe<br>on both occasions is confirm | DIABETIC: > 0R = 126.0<br>prediabetic. A fasting and post-prandial blood                               |

KOS Diagnostic Lab (A Unit of KOS Healthcare)





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)

