

TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT.



		Chopra y & Microbiology) Consultant Pathologist	Dr. Yugam MD (CEO & Consultant	(Pathology)
IAME	: Mr. SANDEEP SHARMA			
GE/ GENDER	: 53 YRS/MALE	PATI	ENT ID	: 1791517
COLLECTED BY	:	REG.	NO./LAB NO.	: 012503150005
REFERRED BY	:	REGI	STRATION DATE	: 15/Mar/2025 08:21 AM
BARCODE NO.	:01527094	COLL	ECTION DATE	: 15/Mar/2025 08:22AM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	REPO	ORTING DATE	: 15/Mar/2025 11:16AM
LIENT ADDRESS	: 6349/1, NICHOLSON ROA	AD, AMBALA CANTT		
Fest Name		Value	Unit	Biological Reference interval
	CLIN	ICAL CHEMISTRY	/BIOCHEMIST	RY
		GLUCOSE FAS	ГING (F)	
GLUCOSE FASTING	G (F): PLASMA SE - PEROXIDASE (GOD-POD)	136.77 ^H	mg/dL	NORMAL: < 100.0 PREDIABETIC: 100.0 - 125.0
A ACCORDANCE WIT A fasting plasma g A fasting plasma g	H AMERICAN DIABETES ASSOC lucose level below 100 mg/dl lucose level between 100 - 12	is considered normal. 25 mg/dl is considered as o	lucose intolerant or p	DIABETIC: $> 0R = 126.0$ prediabetic. A fasting and post-prandial blood
 A fasting plasma g A fasting plasma g est (after consumpt) 	lucose level below 100 mg/dl lucose level between 100 - 12 ion of 75 ams of alucose) is red	is considered normal. 25 mg/dl is considered as g commended for all such pa /dl is highly suggestive of c cess of 125 mg/dl on both	itients. liabetic state. A repea occasions is confirma	DIABETIC: $> 0R = 126.0$
N ACCORDANCE WIT 1. A fasting plasma g 2. A fasting plasma g est (after consumpt	lucose level below 100 mg/dl lucose level between 100 - 12 ion of 75 ams of alucose) is red	is considered normal. 25 mg/dl is considered as g commended for all such pa	itients. liabetic state. A repea occasions is confirma	DIABETIC: > 0R = 126.0
N ACCORDANCE WIT 1. A fasting plasma g 2. A fasting plasma g est (after consumpt	lucose level below 100 mg/dl lucose level between 100 - 12 ion of 75 ams of alucose) is red	is considered normal. 25 mg/dl is considered as g commended for all such pa /dl is highly suggestive of c cess of 125 mg/dl on both	itients. liabetic state. A repea occasions is confirma	DIABETIC: > 0R = 126.0
N ACCORDANCE WIT 1. A fasting plasma g 2. A fasting plasma g est (after consumpt	lucose level below 100 mg/dl lucose level between 100 - 12 ion of 75 ams of alucose) is red	is considered normal. 25 mg/dl is considered as g commended for all such pa /dl is highly suggestive of c cess of 125 mg/dl on both	itients. liabetic state. A repea occasions is confirma	DIABETIC: > 0R = 126.0
N ACCORDANCE WIT 1. A fasting plasma g 2. A fasting plasma g est (after consumpt	lucose level below 100 mg/dl lucose level between 100 - 12 ion of 75 ams of alucose) is red	is considered normal. 25 mg/dl is considered as g commended for all such pa /dl is highly suggestive of c cess of 125 mg/dl on both	itients. liabetic state. A repea occasions is confirma	DIABETIC: > 0R = 126.0
N ACCORDANCE WIT 1. A fasting plasma g 2. A fasting plasma g est (after consumpt	lucose level below 100 mg/dl lucose level between 100 - 12 ion of 75 ams of alucose) is red	is considered normal. 25 mg/dl is considered as g commended for all such pa /dl is highly suggestive of c cess of 125 mg/dl on both	itients. liabetic state. A repea occasions is confirma	DIABETIC: > 0R = 126.0
N ACCORDANCE WIT 1. A fasting plasma g 2. A fasting plasma g est (after consumpt	lucose level below 100 mg/dl lucose level between 100 - 12 ion of 75 ams of alucose) is red	is considered normal. 25 mg/dl is considered as g commended for all such pa /dl is highly suggestive of c cess of 125 mg/dl on both	itients. liabetic state. A repea occasions is confirma	DIABETIC: > 0R = 126.0
N ACCORDANCE WIT 1. A fasting plasma g 2. A fasting plasma g est (after consumpt	lucose level below 100 mg/dl lucose level between 100 - 12 ion of 75 ams of alucose) is red	is considered normal. 25 mg/dl is considered as g commended for all such pa /dl is highly suggestive of c cess of 125 mg/dl on both	itients. liabetic state. A repea occasions is confirma	DIABETIC: > 0R = 126.0
N ACCORDANCE WIT 1. A fasting plasma g 2. A fasting plasma g est (after consumpt	lucose level below 100 mg/dl lucose level between 100 - 12 ion of 75 ams of alucose) is red	is considered normal. 25 mg/dl is considered as g commended for all such pa /dl is highly suggestive of c cess of 125 mg/dl on both	itients. liabetic state. A repea occasions is confirma	DIABETIC: > 0R = 126.0
N ACCORDANCE WIT 1. A fasting plasma g 2. A fasting plasma g est (after consumpt	lucose level below 100 mg/dl lucose level between 100 - 12 ion of 75 ams of alucose) is red	is considered normal. 25 mg/dl is considered as g commended for all such pa /dl is highly suggestive of c cess of 125 mg/dl on both	itients. liabetic state. A repea occasions is confirma	DIABETIC: > 0R = 126.0

KOS Diagnostic Lab (A Unit of KOS Healthcare)





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)

 KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana

 KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt -133 001, Haryana

 0171-2643898, +91 99910 43898
 care@koshealthcare.com
 www.koshealthcare.com

