

Dr. Vinay Chopra
 MD (Pathology & Microbiology)
 Chairman & Consultant Pathologist

Dr. Yugam Chopra
 MD (Pathology)
 CEO & Consultant Pathologist

NAME	: Mrs. RAJNI JAIN	PATIENT ID	: 1796967
AGE/ GENDER	: 66 YRS/FEMALE	REG. NO./LAB NO.	: 012503180055
COLLECTED BY	: SURJESH	REGISTRATION DATE	: 18/Mar/2025 07:21 PM
REFERRED BY	:	COLLECTION DATE	: 18/Mar/2025 07:23PM
BARCODE NO.	: 01527357	REPORTING DATE	: 19/Mar/2025 12:28PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB		
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

TUMOUR MARKER

CARCINO EMBRYONIC ANTIGEN (CEA)

CARCINO EMBRYONIC ANTIGEN (CEA): SERUM	6.24^H	ng/mL	< 5.0
--	-------------------------	-------	-------

by CLIA (CHEMILUMINESCENCE IMMUNOASSAY)

INTERPRETATION:

1. Carcinoembryonic antigen (CEA) is a glycoprotein normally found in embryonic endodermal epithelium.
2. Increased levels may be found in patients with primary colorectal cancer or other malignancies including medullary thyroid carcinoma and breast, gastrointestinal tract, liver, lung, ovarian, pancreatic, and prostatic cancers.
3. Serial monitoring of CEA should begin prior to initiation of cancer therapy to verify post therapy decrease in concentration and to establish a baseline for evaluating possible recurrence. Levels generally return to normal within 1 to 4 months after removal of cancerous tissue.

CLINICAL SIGNIFICANCE:

1. Monitoring colorectal cancer and selected other cancers such as medullary thyroid carcinoma
2. May be useful in assessing the effectiveness of chemotherapy or radiation treatment.

NOTE:

1. Carcinoembryonic antigen levels should not be used for screening of the general population for undetected cancers.
2. Grossly elevated carcino-embryonic antigen (CEA) concentrations (>20 ng/mL) in a patient with compatible symptoms are strongly suggestive of the presence of cancer and also suggest metastasis.
3. Most healthy subjects (97%) have values < or =3.0 ng/mL.
4. After removal of a colorectal tumor, the serum CEA concentration should return to normal by 6 weeks, unless there is residual tumor.
5. Increases in test values over time in a patient with a history of cancer suggest tumor recurrence.

*** End Of Report ***




 DR. VINAY CHOPRA

CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)


 DR. YUGAM CHOPRA

CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY)

