

KOS Diagnostic Lab (A Unit of KOS Healthcare)



Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologist

Dr. Yugam Chopra MD (Pathology) CEO & Consultant Pathologist

NAME : Mr. IQBAL

AGE/ GENDER : 54 YRS/MALE **PATIENT ID** : 1811741

COLLECTED BY : 012503300005 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 30/Mar/2025 08:15 AM BARCODE NO. :01528006 **COLLECTION DATE** : 30/Mar/2025 08:48AM CLIENT CODE. : KOS DIAGNOSTIC LAB REPORTING DATE :30/Mar/2025 11:56AM

CLIENT ADDRESS : 6349/1, NICHOLSON ROAD, AMBALA CANTT

Value Unit Test Name **Biological Reference interval**

CLINICAL CHEMISTRY/BIOCHEMISTRY GLUCOSE FASTING (F) AND POST PRANDIAL (PP)

GLUCOSE FASTING (F): PLASMA mg/dL NORMAL: < 100.0 109.02^{H}

by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD) PREDIABETIC: 100.0 - 125.0

DIABETIC: > 0R = 126.0

GLUCOSE POST PRANDIAL (PP): PLASMA 182.82H mg/dL NORMAL: < 140.00

by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD) PREDIABETIC: 140.0 - 200.0

DIABETIC: > 0R = 200.0

INTERPRETATION:

IN ACCORDANCE WITH AMERICAN DIABETES ASSOCIATION GUIDELINES:

- 1. A fasting plasma glucose below 100 mg/dL and post-prandial plasma glucose level below 140 mg/dl is considered normal.
- 2. A fasting plasma glucose level between 100 125 mg/dl and post-prandial plasma glucose level between 140 200 mg/dL is considered as glucose intolerant or pre diabetic. A fasting and post-prandial blood test (after consumption of 75 gms of glucose) is recommended for all such patients.
- 3. A fasting plasma glucose level of above 125 mg/dL and post-prandial plasma glucose level above 200 mg/dL is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for all such patients. A fasting plasma glucose level in excess of 125 mg/dl on both occasions is confirmatory for diabetic state.

* End Of Report ***



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST

