Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra
MD (Pathology)
CEO & Consultant Pathologist

NAME	**: Mrs. POONAM**		
AGE/ GENDER	: 47 YRS/FEMALE	**PATIENT ID**	: 1816421
COLLECTED BY	: SURJESH	**REG. NO./LAB NO.**	**: 012504030041**
REFERRED BY	: CENTRAL PHOENIX CLUB (AMBALA CANTT)	**REGISTRATION DATE**	: 03/Apr/2025 12:31 PM
BARCODE NO.	: 01528297	**COLLECTION DATE**	: 03/Apr/2025 01:38PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	**REPORTING DATE**	: 03/Apr/2025 03:51PM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

CLINICAL CHEMISTRY/BIOCHEMISTRY

IRON PROFILE

Test Name	Value	Unit	Biological Reference interval
IRON: SERUM *by FERROZINE, SPECTROPHOTOMETRY*	70.79	µg/dL	50.0 - 170.0
UNSATURATED IRON BINDING CAPACITY (UIBC) :SERUM *by FERROZINE, SPECTROPHOTOMETERY*	172.36	µg/dL	150.0 - 336.0
TOTAL IRON BINDING CAPACITY (TIBC) :SERUM *by SPECTROPHOTOMETERY*	243.15	µg/dL	230 - 430
%TRANSFERRIN SATURATION: SERUM *by CALCULATED, SPECTROPHOTOMETERY (FERENE)*	29.11	%	15.0 - 50.0
TRANSFERRIN: SERUM *by SPECTROPHOTOMETERY (FERENE)*	**172.64**[L]	mg/dL	200.0 - 350.0

INTERPRETATION:-

VARIABLES	*ANEMIA OF CHRONIC DISEASE*	*IRON DEFICIENCY ANEMIA*	*THALASSEMIA α/β TRAIT*
SERUM IRON:	Normal to Reduced	Reduced	Normal
TOTAL IRON BINDING CAPACITY:	Decreased	Increased	Normal
% TRANSFERRIN SATURATION:	Decreased	Decreased < 12-15 %	Normal
SERUM FERRITIN:	Normal to Increased	Decreased	Normal or Increased

IRON:
1.Serum iron studies is recommended for differential diagnosis of microcytic hypochromic anemia.i.e iron deficiency anemia, zinc deficiency anemia,anemia of chronic disease and thalassemia syndromes.
2. It is essential to isolate iron deficiency anemia from Beta thalassemia syndromes because during iron replacement which is therapeutic for iron deficiency anemia, is severely contra-indicated in Thalassemia.

TOTAL IRON BINDING CAPACITY (TIBC):
1.It is a direct measure of protein transferrin which transports iron from the gut to storage sites in the bone marrow.

% TRANSFERRIN SATURATION:
1.Occurs in idiopathic hemochromatosis and transfusional hemosiderosis where no unsaturated iron binding capacity is available for iron mobilization. Similar condition is seen in congenital deficiency of transferrin.

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)

Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra
MD (Pathology)
CEO & Consultant Pathologist

NAME	**: Mrs. POONAM**		
AGE/ GENDER	: 47 YRS/FEMALE	**PATIENT ID**	: 1816421
COLLECTED BY	: SURJESH	**REG. NO./LAB NO.**	**: 012504030041**
REFERRED BY	: CENTRAL PHOENIX CLUB (AMBALA CANTT)	**REGISTRATION DATE**	: 03/Apr/2025 12:31 PM
BARCODE NO.	: 01528297	**COLLECTION DATE**	: 03/Apr/2025 01:38PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	**REPORTING DATE**	: 03/Apr/2025 04:25PM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name	Value	Unit	Biological Reference interval

ENDOCRINOLOGY

THYROID FUNCTION TEST: TOTAL

TRIIODOTHYRONINE (T3): SERUM *by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)*	0.82	ng/mL	0.35 - 1.93
THYROXINE (T4): SERUM *by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)*	**3.94**^{L}	µgm/dL	4.87 - 12.60
THYROID STIMULATING HORMONE (TSH): SERUM *by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)*	1.858	µIU/mL	0.35 - 5.50

3rd GENERATION, ULTRASENSITIVE

INTERPRETATION:

TSH levels are subject to circadian variation, reaching peak levels between 2-4 a.m and at a minimum between 6-10 pm. The variation is of the order of 50%.Hence time of the day has influence on the measured serum TSH concentrations. TSH stimulates the production and secretion of the metabolically active hormones, thyroxine (T4)and triiodothyronine (T3).Failure at any level of regulation of the hypothalamic-pituitary-thyroid axis will result in either underproduction (hypothyroidism) or overproduction(hyperthyroidism) of T4 and/or T3.

CLINICAL CONDITION	T3	T4	TSH
Primary Hypothyroidism:	Reduced	Reduced	Increased (Significantly)
Subclinical Hypothyroidism:	Normal or Low Normal	Normal or Low Normal	High
Primary Hyperthyroidism:	Increased	Increased	Reduced (at times undetectable)
Subclinical Hyperthyroidism:	Normal or High Normal	Normal or High Normal	Reduced

LIMITATIONS:-

1. T3 and T4 circulates in reversibly bound form with Thyroid binding globulins (TBG),and to a lesser extent albumin and Thyroid binding Pre Albumin so conditions in which TBG and protein levels alter such as pregnancy, excess estrogens, androgens, anabolic steroids and glucocorticoids may falsely affect the T3 and T4 levels and may cause false thyroid values for thyroid function tests.
2. Normal levels of T4 can also be seen in Hyperthyroid patients with :T3 Thyrotoxicosis, Decreased binding capacity due to hypoproteinemia or ingestion of certain drugs (e.g.: phenytoin , salicylates).
3. Serum T4 levels in neonates and infants are higher than values in the normal adult , due to the increased concentration of TBG in neonate serum.
4. TSH may be normal in central hypothyroidism , recent rapid correction of hyperthyroidism or hypothyroidism , pregnancy , phenytoin therapy.

TRIIODOTHYRONINE (T3)		THYROXINE (T4)		THYROID STIMULATING HORMONE (TSH)	
Age	**Refferance Range (ng/mL)**	**Age**	**Refferance Range (µg/dL)**	**Age**	**Reference Range (µIU/mL)**
0 - 7 Days	**0.20 - 2.65**	**0 - 7 Days**	**5.90 – 18.58**	**0 - 7 Days**	**2.43 - 24.3**
7 Days - 3 Months	**0.36 - 2.59**	**7 Days - 3 Months**	**6.39 - 17.66**	**7 Days - 3 Months**	**0.58 - 11.00**
3 - 6 Months	**0.51 - 2.52**	**3 - 6 Months**	**6.75 – 17.04**	**3 Days – 6 Months**	**0.70 - 8.40**

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)

Dr. Vinay Chopra
MD (Pathology & Microbiology)
Chairman & Consultant Pathologist

Dr. Yugam Chopra
MD (Pathology)
CEO & Consultant Pathologist

NAME	**: Mrs. POONAM**		
AGE/ GENDER	: 47 YRS/FEMALE	**PATIENT ID**	: 1816421
COLLECTED BY	: SURJESH	**REG. NO./LAB NO.**	**: 012504030041**
REFERRED BY	: CENTRAL PHOENIX CLUB (AMBALA CANTT)	**REGISTRATION DATE**	: 03/Apr/2025 12:31 PM
BARCODE NO.	: 01528297	**COLLECTION DATE**	: 03/Apr/2025 01:38PM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	**REPORTING DATE**	: 03/Apr/2025 04:25PM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name			Value	Unit	Biological Reference interval
6 - 12 Months	0.74 - 2.40	6 - 12 Months	7.10 – 16.16	6 – 12 Months	0.70 - 7.00
1 - 10 Years	0.92 - 2.28	1 - 10 Years	6.00 - 13.80	1 – 10 Years	0.60 - 5.50
11- 19 Years	0.35 - 1.93	11 - 19 Years	4.87- 13.20	11 – 19 Years	0.50 – 5.50
> 20 years (Adults)	0.35 - 1.93	> 20 Years (Adults)	4.87 - 12.60	> 20 Years (Adults)	0.35– 5.50
RECOMMENDATIONS OF TSH LEVELS DURING PREGNANCY (µIU/mL)					
1st Trimester			0.10 – 2.50		
2nd Trimester			0.20 – 3.00		
3rd Trimester			0.30 – 4.10		

INCREASED TSH LEVELS:

1.Primary or untreated hypothyroidism may vary from 3 times to more than 100 times normal depending upon degree of hypofunction.
2.Hypothyroid patients receiving insufficient thyroid replacement therapy.
3.Hashimotos thyroiditis
4.DRUGS: Amphetamines, iodine containing agents & dopamine antagonist.
5.Neonatal period, increase in 1st 2-3 days of life due to post-natal surge

DECREASED TSH LEVELS:

1.Toxic multi-nodular goiter & Thyroiditis.
2.Over replacement of thyroid hormone in treatment of hypothyroidism.
3.Autonomously functioning Thyroid adenoma
4.Secondary pituitary or hypothalamic hypothyroidism
5.Acute psychiatric illness
6.Severe dehydration.
7.DRUGS: Glucocorticoids, Dopamine, Levodopa, T4 replacement therapy, Anti-thyroid drugs for thyrotoxicosis.
8.Pregnancy: 1st and 2nd Trimester

*** End Of Report ***

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)