



Dr. Vinay ChopraDr. Yugam ChopraMD (Pathology & Microbiology)MD (Pathology)Chairman & Consultant PathologistCEO & Consultant Pathologist					
NAME	: Mrs. SEEMA				
AGE/ GENDER	: 42 YRS/FEMALE	PA	TIENT ID	: 1819921	
COLLECTED BY	: SURJESH	RE	G. NO./LAB NO.	: 012504060029	
REFERRED BY	:	RE	GISTRATION DATE	: 06/Apr/2025 09:59 AM	
BARCODE NO.	: 01528455	CO	LLECTION DATE	:06/Apr/2025 10:00AM	
CLIENT CODE.	: KOS DIAGNOSTIC LAB	RE	PORTING DATE	:06/Apr/2025 11:48AM	
CLIENT ADDRESS	: 6349/1, NICHOLSON ROA	D, AMBALA CANTT			
Test Name		Value	Unit	Biological Reference interval	
		ENDOCRI	NOLOGY		
	Т	HYROID FUNCTION	ON TEST: TOTAL		
TRIIODOTHYRON by CMIA (CHEMILUMIN	INE (T3): SERUM IESCENT MICROPARTICLE IMMUN	0.922 OASSAY)	ng/mL	0.35 - 1.93	
THYROXINE (T4): SERUM 8.76 by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)			µgm/dL	4.87 - 12.60	
	ATING HORMONE (TSH):		µIU/mL	0.35 - 5.50	
3rd GENERATION, ULT <u>INTERPRETATION</u> :					
day has influence on the trilodothyronine (T3).Fai	measured serum TSH concentrations	. TSH stimulates the produc	tion and secretion of the m	m. The variation is of the order of 50%.Hence time of the tabolically active hormones, thyroxine (T4)and er underproduction (hypothyroidism) or	
CLINICAL CONDITION	T3		T4	TSH	

CLINICAL CONDITION	T3	T4	TSH
Primary Hypothyroidism:	Reduced	Reduced	Increased (Significantly)
Subclinical Hypothyroidism:	Normal or Low Normal	Normal or Low Normal	High
Primary Hyperthyroidism:	Increased	Increased	Reduced (at times undetectable)
Subclinical Hyperthyroidism:	Normal or High Normal	Normal or High Normal	Reduced

LIMITATIONS:-

1. T3 and T4 circulates in reversibly bound form with Thyroid binding globulins (TBG), and to a lesser extent albumin and Thyroid binding Pre Albumin so conditions in which TBG and protein levels alter such as pregnancy, excess estrogens, androgens, anabolic steroids and glucocorticoids may falsely affect the T3 and T4 levels and may cause false thyroid values for thyroid function tests.

2. Normal levels of T4 can also be seen in Hyperthyroid patients with :T3 Thyrotoxicosis, Decreased binding capacity due to hypoproteinemia or ingestion of certain drugs (e.g.: phenytoin , salicylates).

3. Serum T4 levels in neonates and infants are higher than values in the normal adult , due to the increased concentration of TBG in neonate serum.

4. TSH may be normal in central hypothyroidism, recent rapid correction of hyperthyroidism or hypothyroidism, pregnancy, phenytoin therapy.

TRIIODOTHYRONINE (T3)		THYROXINE (T4)		THYROID STIMULATING HORMONE (TSH)		
Age	Refferance Range (ng/mL)	Age	Refferance Range (µg/dL)	Age	Reference Range (μIU/mL)	
0 - 7 Days	0.20 - 2.65	0 - 7 Days	5.90 - 18.58	0 - 7 Days	2.43 - 24.3	
7 Days - 3 Months	0.36 - 2.59	7 Days - 3 Months	6.39 - 17.66	7 Days - 3 Months	0.58 - 11.00	
3 - 6 Months	0.51 - 2.52	3 - 6 Months	6.75 - 17.04	3 Days – 6 Months	0.70 - 8.40	





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)



TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT







	Dr. Vinay Chopra MD (Pathology & Microbiology) Chairman & Consultant Pathologis		Pathology)
NAME	: Mrs. SEEMA		
AGE/ GENDER	: 42 YRS/FEMALE	PATIENT ID	: 1819921
COLLECTED BY	: SURJESH	REG. NO./LAB NO.	: 012504060029
REFERRED BY	:	REGISTRATION DATE	: 06/Apr/2025 09:59 AM
BARCODE NO.	: 01528455	COLLECTION DATE	:06/Apr/2025 10:00AM
CLIENT CODE.	: KOS DIAGNOSTIC LAB	REPORTING DATE	:06/Apr/2025 11:48AM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, AMBALA CANTT		

Test Name			Value	Unit	t	Biological Reference interval
6 - 12 Months	0.74 - 2.40	6 - 12 Months	7.10 - 16.16	6 - 12 Months	0.70 - 7.00	
1 - 10 Years	0.92 - 2.28	1 - 10 Years	6.00 - 13.80	1 – 10 Years	0.60 - 5.50	
11-19 Years	0.35 - 1.93	11 - 19 Years	4.87- 13.20	11 – 19 Years	0.50 - 5.50	
> 20 years (Adults)	0.35 - 1.93	> 20 Years (Adults)	4.87 - 12.60	> 20 Years (Adults)	0.35-5.50	
	RECOM	MENDATIONS OF TSH L	EVELS DURING PRE	GNANCY (µIU/mL)		
1st Trimester				0.10 - 2.50		
2nd Trimester			0.20 - 3.00			
3rd Trimester			0.30 - 4.10			

INCREASED TSH LEVELS:

1.Primary or untreated hypothyroidism may vary from 3 times to more than 100 times normal depending upon degree of hypofunction.

2. Hypothyroid patients receiving insufficient thyroid replacement therapy.

3.Hashimotos thyroiditis

4.DRUGS: Amphetamines, iodine containing agents & dopamine antagonist.

5.Neonatal period, increase in 1st 2-3 days of life due to post-natal surge

DECREASED TSH LEVELS:

1.Toxic multi-nodular goiter & Thyroiditis.

2. Over replacement of thyroid hormone in treatment of hypothyroidism.

3. Autonomously functioning Thyroid adenoma

4.Secondary pituitary or hypothalamic hypothyroidism

5. Acute psychiatric illness

6.Severe dehydration.

7.DRUGS: Glucocorticoids, Dopamine, Levodopa, T4 replacement therapy, Anti-thyroid drugs for thyrotoxicosis.

8.Pregnancy: 1st and 2nd Trimester

*** End Of Report **





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)

