

TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT.



	MD	: Vinay Cho) (Pathology & I airman & Consu			(Pathology)
NAME AGE/ GENDER COLLECTED BY REFERRED BY BARCODE NO. CLIENT CODE. CLIENT ADDRESS	: Mr. AMI LAL : 62 YRS/MALE : : : 01528559 : KOS DIAGNOSTI : 6349/1, NICHOI		MBALA CANTT	PATIENT ID REG. NO./LAB NO. REGISTRATION DATE COLLECTION DATE REPORTING DATE	: 1822154 : 012504080006 : 08/Apr/2025 07:11 AM : 08/Apr/2025 08:33AM : 08/Apr/2025 12:51PM
Test Name			Value	Unit	Biological Reference interval
		CLINICA	L CHEMIS	STRY/BIOCHEMIS	STRY
				E FASTING (F)	
GLUCOSE FASTIN by GLUCOSE OXIDAS	IG (F): PLASMA E - PEROXIDASE (GOL	D-POD)	148.13 ^H	mg/dL	NORMAL: < 100.0 PREDIABETIC: 100.0 - 125.0
N ACCORDANCE WIT 1. A fasting plasma g 2. A fasting plasma g est (after consumpt 3. A fasting plasma g	lucose level below 1 lucose level betwee ion of 75 gms of gluc lucose level of abov	100 mg/dl is co en 100 - 125 mg cose) is recom re 125 mg/dl is	onsidered norma g/dl is considere mended for all s highly suggestiv	al. ed as glucose intolerant or uch patients.	at post-prandial is strongly recommended for
test (after consumpt 3. A fasting plasma g	lucose level below 1 lucose level betwee ion of 75 gms of gluc lucose level of abov	100 mg/dl is co en 100 - 125 mg cose) is recomi re 125 mg/dl is level in excess	onsidered norma g/dl is considered mended for all s highly suggestiv of 125 mg/dl or	al. ed as glucose intolerant or uch patients. ve of diabetic state. A repe n both occasions is confirm	prediabetic. A fasting and post-prandial blood at post-prandial is strongly recommended for a
N ACCORDANCE WIT 1. A fasting plasma g 2. A fasting plasma g test (after consumpt 3. A fasting plasma g	lucose level below 1 lucose level betwee ion of 75 gms of gluc lucose level of abov	100 mg/dl is co en 100 - 125 mg cose) is recomi re 125 mg/dl is level in excess	onsidered norma g/dl is considere mended for all s highly suggestiv	al. ed as glucose intolerant or uch patients. ve of diabetic state. A repe n both occasions is confirm	prediabetic. A fasting and post-prandial blood at post-prandial is strongly recommended for a
IN ACCORDANCE WIT 1. A fasting plasma g 2. A fasting plasma g test (after consumpt 3. A fasting plasma g	lucose level below 1 lucose level betwee ion of 75 gms of gluc lucose level of abov	100 mg/dl is co en 100 - 125 mg cose) is recomi re 125 mg/dl is level in excess	onsidered norma g/dl is considered mended for all s highly suggestiv of 125 mg/dl or	al. ed as glucose intolerant or uch patients. ve of diabetic state. A repe n both occasions is confirm	prediabetic. A fasting and post-prandial blood at post-prandial is strongly recommended for a
N ACCORDANCE WIT 1. A fasting plasma g 2. A fasting plasma g test (after consumpt 3. A fasting plasma g	lucose level below 1 lucose level betwee ion of 75 gms of gluc lucose level of abov	100 mg/dl is co en 100 - 125 mg cose) is recomi re 125 mg/dl is level in excess	onsidered norma g/dl is considered mended for all s highly suggestiv of 125 mg/dl or	al. ed as glucose intolerant or uch patients. ve of diabetic state. A repe n both occasions is confirm	prediabetic. A fasting and post-prandial blood at post-prandial is strongly recommended for a
N ACCORDANCE WIT 1. A fasting plasma g 2. A fasting plasma g test (after consumpt 3. A fasting plasma g	lucose level below 1 lucose level betwee ion of 75 gms of gluc lucose level of abov	100 mg/dl is co en 100 - 125 mg cose) is recomi re 125 mg/dl is level in excess	onsidered norma g/dl is considered mended for all s highly suggestiv of 125 mg/dl or	al. ed as glucose intolerant or uch patients. ve of diabetic state. A repe n both occasions is confirm	prediabetic. A fasting and post-prandial blood at post-prandial is strongly recommended for a
N ACCORDANCE WIT 1. A fasting plasma g 2. A fasting plasma g test (after consumpt 3. A fasting plasma g	lucose level below 1 lucose level betwee ion of 75 gms of gluc lucose level of abov	100 mg/dl is co en 100 - 125 mg cose) is recomi re 125 mg/dl is level in excess	onsidered norma g/dl is considered mended for all s highly suggestiv of 125 mg/dl or	al. ed as glucose intolerant or uch patients. ve of diabetic state. A repe n both occasions is confirm	prediabetic. A fasting and post-prandial blood at post-prandial is strongly recommended for a
N ACCORDANCE WIT 1. A fasting plasma g 2. A fasting plasma g test (after consumpt 3. A fasting plasma g	lucose level below 1 lucose level betwee ion of 75 gms of gluc lucose level of abov	100 mg/dl is co en 100 - 125 mg cose) is recomi re 125 mg/dl is level in excess	onsidered norma g/dl is considered mended for all s highly suggestiv of 125 mg/dl or	al. ed as glucose intolerant or uch patients. ve of diabetic state. A repe n both occasions is confirm	prediabetic. A fasting and post-prandial blood at post-prandial is strongly recommended for a
IN ACCORDANCE WIT 1. A fasting plasma g 2. A fasting plasma g test (after consumpt 3. A fasting plasma g	lucose level below 1 lucose level betwee ion of 75 gms of gluc lucose level of abov	100 mg/dl is co en 100 - 125 mg cose) is recomi re 125 mg/dl is level in excess	onsidered norma g/dl is considered mended for all s highly suggestiv of 125 mg/dl or	al. ed as glucose intolerant or uch patients. ve of diabetic state. A repe n both occasions is confirm	prediabetic. A fasting and post-prandial blood at post-prandial is strongly recommended for a

DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

回路

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)

KOS Central Lab: 6349/1, Nicholson Road, Ambala Cantt -133 001, Haryana KOS Molecular Lab: IInd Floor, Parry Hotel, Staff Road, Opp. GPO, Ambala Cantt -133 001, Haryana 0171-2643898, +91 99910 43898 | care@koshealthcare.com | www.koshealthcare.com

