



	Dr. Vinay Cho MD (Pathology & Chairman & Cons	Microbiology)		(Pathology)
NAME	: Mr. RANDHIR SINGH			
AGE/ GENDER	: 55 YRS/MALE		PATIENT ID	: 1554189
COLLECTED BY	:		REG. NO./LAB NO.	: 042407190002
REFERRED BY	:		REGISTRATION DATE	: 19/Jul/2024 01:54 PM
BARCODE NO.	: A0465998		COLLECTION DATE	: 19/Jul/2024 03:34PM
CLIENT CODE.	: KOS DIAGNOSTIC SHAHBAD		REPORTING DATE	: 19/Jul/2024 04:56PM
CLIENT ADDRESS	: 6349/1, NICHOLSON ROAD, A	MBALA CANTT		
Test Name		Value	Unit	Biological Reference interval
GLUCOSE RANDOM (R): PLASMA by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD)		185.39 <sup>H</sup>	mg/dL	PREDIABETIC: 140.0 - 200.0 DIABETIC: > 0R = 200.0
1. A random plasma 2. A random glucose (after consumption of 3. A random glucose	of 75 gms of glucose) is recommen	considered norm considered as glu ded for all such p suggestive of dia	ucose intolerant or prediat patients. abetic state. A repeat post	petic. A fasting and post-prnadial blood test -prandial is strongly recommended for all such for diabetic state.
IN ACCORDANCE WIT 1. A random plasma 2. A random glucose (after consumption of 3. A random glucose	glucose level below 140 mg/dl is o level between 140 - 200 mg/dl is o of 75 gms of glucose) is recommen level of above 200 mg/dl is highly lasma glucose level in excess of 12	considered norm considered as glu ded for all such p suggestive of dia	acose intolerant or prediab patients. abetic state. A repeat post occasions is confirmatory	-prandial is strongly recommended for all such
IN ACCORDANCE WIT 1. A random plasma 2. A random glucose (after consumption of 3. A random glucose	glucose level below 140 mg/dl is o level between 140 - 200 mg/dl is o of 75 gms of glucose) is recommen level of above 200 mg/dl is highly lasma glucose level in excess of 12	considered norm considered as glu ded for all such p suggestive of dia 25 mg/dl on both	acose intolerant or prediab patients. abetic state. A repeat post occasions is confirmatory	-prandial is strongly recommended for all such
IN ACCORDANCE WIT 1. A random plasma 2. A random glucose (after consumption of 3. A random glucose	glucose level below 140 mg/dl is o level between 140 - 200 mg/dl is o of 75 gms of glucose) is recommen level of above 200 mg/dl is highly lasma glucose level in excess of 12	considered norm considered as glu ded for all such p suggestive of dia 25 mg/dl on both	acose intolerant or prediab patients. abetic state. A repeat post occasions is confirmatory	-prandial is strongly recommended for all such
IN ACCORDANCE WIT 1. A random plasma 2. A random glucose (after consumption of 3. A random glucose	glucose level below 140 mg/dl is o level between 140 - 200 mg/dl is o of 75 gms of glucose) is recommen level of above 200 mg/dl is highly lasma glucose level in excess of 12	considered norm considered as glu ded for all such p suggestive of dia 25 mg/dl on both	acose intolerant or prediab patients. abetic state. A repeat post occasions is confirmatory	-prandial is strongly recommended for all such





**DR.VINAY CHOPRA** CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)

