

**Dr. Vinay Chopra**  
 MD (Pathology & Microbiology)  
 Chairman & Consultant Pathologist

**Dr. Yugam Chopra**  
 MD (Pathology)  
 CEO & Consultant Pathologist

|                       |                                        |                          |                        |
|-----------------------|----------------------------------------|--------------------------|------------------------|
| <b>NAME</b>           | : Mr. NARESH ARORA                     | <b>PATIENT ID</b>        | : 1690277              |
| <b>AGE/ GENDER</b>    | : 59 YRS/MALE                          | <b>REG. NO./LAB NO.</b>  | : 042503280007         |
| <b>COLLECTED BY</b>   | :                                      | <b>REGISTRATION DATE</b> | : 28/Mar/2025 05:51 PM |
| <b>REFERRED BY</b>    | :                                      | <b>COLLECTION DATE</b>   | : 28/Mar/2025 06:37PM  |
| <b>BARCODE NO.</b>    | : A1260750                             | <b>REPORTING DATE</b>    | : 28/Mar/2025 06:50PM  |
| <b>CLIENT CODE.</b>   | : KOS DIAGNOSTIC SHAHBAD               |                          |                        |
| <b>CLIENT ADDRESS</b> | : 6349/1, NICHOLSON ROAD, AMBALA CANTT |                          |                        |

| Test Name | Value | Unit | Biological Reference interval |
|-----------|-------|------|-------------------------------|
|-----------|-------|------|-------------------------------|

## CLINICAL CHEMISTRY/BIOCHEMISTRY

### URIC ACID

|                  |                   |       |             |
|------------------|-------------------|-------|-------------|
| URIC ACID: SERUM | 2.82 <sup>L</sup> | mg/dL | 3.60 - 7.70 |
|------------------|-------------------|-------|-------------|

by URICASE - OXIDASE PEROXIDASE

#### INTERPRETATION:-

1.GOUT occurs when high levels of Uric Acid in the blood cause crystals to form & accumulate around a joint.  
 2.Uric Acid is the end product of purine metabolism . Uric acid is excreted to a large degree by the kidneys and to a smaller degree in the intestinal tract by microbial degradation.

#### INCREASED:-

##### (A).DUE TO INCREASED PRODUCTION:-

- 1.Idiopathic primary gout.
- 2.Excessive dietary purines (organ meats,legumes,anchovies, etc).
- 3.Cytolytic treatment of malignancies especially leukemias & lymphomas.
- 4.Polycythemia vera & myeloid metaplasia.
- 5.Psoriasis.
- 6.Sickle cell anaemia etc.

##### (B).DUE TO DECREASED EXCRETION (BY KIDNEYS)

- 1.Alcohol ingestion.
- 2.Thiazide diuretics.
- 3.Lactic acidosis.
- 4.Aspirin ingestion (less than 2 grams per day ).
- 5.Diabetic ketoacidosis or starvation.
- 6.Renal failure due to any cause etc.

#### DECREASED:-

##### (A).DUE TO DIETARY DEFICIENCY

- 1.Dietary deficiency of Zinc, Iron and molybdenum.
- 2.Fanconi syndrome & Wilsons disease.
- 3.Multiple sclerosis .
- 4.Syndrome of inappropriate antidiuretic hormone (SIADH) secretion & low purine diet etc.

##### (B).DUE TO INCREASED EXCRETION

- 1.Drugs:-Probenecid , sulphinpyrazone, aspirin doses (more than 4 grams per day), corticosteroids and ACTH, anti-coagulants and estrogens etc.

\*\*\* End Of Report \*\*\*





**DR.VINAY CHOPRA**  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)



**DR.YUGAM CHOPRA**  
 CONSULTANT PATHOLOGIST  
 MBBS, MD (PATHOLOGY)

