A PIONEER DIAGNOSTIC CENTRE

【 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. BALWINDER KAUR			
AGE/ GENDER	: 66 YRS/FEMALE	PAT	IENT ID	: 1549115
COLLECTED BY	:	REG	. NO./LAB NO.	: 122407150001
REFERRED BY	:	REG	ISTRATION DATE	: 15/Jul/2024 08:47 AM
BARCODE NO.	: 12503595	COL	LECTION DATE	: 15/Jul/2024 09:12AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITU	ITE REP	ORTING DATE	: 15/Jul/2024 04:49PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBA	LA CITY - HARYAN	NA	
Test Name		Value	Unit	Biological Reference interval
	SWAS	THYA WELLN	ESS PANEL: 1.5	
	CON	IPLETE BLOOD	COUNT (CBC)	
RED BLOOD CELLS (F	RBCS) COUNT AND INDICES			
HAEMOGLOBIN (HB)		11.8 ^L	gm/dL	12.0 - 16.0
RED BLOOD CELL (RE	COUNT	4.32	Millions/cr	mm 3.50 - 5.00
PACKED CELL VOLUN		35.2 ^L	%	37.0 - 50.0
MEAN CORPUSCULA		81.5	fL	80.0 - 100.0
MEAN CORPUSCULA	R HAEMOGLOBIN (MCH) UTOMATED HEMATOLOGY ANALYZER	27.3	pg	27.0 - 34.0
MEAN CORPUSCULA	R HEMOGLOBIN CONC. (MCHC)	33.5	g/dL	32.0 - 36.0
	ION WIDTH (RDW-CV) UTOMATED HEMATOLOGY ANALYZER	13	%	11.00 - 16.00
	ION WIDTH (RDW-SD) UTOMATED HEMATOLOGY ANALYZER	40.4	fL	35.0 - 56.0
MENTZERS INDEX by CALCULATED		18.87	RATIO	BETA THALASSEMIA TRAIT: < 13 IRON DEFICIENCY ANEMIA: >13
GREEN & KING INDE by calculated	Х	24.51	RATIO	BETA THALASSEMIA TRAIT: < = 65.0 IRON DEFICIENCY ANEMIA: > 65
WHITE BLOOD CELLS	<u>S (WBCS)</u>			IKON DEI IGIENGT ANEIVIIA. 203
	Y BY SF CUBE & MICROSCOPY	10740	/cmm	4000 - 11000
DIFFERENTIAL LEUCO	<u>JCYTE COUNT (DLC)</u>	()	~	50.70
NEUTROPHILS by FLOW CYTOMETRY	Y BY SF CUBE & MICROSCOPY	63	%	50 - 70
LYMPHOCYTES		29	%	20 - 40
by FLOW CYTOMETRY EOSINOPHILS	Y BY SF CUBE & MICROSCOPY	2	%	1-6

DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS , MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST



A PIONEER DIAGNOSTIC CENTRE

【 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. BALWINDER KAUR		
AGE/ GENDER	: 66 YRS/FEMALE	PATIENT ID	: 1549115
COLLECTED BY	:	REG. NO./LAB NO.	: 122407150001
REFERRED BY	:	REGISTRATION DATE	: 15/Jul/2024 08:47 AM
BARCODE NO.	: 12503595	COLLECTION DATE	: 15/Jul/2024 09:12AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	REPORTING DATE	: 15/Jul/2024 04:49PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY	- HARYANA	

Test Name	Value	Unit	Biological Reference interval
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY			
MONOCYTES	6	%	2 - 12
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY			
BASOPHILS	0	%	0 - 1
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY			
ABSOLUTE LEUKOCYTES (WBC) COUNT			
ABSOLUTE NEUTROPHIL COUNT	6766	/cmm	2000 - 7500
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY			
ABSOLUTE LYMPHOCYTE COUNT	3115 ^L	/cmm	800 - 4900
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY			
ABSOLUTE EOSINOPHIL COUNT	215	/cmm	40 - 440
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	A PAR /		
ABSOLUTE MONOCYTE COUNT	644	/cmm	80 - 880
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	0	1	0 110
ABSOLUTE BASOPHIL COUNT by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	0	/cmm	0 - 110
PLATELETS AND OTHER PLATELET PREDICTIVE MARKED	DC		
PLATELET COUNT (PLT)	234000	/cmm	150000 - 450000
by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	0.07	04	0.400.0/
	0.26	%	0.10 - 0.36
by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	11	fl	(50, 12.0
MEAN PLATELET VOLUME (MPV) by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	11	fL	6.50 - 12.0
PLATELET LARGE CELL COUNT (P-LCC)	87000	/cmm	30000 - 90000
by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	07000	/ CITIITI	30000 - 70000
PLATELET LARGE CELL RATIO (P-LCR)	37.2	%	11.0 - 45.0
by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE		· -	
PLATELET DISTRIBUTION WIDTH (PDW)	16.6	%	15.0 - 17.0
by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE			
NOTE: TEST CONDUCTED ON EDTA WHOLE BLOOD			





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS , MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST





NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

💟 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. BALWINDER KAUR			
AGE/ GENDER	: 66 YRS/FEMALE	PATIE	NT ID	: 1549115
COLLECTED BY	:	REG. N	O./LAB NO.	: 122407150001
REFERRED BY	:	REGIST	FRATION DATE	: 15/Jul/2024 08:47 AM
BARCODE NO.	: 12503595	COLLE	CTION DATE	: 15/Jul/2024 09:12AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INS	TITUTE REPOR	TING DATE	: 15/Jul/2024 04:50PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AM	MBALA CITY - HARYANA		
Test Name		Value	Unit	Biological Reference interval
GLYCOSYLATED HAEM(WHOLE BLOOD		YCOSYLATED HAEMOG. 8.9 ^H	LOBIN (HBA1C) %	4.0 - 6.4
WHOLE BLOOD by HPLC (HIGH PERFORM ESTIMATED AVERAGE F	OGLOBIN (HbA1c): Mance liquid chromatography)			4.0 - 6.4 60.00 - 140.00
WHOLE BLOOD by HPLC (HIGH PERFORM ESTIMATED AVERAGE F by HPLC (HIGH PERFORM	OGLOBIN (HbA1c): MANCE LIQUID CHROMATOGRAPHY) PLASMA GLUCOSE MANCE LIQUID CHROMATOGRAPHY)	8.9 ^H 208.73 ^H	%	
WHOLE BLOOD by HPLC (HIGH PERFORM ESTIMATED AVERAGE F by HPLC (HIGH PERFORM INTERPRETATION:	OGLOBIN (HbA1c): MANCE LIQUID CHROMATOGRAPHY) PLASMA GLUCOSE MANCE LIQUID CHROMATOGRAPHY)	8.9 ^H 208.73 ^H SETES ASSOCIATION (ADA):	% mg/dL	60.00 - 140.00
WHOLE BLOOD by HPLC (HIGH PERFORM ESTIMATED AVERAGE F by HPLC (HIGH PERFORM INTERPRETATION: RE	OGLOBIN (HbA1c): MANCE LIQUID CHROMATOGRAPHY) PLASMA GLUCOSE MANCE LIQUID CHROMATOGRAPHY) AS PER AMERICAN DIAE	8.9 ^H 208.73 ^H SETES ASSOCIATION (ADA):	%	60.00 - 140.00
WHOLE BLOOD by HPLC (HIGH PERFORM ESTIMATED AVERAGE F by HPLC (HIGH PERFORM INTERPRETATION: RE Non diabu	OGLOBIN (HbA1c): MANCE LIQUID CHROMATOGRAPHY) PLASMA GLUCOSE MANCE LIQUID CHROMATOGRAPHY) AS PER AMERICAN DIAE FERENCE GROUP	8.9 ^H 208.73 ^H BETES ASSOCIATION (ADA): GLYCOSYLATED HI	% mg/dL EMOGLOGIB (HBAIC) in 9	60.00 - 140.00
WHOLE BLOOD by HPLC (HIGH PERFORM ESTIMATED AVERAGE F by HPLC (HIGH PERFORM INTERPRETATION: REI Non diab At R	OGLOBIN (HbA1c): MANCE LIQUID CHROMATOGRAPHY) PLASMA GLUCOSE MANCE LIQUID CHROMATOGRAPHY) AS PER AMERICAN DIAE FERENCE GROUP etic Adults >= 18 years	8.9 ^H 208.73 ^H BETES ASSOCIATION (ADA): GLYCOSYLATED HI	% mg/dL EMOGLOGIB (HBAIC) in 9	60.00 - 140.00
WHOLE BLOOD by HPLC (HIGH PERFORM ESTIMATED AVERAGE F by HPLC (HIGH PERFORM INTERPRETATION: REI Non diab At R	OGLOBIN (HbA1c): MANCE LIQUID CHROMATOGRAPHY) PLASMA GLUCOSE MANCE LIQUID CHROMATOGRAPHY) AS PER AMERICAN DIAE FERENCE GROUP etic Adults >= 18 years Risk (Prediabetes)	8.9 ^H 208.73 ^H BETES ASSOCIATION (ADA): GLYCOSYLATED HI S Age	% mg/dL <5.7 5.7 - 6.4 >= 6.5 > 19 Years	60.00 - 140.00
WHOLE BLOOD by HPLC (HIGH PERFORM ESTIMATED AVERAGE F by HPLC (HIGH PERFORM INTERPRETATION: REI Non diab At R Diag	OGLOBIN (HbA1c): MANCE LIQUID CHROMATOGRAPHY) PLASMA GLUCOSE MANCE LIQUID CHROMATOGRAPHY) AS PER AMERICAN DIAE FERENCE GROUP etic Adults >= 18 years Risk (Prediabetes) gnosing Diabetes	8.9 ^H 208.73 ^H BETES ASSOCIATION (ADA): GLYCOSYLATED HI	% mg/dL <5.7 5.7 - 6.4 >= 6.5	60.00 - 140.00
WHOLE BLOOD by HPLC (HIGH PERFORM ESTIMATED AVERAGE F by HPLC (HIGH PERFORM INTERPRETATION: RE Non diab At R Diag	OGLOBIN (HbA1c): MANCE LIQUID CHROMATOGRAPHY) PLASMA GLUCOSE MANCE LIQUID CHROMATOGRAPHY) AS PER AMERICAN DIAE FERENCE GROUP etic Adults >= 18 years Risk (Prediabetes)	8.9 ^H 208.73 ^H BETES ASSOCIATION (ADA): GLYCOSYLATED HI GLYCOSYLATED HI Goals of Therapy: Actions Suggested:	% mg/dL <5.7 5.7 - 6.4 >= 6.5 :> 19 Years <7.0 >8.0	60.00 - 140.00
WHOLE BLOOD by HPLC (HIGH PERFORM ESTIMATED AVERAGE F by HPLC (HIGH PERFORM INTERPRETATION: REI Non diab At R Diag	OGLOBIN (HbA1c): MANCE LIQUID CHROMATOGRAPHY) PLASMA GLUCOSE MANCE LIQUID CHROMATOGRAPHY) AS PER AMERICAN DIAE FERENCE GROUP etic Adults >= 18 years Risk (Prediabetes) gnosing Diabetes	8.9 ^H 208.73 ^H BETES ASSOCIATION (ADA): GLYCOSYLATED HI GLYCOSYLATED HI Goals of Therapy: Actions Suggested:	% mg/dL <5.7 5.7 - 6.4 >= 6.5 >> 19 Years < 7.0	60.00 - 140.00

COMMENTS:

1.Glycosylated hemoglobin (HbA1c) test is three monthly monitoring done to assess compliace with therapeutic regimen in diabetic patients.

2.Since Hb1c reflects long term fluctuations in blood glucose concentration, a diabetic patient who has recently under good control may still have high concentration of HbAlc. Converse is true for a diabetic previously under good control but now poorly controlled.

3. Target goals of < 7.0 % may be beneficial in patients with short duration of diabetes, long life expectancy and no significant cardiovascular disease. In patients with significant complications of diabetes, limited life expectancy or extensive co-morbid conditions, targetting a goal of < 7.0% may not be appropriate. 4. High

HbA1c (>9.0 -9.5 %) is strongly associated with risk of development and rapid progression of microvascular and nerve complications 5.Any condition that shorten RBC life span like acute blood loss, hemolytic anemia falsely lower HbA1c results.

6.HbA1c results from patients with HbSS,HbSC and HbD must be interpreted with caution, given the pathological processes including anemia, increased red cell turnover, and transfusion requirement that adversely impact HbA1c as a marker of long-term gycemic control.

7. Specimens from patients with polycythemia or post-splenctomy may exhibit increse in HbA1c values due to a somewhat longer life span of the red cells.






DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)



🔽 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. BALWINDER KAUR		
AGE/ GENDER	: 66 YRS/FEMALE	PATIENT ID	: 1549115
COLLECTED BY	:	REG. NO./LAB NO.	: 122407150001
REFERRED BY	:	REGISTRATION DATE	: 15/Jul/2024 08:47 AM
BARCODE NO.	: 12503595	COLLECTION DATE	: 15/Jul/2024 09:12AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	REPORTING DATE	: 15/Jul/2024 04:50PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY -	HARYANA	
Test Name	Value	Unit	Biological Reference interval







DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS , MD (PATHOLOGY)

DR.YUGAM CHOPRA

CONSULTANT PATHOLOGIST





A PIONEER DIAGNOSTIC CENTRE

🔽 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. BALWINDER KAUR				
AGE/ GENDER	: 66 YRS/FEMALE	РАТ	TENT ID	: 1549115	
COLLECTED BY	:	REG	. NO./LAB NO.	: 122407150001	
REFERRED BY	:	REG	ISTRATION DATE	: 15/Jul/2024 08:47 AM	
BARCODE NO.	: 12503595	COL	LECTION DATE	: 15/Jul/2024 09:12AM	
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTIT	UTE Rep	ORTING DATE	: 15/Jul/2024 04:10PM	
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBA	LA CITY - HARYA	NA		
Test Name		Value	Unit	Biological Reference interval	
				_	
	ERYTHRC	OCYTE SEDIMEN	ITATION RATE (ES	R)	
	MENTATION RATE (ESR)	57 ^H	mm/1st l	hr 0 - 20	
by MODIFIED WESTER	RGREN AUTOMATED METHOD				
1. ESR is a non-specific test because an elevated result often indicates the presence of inflammation associated with infection, cancer and auto-					
immune disease, but	does not tell the health practitioner	exactly where the	inflammation is in the	e body or what is causing it.	
		ammation. For thi	s reason, the ESR is ty	picallý used in conjunction with other test such	
as C-reactive protein	he used to monitor disease activity	and rosponso to th	orany in both of the a	as daus aradto amos as llow as sozeaith aud	
ouctomic lunus on the	3. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as some others, such as				

systemic lupus erythematosus

CONDITION WITH LOW ESR

A low ESR can be seen with conditions that inhibit the normal sedimentation of red blood cells, such as a high red blood cell count

(polycythaemia), significantly high white blood cell count (leucocytosis), and some protein abnormalities. Some changes in red cell shape (such as sickle cells in sickle cell anaemia) also lower the ESR.

NOTE:

 ESR and C - reactive protein (C-RP) are both markers of inflammation.
 Generally, ESR does not change as rapidly as does CRP, either at the start of inflammation or as it resolves.
 CRP is not affected by as many other factors as is ESR, making it a better marker of inflammation.
 If the ESR is elevated, it is typically a result of two types of proteins, globulins or fibrinogen.
 Women tend to have a higher ESR, and menstruation and pregnancy can cause temporary elevations.
 Drugs such as dextran, methyldopa, oral contraceptives, penicillamine procainamide, theophylline, and vitamin A can increase ESR, while service contractions and pregnancy can be added and the start of the s aspirin, cortisone, and quinine may decrease it





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)





A PIONEER DIAGNOSTIC CENTRE

💟 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. BALWINDER KAUR		
AGE/ GENDER	: 66 YRS/FEMALE	PATIENT ID	: 1549115
COLLECTED BY	:	REG. NO./LAB NO.	: 122407150001
REFERRED BY	:	REGISTRATION DATE	: 15/Jul/2024 08:47 AM
BARCODE NO.	: 12503595	COLLECTION DATE	: 15/Jul/2024 09:12AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	REPORTING DATE	: 15/Jul/2024 04:52PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - H	ARYANA	

PERIPHERAL BLOOD SMEAR

TEST NAME:

PERIPHERAL BLOOD FILM/SMEAR (PBF)

RED BLOOD CELLS (RBC'S):

RBCs mostly appear normocytic & normochromic.No polychromatic cells or normoblasts noted.

WHITE BLOOD CELLS (WBC'S)

No immature leucocytes seen.

PLATELETS:

Platelets are adequate.

HEMOPARASITES:

NOT SEEN.

IMPRESSION:

Normocytic normochromic picture.





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)

NOT VALID FOR MEDICO LEGAL PURPOSE



A PIONEER DIAGNOSTIC CENTRE

🔽 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. BALWINDER KAUR			
AGE/ GENDER	: 66 YRS/FEMALE	РАТ	IENT ID	: 1549115
COLLECTED BY	:	REG	. NO./LAB NO.	: 122407150001
REFERRED BY	:	REG	ISTRATION DATE	: 15/Jul/2024 08:47 AM
BARCODE NO.	: 12503595	COL	LECTION DATE	: 15/Jul/2024 09:12AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INS	STITUTE REP	ORTING DATE	: 15/Jul/2024 04:50PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, A	MBALA CITY - HARYAN	NA	
Test Name		Value	Unit	Biological Reference interval
	CLIN	ICAL CHEMISTRY GLUCOSE FAS		Y
GLUCOSE FASTING (I by GLUCOSE OXIDAS	F): PLASMA E - PEROXIDASE (GOD-POD)	141.42 ^H	mg/dL	NORMAL: < 100.0 PREDIABETIC: 100.0 - 125.0 DIABETIC: > 0R = 126.0
	H AMERICAN DIABETES ASSOCIAT			

A fasting plasma glucose level below 100 mg/di is considered as glucose intolerant or prediabetic. A fasting and post-prandial blood test (after consumption of 75 gms of glucose) is recommended for all such patients.
 A fasting plasma glucose level of above 125 mg/dl is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for all such patients. A fasting plasma glucose level in excess of 125 mg/dl on both occasions is confirmatory for diabetic state.





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)

NOT VALID FOR MEDICO LEGAL PURPOSE



【 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. BALWINDER KAUR			
AGE/ GENDER	: 66 YRS/FEMALE		PATIENT ID	: 1549115
COLLECTED BY	:		REG. NO./LAB NO.	: 122407150001
REFERRED BY	:		REGISTRATION DATE	: 15/Jul/2024 08:47 AM
BARCODE NO.	: 12503595		COLLECTION DATE	: 15/Jul/2024 09:12AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INS	TITUTE	REPORTING DATE	: 15/Jul/2024 05:37PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AM	/IBALA CITY - H/	ARYANA	
Test Name		Value	Unit	Biological Reference interval
		LIPID PR	OFILE : BASIC	
CHOLESTEROL TOTA by CHOLESTEROL OX		257.79 ^H	mg/dL	OPTIMAL: < 200.0 BORDERLINE HIGH: 200.0 - 239.0 HIGH CHOLESTEROL: > OR = 240.0
TRIGLYCERIDES: SER by GLYCEROL PHOSE	UM HATE OXIDASE (ENZYMATIC)	387.2 ^H	mg/dL	OPTIMAL: < 150.0 BORDERLINE HIGH: 150.0 - 199.0 HIGH: 200.0 - 499.0 VERY HIGH: > OR = 500.0
HDL CHOLESTEROL (by SELECTIVE INHIBITI		47.77	mg/dL	LOW HDL: < 30.0 BORDERLINE HIGH HDL: 30.0 - 60.0 HIGH HDL: > OR = 60.0
LDL CHOLESTEROL: S by CALCULATED, SPE		132.58 ^H	mg/dL	OPTIMAL: < 100.0 ABOVE OPTIMAL: 100.0 - 129.0 BORDERLINE HIGH: 130.0 - 159.0 HIGH: 160.0 - 189.0 VERY HIGH: > OR = 190.0
NON HDL CHOLESTE by CALCULATED, SPE		210.02 ^H	mg/dL	OPTIMAL: < 130.0 ABOVE OPTIMAL: 130.0 - 159.0 BORDERLINE HIGH: 160.0 - 189.0 HIGH: 190.0 - 219.0 VERY HIGH: > OR = 220.0
VLDL CHOLESTEROL:		77.44 ^H	mg/dL	0.00 - 45.00
by CALCULATED, SPE TOTAL LIPIDS: SERUI by CALCULATED, SPE	M	902.78 ^H	mg/dL	350.00 - 700.00
CHOLESTEROL/HDL I by CALCULATED, SPE	RATIO: SERUM	5.4 ^H	RATIO	LOW RISK: 3.30 - 4.40 AVERAGE RISK: 4.50 - 7.0 MODERATE RISK: 7.10 - 11.0 HIGH RISK: > 11.0
LDL/HDL RATIO: SER by CALCULATED, SPE		2.78	RATIO	LOW RISK: 0.50 - 3.0 MODERATE RISK: 3.10 - 6.0 HIGH RISK: > 6.0

DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS , MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST

NOT VALID FOR MEDICO LEGAL PURPOSE

440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. **REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)**



Page 8 of 26



A PIONEER DIAGNOSTIC CENTRE

🔽 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. BALWINDER KAUR		
AGE/ GENDER	: 66 YRS/FEMALE	PATIENT ID	: 1549115
COLLECTED BY	:	REG. NO./LAB NO.	: 122407150001
REFERRED BY	:	REGISTRATION DATE	: 15/Jul/2024 08:47 AM
BARCODE NO.	: 12503595	COLLECTION DATE	: 15/Jul/2024 09:12AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	REPORTING DATE	: 15/Jul/2024 05:37PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY -	HARYANA	
Test Name	Value	Unit	Biological Reference interval

	Value	onit	biological Reference interval
TRIGLYCERIDES/HDL RATIO: SERUM	8.11 ^H	RATIO	3.00 - 5.00
by CALCULATED, SPECTROPHOTOMETRY			

INTERPRETATION:

1.Measurements in the same patient can show physiological& analytical variations. Three serial samples 1 week apart are recommended for Total Cholesterol, Triglycerides, HDL & LDL Cholesterol.

2. As per NLA-2014 guidelines, all adults above the age of 20 years should be screened for lipid status. Selective screening of children above the age of 2 years with a family history of premature cardiovascular disease or those with at least one parent with high total cholesterol is recommended.

3. Low HDL levels are associated with increased risk for Atherosclerotic Cardiovascular disease (ASCVD) due to insufficient HDL being available to participate in reverse cholesterol transport, the process by which cholesterol is eliminated from peripheral tissues. 4. NLA-2014 identifies Non HDL Cholesterol (an indicator of all atherogeniclipoproteins such as LDL, VLDL, IDL, Lpa, Chylomicron remnants) along with LDL-cholesterol as co- primary target for cholesterol lowering therapy. Note that major risk factors can modify treatment goals for LDL & Non HDL

5. Additional testing for Apolipoprotein B, hsCRP,Lp(a) & LP-PLA2 should be considered among patients with moderate risk for ASCVD for risk refinement



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)





A PIONEER DIAGNOSTIC CENTRE

🔽 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. BALWINDER KAUR		
AGE/ GENDER	: 66 YRS/FEMALE	PATIENT ID	: 1549115
COLLECTED BY	:	REG. NO./LAB NO.	: 122407150001
REFERRED BY	:	REGISTRATION DATE	: 15/Jul/2024 08:47 AM
BARCODE NO.	: 12503595	COLLECTION DATE	: 15/Jul/2024 09:12AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	REPORTING DATE	: 15/Jul/2024 05:37PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CIT	Y - HARYANA	
Test Name	Value	e Unit	Biological Reference interval
	LIVER FUN	CTION TEST (COMPLETE)	
BILIRUBIN TOTAL: S	ERUM 0.48 PECTROPHOTOMETRY	mg/dL	INFANT: 0.20 - 8.00 ADLILT: 0.00 - 1.20

by DIAZOTIZATION, SPECTROPHOTOMETRY	0.40	ing/ de	ADULT: 0.00 - 1.20	
BILIRUBIN DIRECT (CONJUGATED): SERUM by DIAZO MODIFIED, SPECTROPHOTOMETRY	0.15	mg/dL	0.00 - 0.40	
BILIRUBIN INDIRECT (UNCONJUGATED): SERUM by CALCULATED, SPECTROPHOTOMETRY	0.33	mg/dL	0.10 - 1.00	
SGOT/AST: SERUM by IFCC, WITHOUT PYRIDOXAL PHOSPHATE	14.52	U/L	7.00 - 45.00	
SGPT/ALT: SERUM	12.94	U/L	0.00 - 49.00	
by IFCC, WITHOUT PYRIDOXAL PHOSPHATE				
AST/ALT RATIO: SERUM	1.12	RATIO	0.00 - 46.00	
by CALCULATED, SPECTROPHOTOMETRY				
ALKALINE PHOSPHATASE: SERUM	187.57 ^H	U/L	40.0 - 130.0	
by PARA NITROPHENYL PHOSPHATASE BY AMINO METHYL PROPANOL	L			
GAMMA GLUTAMYL TRANSFERASE (GGT): SERUM by SZASZ, SPECTROPHTOMETRY	35.19	U/L	0.00 - 55.0	
TOTAL PROTEINS: SERUM by BIURET, SPECTROPHOTOMETRY	7.4	gm/dL	6.20 - 8.00	
ALBUMIN: SERUM	4.38	gm/dL	3.50 - 5.50	
by BROMOCRESOL GREEN	0.00	<i>i</i>	0.00 0.50	
GLOBULIN: SERUM	3.02	gm/dL	2.30 - 3.50	
by CALCULATED, SPECTROPHOTOMETRY				
A : G RATIO: SERUM	1.45	RATIO	1.00 - 2.00	
by CALCULATED, SPECTROPHOTOMETRY				

INTERPRETATION

NOTE:- To be correlated in individuals having SGOT and SGPT values higher than Normal Referance Range.

USE: - Differential diagnosis of diseases of hepatobiliary system and pancreas.

INCREASED:

DRUG HEPATOTOXICITY	> 2
ALCOHOLIC HEPATITIS	> 2 (Highly Suggestive)
CIRRHOSIS	1.4 - 2.0
INTRAHEPATIC CHOLESTATIS	> 1.5
HEPATOCELLULAR CARCINOMA & CHRONIC HEPATITIS	> 1.3 (Slightly Increased)





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA

CONSULTANT PATHOLOGIST

MBBS , MD (PATHOLOGY)







A PIONEER DIAGNOSTIC CENTRE

【 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. BALWINDER KAUR				
AGE/ GENDER	: 66 YRS/FEMALE	PATIENT ID	: 1549115		
COLLECTED BY	:	REG. NO./LAB NO.	: 122407150001		
REFERRED BY	:	REGISTRATION DATE	: 15/Jul/2024 08:47 AM		
BARCODE NO.	: 12503595	COLLECTION DATE	: 15/Jul/2024 09:12AM		
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	REPORTING DATE	: 15/Jul/2024 05:37PM		
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA				

rest name Unit Biological Reference Intervi	Test Name	Value	Unit	Biological Reference interva
---	-----------	-------	------	------------------------------

DECREASED:

1. Acute Hepatitis due to virus, drugs, toxins (with AST increased 3 to 10 times upper limit of normal)

2. Extra Hepatic cholestatis: 0.8 (normal or slightly decreased).

PRO	GNO	STIC 3	SIGNIFI	CANCE:	

NORMAL	< 0.65
GOOD PROGNOSTIC SIGN	0.3 - 0.6
POOR PROGNOSTIC SIGN	1.2 - 1.6



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)





A PIONEER DIAGNOSTIC CENTRE

└ 0171-2532620, 8222896961 **□** pkrjainhealthcare@gmail.com

NAME	: Mrs. BALWINDER KAUR		
AGE/ GENDER	: 66 YRS/FEMALE	PATIENT ID	: 1549115
COLLECTED BY	:	REG. NO./LAB NO.	: 122407150001
REFERRED BY	:	REGISTRATION DATE	: 15/Jul/2024 08:47 AM
BARCODE NO.	: 12503595	COLLECTION DATE	: 15/Jul/2024 09:12AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	REPORTING DATE	: 15/Jul/2024 05:19PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - H	HARYANA	
Test Name	Value	Unit	Biological Reference interval
	KIDNEY FUNCT	ION TEST (COMPLETE)	
UREA: SERUM	49.07	mg/dL	10.00 - 50.00

UREA: SERUM	49.07	mg/dL	10.00 - 50.00
by UREASE - GLUTAMATE DEHYDROGENASE (GLDH)			
CREATININE: SERUM by ENZYMATIC, SPECTROPHOTOMETERY	1.42 ^H	mg/dL	0.40 - 1.20
BLOOD UREA NITROGEN (BUN): SERUM by CALCULATED, SPECTROPHOTOMETRY	22.93	mg/dL	7.0 - 25.0
BLOOD UREA NITROGEN (BUN)/CREATININE RATIO: SERUM	16.15	RATIO	10.0 - 20.0
by CALCULATED, SPECTROPHOTOMETRY UREA/CREATININE RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	34.56	RATIO	
URIC ACID: SERUM	5.96 ^H	mg/dL	2.50 - 6.80
CALCIUM: SERUM by ARSENAZO III, SPECTROPHOTOMETRY	9.53	mg/dL	8.50 - 10.60
PHOSPHOROUS: SERUM by phosphomolybdate, spectrophotometry	3.74	mg/dL	2.30 - 4.70
ELECTROLYTES			
SODIUM: SERUM by ISE (ION SELECTIVE ELECTRODE)	142.6	mmol/L	135.0 - 150.0
POTASSIUM: SERUM by ise (ion selective electrode)	5	mmol/L	3.50 - 5.00
CHLORIDE: SERUM by ISE (ION SELECTIVE ELECTRODE)	106.95	mmol/L	90.0 - 110.0
ESTIMATED GLOMERULAR FILTERATION RATE			
ESTIMATED GLOMERULAR FILTERATION RATE (eGFR): SERUM <i>by CALCULATED</i>	40.8		

INTERPRETATION:

To differentiate between pre- and post renal azotemia.

INCREASED RATIO (>20:1) WITH NORMAL CREATININE:

1. Prerenal azotemia (BUN rises without increase in creatinine) e.g. heart failure, salt depletion, dehydration, blood loss) due to decreased glomerular filtration rate.

2. Catabolic states with increased tissue breakdown.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)

NOT VALID FOR MEDICO LEGAL PURPOSE



A PIONEER DIAGNOSTIC CENTRE

🔽 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. BALWINDER KAUR		
AGE/ GENDER	: 66 YRS/FEMALE	PATIENT ID	: 1549115
COLLECTED BY	:	REG. NO./LAB NO.	: 122407150001
REFERRED BY	:	REGISTRATION DATE	: 15/Jul/2024 08:47 AM
BARCODE NO.	: 12503595	COLLECTION DATE	: 15/Jul/2024 09:12AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	REPORTING DATE	: 15/Jul/2024 05:19PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY	- HARYANA	
Test Name	Value	Unit	Biological Reference interval
3. GI haemorrhage.		Unit	Biological Reference Interval
3. GI haemorrhage. 4. High protein intake	÷.	Unit	Biological Reference Interval
 GI haemorrhage. High protein intake Impaired renal fun Excess protein inta 	e. Inction plus ike or production or tissue breakdown (e.g. int		
 GI haemorrhage. High protein intake Impaired renal fun Excess protein inta burns, surgery, cache 	e. Inction plus Ike or production or tissue breakdown (e.g. interview exia, high fever).		
burns, surgery, cache 7. Urine reabsorption	e. Inction plus Ike or production or tissue breakdown (e.g. in Exia, high fever). I (e.g. ureter colostomy)		
 GI haemorrhage. High protein intake Impaired renal fun Excess protein inta burns, surgery, cache Urine reabsorption Reduced muscle m 	e. Inction plus Ike or production or tissue breakdown (e.g. interview exia, high fever).		

1. Postrenal azotemia (BUN rises disproportionately more than creatinine) (e.g. obstructive uropathy).

2. Prerenal azotemia superimposed on renal disease.

DECREASED RATIO (<10:1) WITH DECREASED BUN :

1. Acute tubular necrosis.

- 2. Low protein diet and starvation.
- 3. Severe liver disease.
- 4. Other causes of decreased urea synthesis.

5. Repeated dialysis (urea rather than creatinine diffuses out of extracellular fluid).

- 6. Inherited hyperammonemias (urea is virtually absent in blood).
- 7. SIADH (syndrome of inappropiate antidiuretic harmone) due to tubular secretion of urea.
- 8. Pregnancy.

DECREASED RATIO (<10:1) WITH INCREASED CREATININE:

1. Phenacimide therapy (accelerates conversion of creatine to creatinine).

2. Rhabdomyolysis (releases muscle creatinine).

3. Muscular patients who develop renal failure.

INAPPROPIATE RATIO:

1. Diabetic ketoacidosis (acetoacetate causes false increase in creatinine with certain methodologies, resulting in normal ratio when dehydration should produce an increased BUN/creatinine ratio).

2. Cephalosporin therapy (interferes with creatinine measurement).

CKD STAGE	DESCRIPTION	GFR (mL/min/1.73m2)	ASSOCIATED FINDINGS
G1	Normal kidney function	>90	No proteinuria
G2	Kidney damage with normal or high GFR	>90	Presence of Protein , Albumin or cast in urine
G3a	Mild decrease in GFR	60 -89	
G3b	Moderate decrease in GFR	30-59	
G4	Severe decrease in GFR	15-29	
G5	Kidney failure	<15	





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)





A PIONEER DIAGNOSTIC CENTRE

0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. BALWINDER KAUR				
AGE/ GENDER	: 66 YRS/FEMALE	PATIENT ID	: 1549115		
COLLECTED BY	:	REG. NO./LAB NO.	: 122407150001		
REFERRED BY	:	REGISTRATION DATE	: 15/Jul/2024 08:47 AM		
BARCODE NO.	: 12503595	COLLECTION DATE	: 15/Jul/2024 09:12AM		
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	REPORTING DATE	: 15/Jul/2024 05:19PM		
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA				

Test Name	Value	Unit	Biological Reference interval

COMMENTS:

1. Estimated Glomerular filtration rate (eGFR) is the sum of filtration rates in all functioning nephrons and so an estimation of the GFR provides a measure of functioning nephrons of the kidney. 2. eGFR calculated using the 2009 CKD-EPI creatinine equation and GFR category reported as per KDIGO guideline 2012

3. In patients, with eGFR creatinine between 45-59 ml/min/1.73 m2 (G3) and without any marker of Kidney damage, It is recommended to measure eGFR with Cystatin C for confirmation of CKD

4. eGFR category G1 OR G2 does not fullfill the criteria for CKD, in the absence of evidence of Kidney Damage 5. In a suspected case of Acute Kidney Injury (AKI), measurement of eGFR should be done after 48-96 hours of any Intervention or procedure 6. eGFR calculated by Serum Creatinine may be less accurate due to certain factors like Race, Muscle Mass, Diet, Certain Drugs. In such cases, eGFR should be calculated using Serum Cystatin C 7. A decrease in eGFR implies either progressive renal disease, or a reversible process causing decreased nephron function (eg, severe dehydration).

ADVICE:

KDIGO guideline, 2012 recommends Chronic Kidney Disease (CKD) should be classified based on cause, eGFR category and Albuminuria (ACR) category. GFR & ACR category combined together reflect risk of progression and helps Clinician to identify the individual who are progressing at more rapid rate than anticipated



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)





🔽 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. BALWINDER KAUR		
AGE/ GENDER	: 66 YRS/FEMALE	PATIENT ID	: 1549115
COLLECTED BY	:	REG. NO./LAB NO.	: 122407150001
REFERRED BY	:	REGISTRATION DATE	: 15/Jul/2024 08:47 AM
BARCODE NO.	: 12503595	COLLECTION DATE	: 15/Jul/2024 09:12AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	REPORTING DATE	: 15/Jul/2024 04:50PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY	- HARYANA	

Test Name	Value	Unit	Biological Reference interval
	IRON PROP	FILE	
IRON: SERUM by FERROZINE, SPECTROPHOTOMETRY	106.8	μg/dL	37.0 - 145.0
UNSATURATED IRON BINDING CAPACITY (UIBC) :SERUM by FERROZINE, SPECTROPHOTOMETERY	187.72	µg/dL	150.0 - 336.0
TOTAL IRON BINDING CAPACITY (TIBC) :SERUM by SPECTROPHOTOMETERY	294.52 ^L	μg/dL	230 - 430
%TRANSFERRIN SATURATION: SERUM by CALCULATED, SPECTROPHOTOMETERY (FERENE)	36.26	%	15.0 - 50.0
TRANSFERRIN: SERUM by SPECTROPHOTOMETERY (FERENE)	209.11	mg/dL	200.0 - 350.0

VARIABLES	ANEMIA OF CHRONIC DISEASE	IRON DEFICIENCY ANEMIA	THALASSEMIA α/β TRAIT
SERUM IRON:	Normal to Reduced	Reduced	Normal
TOTAL IRON BINDING CAPACITY:	Decreased	Increased	Normal
% TRANSFERRIN SATURATION:	Decreased	Decreased < 12-15 %	Normal
SERUM FERRITIN:	Normal to Increased	Decreased	Normal or Increased
SERUM FERRITIN:	Normal to Increased	Decreased	Normal or Increase

IRON:

1.Serum iron studies is recommended for differential diagnosis of microcytic hypochromic anemia.i.e iron deficiency anemia, zinc deficiency anemia, anemia of chronic disease and thalassemia syndromes.

2. It is essential to isolate iron deficiency anemia from Beta thalassemia syndromes because during iron replacement which is therapeutic for TOTAL IRON BINDING CAPACITY (TIBC): 1.1t is a direct measure of protein transferrin which transports iron from the gut to storage sites in the bone marrow.

% TRANSFERRIN SATURATION:

1. Occurs in idiopathic hemochromatosis and transfusional hemosiderosis where no unsaturated iron binding capacity is available for iron mobilization. Similar condition is seen in congenital deficiency of transferrin.





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)

NOT VALID FOR MEDICO LEGAL PURPOSE





A PIONEER DIAGNOSTIC CENTRE

🕻 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. BALWINDER KAUR			
AGE/ GENDER	: 66 YRS/FEMALE	PATIE	INT ID	: 1549115
COLLECTED BY	:	REG. N	IO./LAB NO.	: 122407150001
REFERRED BY	:	REGIS	TRATION DATE	: 15/Jul/2024 08:47 AM
BARCODE NO.	: 12503595	COLLE	ECTION DATE	: 15/Jul/2024 09:12AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITU	TE REPO	RTING DATE	: 15/Jul/2024 05:13PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBAL	A CITY - HARYANA		
Test Name		Value	Unit	Biological Reference interval
Test Name				Biological Reference interval
Test Name		Value ENDOCRINO		Biological Reference interval
Test Name	ТНУК		LOGY	Biological Reference interval
TRIIODOTHYRONINE		ENDOCRINO ROID FUNCTION 1.035	LOGY	Biological Reference interval
TRIIODOTHYRONINE by cmia (chemilumin THYROXINE (T4): SEF	: (T3): SERUM ESCENT MICROPARTICLE IMMUNOASSAY)	ENDOCRINO COID FUNCTION 1.035 8.58	LOGY TEST: TOTAL	

INTERPRETATION:

TSH levels are subject to circadian variation, reaching peak levels between 2-4 a.m and at a minimum between 6-10 pm. The variation is of the order of 50%. Hence time of the day has influence on the measured serum TSH concentrations. TSH stimulates the production and secretion of the metabolically active hormones, thyroxine (T4) and trilodothyronine (T3). Failure at any level of regulation of the hypothalamic-pituitary-thyroid axis will result in either underproduction (hypothyroidism) or overproduction(hyperthyroidism) of T4 and/or T3.

CLINICAL CONDITION	T3	T4	TSH
Primary Hypothyroidism:	Reduced	Reduced	Increased (Significantly)
Subclinical Hypothyroidism:	Normal or Low Normal	Normal or Low Normal	High
Primary Hyperthyroidism:	Increased	Increased	Reduced (at times undetectable)
Subclinical Hyperthyroidism:	Normal or High Normal	Normal or High Normal	Reduced

LIMITATIONS:-

1. T3 and T4 circulates in reversibly bound form with Thyroid binding globulins (TBG), and to a lesser extent albumin and Thyroid binding Pre Albumin so conditions in which TBG and protein levels alter such as pregnancy, excess estrogens, androgens, anabolic steroids and glucocorticoids may falsely affect the T3 and T4 levels and may cause false thyroid values for thyroid function tests.

2. Normal levels of T4 can also be seen in Hyperthyroid patients with :T3 Thyrotoxicosis, Decreased binding capacity due to hypoproteinemia or ingestion of certain drugs (eg: phenytoin , salicylates).

3. Serum T4 levles in neonates and infants are higher than values in the normal adult , due to the increased concentration of TBG in neonate serum.

4. TSH may be normal in central hypothyroidism, recent rapid correction of hyperthyroidism or hypothroidism, pregnancy, phenytoin therapy.

TRIIODOTH	YRONINE (T3)	THYROXINE (T4)		THYROID STIMULATING HORMONE (TSH		
Age	Refferance Range (ng/mL)	Age	Refferance Range (μg/dL)	Age	Reference Range (μIU/mL)	
0 - 7 Days	0.20 - 2.65	0 - 7 Days	5.90 - 18.58	0 - 7 Days	2.43 - 24.3	
7 Days - 3 Months	0.36 - 2.59	7 Days - 3 Months	6.39 - 17.66	7 Days - 3 Months	0.58 - 11.00	
3 - 6 Months	0.51 - 2.52	3 - 6 Months	6.75 - 17.04	3 Days – 6 Months	0.70 - 8.40	





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)





🔽 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. BALWINDER KAUR		
AGE/ GENDER	: 66 YRS/FEMALE	PATIENT ID	: 1549115
COLLECTED BY	:	REG. NO./LAB NO.	: 122407150001
REFERRED BY	:	REGISTRATION DATE	: 15/Jul/2024 08:47 AM
BARCODE NO.	: 12503595	COLLECTION DATE	: 15/Jul/2024 09:12AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	REPORTING DATE	: 15/Jul/2024 05:13PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - H	ARYANA	

Test Name			Value	Unit		Biolog	ical Reference interval
6 - 12 Months	0.74 - 2.40	6 - 12 Months	7.10 - 16.16	6 – 12 Months	0.70 - 7.00		
1 - 10 Years	0.92 - 2.28	1 - 10 Years	6.00 - 13.80	1 – 10 Years	0.60 - 5.50		•
11- 19 Years	0.35 - 1.93	11 - 19 Years	4.87- 13.20	11 – 19 Years	0.50 - 5.50		•
> 20 years (Adults)	0.35 - 1.93	> 20 Years (Adults)	4.87 - 12.60	> 20 Years (Adults)	0.35-5.50		
	RECON	/IMENDATIONS OF TSH L	EVELS DURING PRE	GNANCY (μIU/mL)			
	1st Trimester			0.10 - 2.50			
	2nd Trimester			0.20 - 3.00			1
	3rd Trimester			0.30 - 4.10			

INCREASED TSH LEVELS:

1.Primary or untreated hypothyroidism may vary from 3 times to more than 100 times normal depending upon degree of hypofunction.

2.Hypothyroid patients receiving insufficient thyroid replacement therapy.

3. Hashimotos thyroiditis

4.DRUGS: Amphetamines, idonie containing agents & dopamine antagonist.

5.Neonatal period, increase in 1st 2-3 days of life due to post-natal surge

DECREASED TSH LEVELS:

1.Toxic multi-nodular goitre & Thyroiditis.

2. Over replacement of thyroid harmone in treatment of hypothyroidism.

3. Autonomously functioning Thyroid adenoma

4.Secondary pituatary or hypothalmic hypothyroidism

5. Acute psychiatric illness

6.Severe dehydration.

7.DRUGS: Glucocorticoids, Dopamine, Levodopa, T4 replacement therapy, Anti-thyroid drugs for thyrotoxicosis.

8. Pregnancy: 1st and 2nd Trimester





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



A PIONEER DIAGNOSTIC CENTRE

【 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

AGE/ GENDER	: Mrs. BALWINDER	KAUR			
	: 66 YRS/FEMALE	P	ATIENT ID	: 1549115	
COLLECTED BY	:	R	EG. NO./LAB NO.	: 122407150001	
REFERRED BY	:	R	EGISTRATION DATE	: 15/Jul/2024 08:47 AM	
BARCODE NO.	: 12503595		OLLECTION DATE	: 15/Jul/2024 09:12AM	
CLIENT CODE.	: P.K.R JAIN HEALTH		EPORTING DATE	: 15/Jul/2024 04:50PM	
CLIENT ADDRESS		ROAD, AMBALA CITY - HARY			
Test Name		Value	Unit	Biological Reference interval	
		VITA	MINS		
		VITAMIN D/25 HYD	DROXY VITAMIN D3		
	ROXY VITAMIN D3): S SESCENCE IMMUNOASS		ng/mL	DEFICIENCY: < 20.0 INSUFFICIENCY: 20.0 - 30.0 SUFFICIENCY: 30.0 - 100.0 TOXICITY: > 100.0	
INTERPRETATION:		00			
DEFIC INSUFF		< 20 21 - 29			
PREFFERE		30 - 100		ng/mL ng/mL	
INTOXIC			> 100 ng/mL gocalciferol (from plants, Vitamin D2), or cholecalciferol (from animals,		
tissue and tightly bou 3.Vitamin D plays a pro- phosphate reabsorpti 4.Severe deficiency m DECREASED: 1.Lack of sunshine exr 2.Inadequate intake, 3.Depressed Hepatic M 4.Secondary to advan	nd by a transport pro rimary role in the mai on, skeletal calcium d ay lead to failure to n posure. malabsorption (celiac Vitamin D 25- hydroxy ced Liver disease econdary Hyperparath	tein while in circulation. Intenance of calcium homeos eposition, calcium mobilization ineralize newly formed osteon disease) lase activity roidism (Mild to Moderate do	tatis. It promotes calcium on, mainly requlated by p oid in bone, resulting in ri eficiency)	port form of Vitamin D, being stored in adipos n absorption, renal calcium absorption and arathyroid harmone (PTH). ickets in children and osteomalacia in adults.	





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS , MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST



A PIONEER DIAGNOSTIC CENTRE

💟 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

AGE/ GENDER	: Mrs. BALWINDER KAUR			
IGE/ GENDEK	: 66 YRS/FEMALE	PAT	IENT ID	: 1549115
COLLECTED BY	:	REG	. NO./LAB NO.	: 122407150001
REFERRED BY	:	REG	ISTRATION DATE	: 15/Jul/2024 08:47 AM
BARCODE NO.			LECTION DATE	: 15/Jul/2024 09:12AM
CLIENT CODE.			ORTING DATE	: 16/Jul/2024 09:11AM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD,	AMBALA CITY - HARYA	NA	
Test Name		Value	Unit	Biological Reference inter
		VITAMIN B12/C	OBALAMIN	
VITAMIN B12/COBA		VITAMIN B12/C >2000.0 ^H	OBALAMIN pg/mL	200.0 - 1100.0
	LAMIN: SERUM NESCENT MICROPARTICLE			200.0 - 1100.0
by CMIA (CHEMILUMIN MMUNOASSAY) INTERPRETATION:-	NESCENT MICROPARTICLE		pg/mL	
by CMIA (CHEMILUMIN MMUNOASSAY) I <u>NTERPRETATION:-</u> INCREAS	NESCENT MICROPARTICLE	>2000.0 ^H		
by CMIA (CHEMILUMII MMUNOASSAY) INTERPRETATION:- INCREAS 1.Ingestion of Vitam	NESCENT MICROPARTICLE ED VITAMIN B12 nin C	>2000.0 ^H	pg/mL DECREASED VITAMIN	IB12
by CMIA (CHEMILUMII MMUNOASSAY) INTERPRETATION:- INCREAS 1.Ingestion of Vitam 2.Ingestion of Estrog	NESCENT MICROPARTICLE ED VITAMIN B12 hin C gen	>2000.0 ^H	pg/mL DECREASED VITAMIN irin, Anti-convulsants,	IB12
by CMIA (CHEMILUMIN MMUNOASSAY) INTERPRETATION:- INCREAS 1.Ingestion of Vitam 3.Ingestion of Vitam	NESCENT MICROPARTICLE ED VITAMIN B12 nin C gen nin A	>2000.0 ^H	pg/mL DECREASED VITAMIN irin, Anti-convulsants, stion	IB12
by CMIA (CHEMILUMIN MMUNOASSAY) INTERPRETATION:- INCREAS 1.Ingestion of Vitam 2.Ingestion of Vitam 3.Ingestion of Vitam 4.Hepatocellular inj	NESCENT MICROPARTICLE ED VITAMIN B12 nin C gen nin A jury	>2000.0 ^H	pg/mL DECREASED VITAMIN irin, Anti-convulsants, stion tive Harmones	IB12
by CMIA (CHEMILUMIN MMUNOASSAY) INTERPRETATION:- INCREAS 1.Ingestion of Vitam 3.Ingestion of Vitam	NESCENT MICROPARTICLE ED VITAMIN B12 nin C gen nin A jury	>2000.0 ^H	pg/mL DECREASED VITAMIN irrin, Anti-convulsants, stion tive Harmones ysis	IB12

6.Serum methylmalonic acid and homocysteine levels are also elevated in vitamin B12 deficiency states.

7.Follow-up testing for antibodies to intrinsic factor (IF) is recommended to identify this potential cause of vitamin B12 malabsorption. **NOTE:**A normal serum concentration of vitamin B12 does not rule out tissue deficiency of vitamin B12. The most sensitive test for vitamin B12 deficiency at the cellular level is the assay for MMA. If clinical symptoms suggest deficiency, measurement of MMA and homocysteine should be considered, even if serum vitamin B12 concentrations are normal.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)





A PIONEER DIAGNOSTIC CENTRE

【 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. BALWINDER KAUR			
AGE/ GENDER	: 66 YRS/FEMALE	РАТ	TENT ID	: 1549115
COLLECTED BY	:	REG	. NO./LAB NO.	: 122407150001
REFERRED BY	:	REG	ISTRATION DATE	: 15/Jul/2024 08:47 AM
BARCODE NO.	: 12503595	COL	LECTION DATE	: 15/Jul/2024 09:12AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INST	TITUTE REP	ORTING DATE	: 15/Jul/2024 05:19PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AM			
Test Name		Value	Unit	Biological Reference interv
		CLINICAL PAT	THOLOGY	
	URINE RC	DUTINE & MICROS	SCOPIC EXAMINAT	ION
PHYSICAL EXAMINAT	ION			
QUANTITY RECIEVED		10	ml	
	ANCE SPECTROPHOTOMETRY			
COLOUR	ANCE SPECTROPHOTOMETRY	AMBER YELLO	<i>I</i> V	PALE YELLOW
TRANSPARANCY	ANCE SI ECHICI HOTOMETRI	HAZY		CLEAR
	ANCE SPECTROPHOTOMETRY			
SPECIFIC GRAVITY		1.01		1.002 - 1.030
by DIP STICK/REFLECT	TANCE SPECTROPHOTOMETRY			
	<u>IION</u>			
REACTION	ANCE SPECTROPHOTOMETRY	ACIDIC		
PROTEIN		Negative		NEGATIVE (-ve)
	ANCE SPECTROPHOTOMETRY	noguno		
SUGAR		Negative		NEGATIVE (-ve)
-	TANCE SPECTROPHOTOMETRY			
pH by DIP STICK/REFLECT	ANCE SPECTROPHOTOMETRY	6.5		5.0 - 7.5
BILIRUBIN		Negative		NEGATIVE (-ve)
-	ANCE SPECTROPHOTOMETRY	-		
		Negative		NEGATIVE (-ve)
UROBILINOGEN	ANCE SPECTROPHOTOMETRY.	Normal	EU/dL	0.2 - 1.0
	ANCE SPECTROPHOTOMETRY	Norma	LO/ GL	0.2 1.0
KETONE BODIES		Negative		NEGATIVE (-ve)
•	TANCE SPECTROPHOTOMETRY	Nogetive		
BLOOD by DIP STICK/REFLECT	ANCE SPECTROPHOTOMETRY	Negative		NEGATIVE (-ve)
ASCORBIC ACID		NEGATIVE (-ve))	NEGATIVE (-ve)
	TANCE SPECTROPHOTOMETRY			、 <i>`</i>
MICROSCOPIC EXAM	INATION			



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS , MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST



A PIONEER DIAGNOSTIC CENTRE

🔽 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. BALWINDER KAUR			
AGE/ GENDER	: 66 YRS/FEMALE	PATIEN	T ID	: 1549115
COLLECTED BY	:	REG. NO)./LAB NO.	: 122407150001
REFERRED BY	:	REGIST	RATION DATE	: 15/Jul/2024 08:47 AM
BARCODE NO.	: 12503595	COLLEC	TION DATE	: 15/Jul/2024 09:12AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTIT	TUTE REPOR	FING DATE	: 15/Jul/2024 05:19PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA			
Test Name		Value	Unit	Biological Reference interval
RED BLOOD CELLS (R	BCs) Sentrieuged urinary sediment	NEGATIVE (-ve)	/HPF	0 - 3

by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT			
PUS CELLS	15-20	/HPF	0 - 5
by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT			
EPITHELIAL CELLS	3-4	/HPF	ABSENT
by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT			
CRYSTALS	NEGATIVE (-ve)		NEGATIVE (-ve)
by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT			
CASTS	NEGATIVE (-ve)		NEGATIVE (-ve)
by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT			
BACTERIA	NEGATIVE (-ve)		NEGATIVE (-ve)
by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT			
OTHERS	NEGATIVE (-ve)		NEGATIVE (-ve)
by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT			
TRICHOMONAS VAGINALIS (PROTOZOA)	ABSENT		ABSENT

by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT

DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS , MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST





TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT.

PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana) A PIONEER DIAGNOSTIC CENTRE

【 0171-2532620, 8222896961 ☑ pkrjainhealthcare@gmail.com

NAME	: Mrs. BALWINDER KAUR				
AGE/ GENDER	: 66 YRS/FEMALE		PATIENT ID	: 1549115	
COLLECTED BY	:		REG. NO./LAB NO.	: 122407150001	
REFERRED BY	:		REGISTRATION DATE	: 15/Jul/2024 08:47 AM	
BARCODE NO.	: 12503595		COLLECTION DATE	: 15/Jul/2024 09:12AM	
CLIENT CODE.	: P.K.R JAIN HEALTHCARE IN	NSTITUTE	REPORTING DATE	: 17/Jul/2024 11:17AM	
CLIENT ADDRESS			ARYANA		
Test Name		Value	Unit	Biological Reference interval	
		MICR	OBIOLOGY		
		BIC BACTERIA	AND ANTIBIOTIC SENSIT	FIVITY: URINE	
CULTURE AND SUSC	EPTIBILITY: URINE				
DATE OF SAMPLE		15-07-20)24		
SPECIMEN SOURCE		URINE			
INCUBATION PERIO		48 HOU	()		
GRAM STAIN		GRAM N	IEGATIVE (-ve)		
by MICROSCOPY		POSITIV			
CULTURE by AUTOMATED BROTH CULTURE		PUSITIV			
ORGANISM		ESCHER	CHIA COLI (E.COLI)		
by AUTOMATED BROT					
AEROBIC SUSCEPTIBILITY: URINE AMOXICILLIN+CLAVULANIC ACID by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 8/4 µg/mL		INTERM	EDIATE		
AMPICILLIN		RESISTA	NT		
	TH MICRODILUTION, CLSI ML	RESISTA			
AMPICILLIN+SULBA by AUTOMATED BRO Concentration: 8/4 μ	TH MICRODILUTION, CLSI	SENSITI	νe		
CHLORAMPHENICOI by AUTOMATED BRO Concentration: 8 µg/I	TH MICRODILUTION, CLSI	SENSITI	VE		
CIPROFLOXACIN <i>by AUTOMATED BRO</i> Concentration: 1 μg/ι	TH MICRODILUTION, CLSI ML	SENSITI	VE		
DOXYCYCLINE		SENSITI	VE		
	DR.VINAY CHOPRA	 DR.YI	GAM CHOPRA		

NOT VALID FOR MEDICO LEGAL PURPOSE

CONSULTANT PATHOLOGIST

MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS , MD (PATHOLOGY)

440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600, REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)

CONSULTANT PATHOLOGIST



NAME	: Mrs. BALWINDER KAUR				
AGE/ GENDER	: 66 YRS/FEMALE	P	ATIENT ID	: 1549115	
COLLECTED BY	:	R	EG. NO./LAB NO.	: 122407150001	
REFERRED BY	:	R	EGISTRATION DATE	: 15/Jul/2024 08:47 AM	
BARCODE NO.	: 12503595	C	OLLECTION DATE	: 15/Jul/2024 09:12AM	
CLIENT CODE.	: P.K.R JAIN HEALTHCARE IN	ISTITUTE R	EPORTING DATE	: 17/Jul/2024 11:17AM	
), AMBALA CITY - HARYANA			
Test Name		Value	Unit	Biological Reference interval	
<i>by AUTOMATED BROTI</i> Concentration: 4 μg/m	H MICRODILUTION, CLSI				
NALIDIXIC ACID		SENSITIVE			
	H MICRODILUTION, CLSI				
Concentration: 16 µg/r	nL				
GENTAMICIN		SENSITIVE			
<i>by AUTOMATED BROTI</i> Concentration: 16 μg/r	H MICRODILUTION, CLSI				
		RESISTANT			
<i>by AUTOMATED BROTH</i> Concentration: 16 μg/r					
NORFLOXACIN	H MICRODILUTION, CLSI	SENSITIVE			
Concentration: 4 µg/m					
MINOCYCLINE		SENSITIVE			
by AUTOMATED BROTH	H MICRODILUTION, CLSI	JEINJITTVE			
Concentration: 4 µg/m	L				
TOBRAMYCIN		SENSITIVE			
	H MICRODILUTION, CLSI				
Concentration: 4 µg/m	L				
AMIKACIN		SENSITIVE			
<i>by AUTOMATED BROTI</i> Concentration: 16 μg/r	H MICRODILUTION, CLSI nl				
AZETREONAM	H MICRODILUTION, CLSI	SENSITIVE			
Concentration: 4 µg/m	-				
CEFAZOLIN		INTERMEDI	ATF		
	H MICRODILUTION, CLSI				
Concentration: 16 µg/r	nL				
สสมระเทศ		4			
	AL.	(he	disa		
	an	Y			
		T			
	DR.VINAY CHOPRA CONSULTANT PATHOLOGIST		A CHOPRA		

440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600, REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)

MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS , MD (PATHOLOGY)



A PIONEER DIAGNOSTIC CENTRE

【 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

AGE/ GENDER : 66 YES/FEMALE PATIENT ID :: 1549115 COLLECTED BY : REG. NO./LAB NO. :: 122407150001 REFERRED BY : REGISTRATION DATE :: 15/Jul/2024 08:47 AM BARCODE NO. :: 12503595 COLLECTION DATE :: 15/Jul/2024 08:47 AM CLIENT CODE. : P.K.R. JAIN HEALTHCARE INSTITUTE REPORTING DATE :: 15/Jul/2024 09:12AM CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA :: 17/Jul/2024 09:12AM :: 17/Jul/2024 09:12AM Test Name Value Unit Biological Refe :: 17/Jul/2024 09:12AM CEFIXIME internetion: 6.100 INTERMEDIATE :: 17/Jul/2024 09:12AM Devalue Unit Biological Refe :: 17/Jul/2024 09:12AM CEFIXIME Value Unit Biological Refe by AUTOMATED BROTH MICRODILUTION, CLSI SENSITIVE :: SENSITIVE by AUTOMATED BROTH MICRODILUTION, CLSI SENSITIVE ::	
REFEREND BY :: REGISTRATION DATE :15/Jul/2024 08:47 AM BARCODE NO. :12503595 COLLECTION DATE :15/Jul/2024 09:12AM CLIENT CODE :: P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE :15/Jul/2024 11:17AM CLIENT ADDRESS :: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA :17/Jul/2024 11:17AM CEFININE Value Unit Biological Refe by AUTOMATED BROTH MICRODILUTION, CLSI INTERMEDIATE : CEFTAIDINE by AUTOMATED BROTH MICRODILUTION, CLSI RESISTANT COCCENTRATION: 8 µg/mL SENSITIVE : CEFTRIAXONE SENSITIVE : by AUTOMATED BROTH MICRODILUTION, CLSI SENSITIVE COCCENTRATION: 64 µg/mL SENSITIVE by AUTOMATED BROTH MICRODILUTION, CLSI SENSITIVE Concentration: 64 µg/mL SENSITIVE by AUTOMATED BROTH MICRODILUTION, CLSI SENSITIVE Concentration: 64 µg/mL SENSITIVE by AUTOMATED BROTH MICRODILUTION, CLSI SENSITIVE Concentration: 2 µg/mL SENSITIVE PIPERACILLIN+TAZOBACTUM SENSITIVE by AUTOMATED BROTH MICRODILUTION, CLSI <td< th=""><th></th></td<>	
BARCODE NO. 12503595 COLLECTION DATE 15/Jul/2024 09:12AM CLIENT CODE P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE 17/Jul/2024 11:17AM CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA INTERMEDIATE 17/Jul/2024 11:17AM Test Name Value Unit Biological Refe CEFIXIME INTERMEDIATE RESISTANT by AUTOMATED BROTH MICRODILUTION, CLSI RESISTANT CEFTAZIDIME SENSITIVE by AUTOMATED BROTH MICRODILUTION, CLSI SENSITIVE COCHTRATION: 64 µg/mL SENSITIVE by AUTOMATED BROTH MICRODILUTION, CLSI SENSITIVE by AUT	
CLIENT CODE : P.K.R. JAIN HEALTHCARE INSTITUTE REPORTING DATE : 17/Jul/2024 11:17AM CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA Init Biological Refe Test Name Value Unit Biological Refe CEFIXIME INTERMEDIATE By AUTOMATED BROTH MICRODILUTION, CLSI INTERMEDIATE Dy AUTOMATED BROTH MICRODILUTION, CLSI RESISTANT SENSITIVE Dy AUTOMATED BROTH MICRODILUTION, CLSI SENSITIVE	
CLEENT ADDRESS : MASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA Test Name Value Unit Biological Refe CEFIXINE INTERMEDIATE by AUTOMATED BROTH MICRODILUTION, CLSI CEFTAZIDINE SENSITIVE by AUTOMATED BROTH MICRODILUTION, CLSI COMMATED BROTH MICRODILUTION, CLSI COMMATED BROTH MICRODILUTION, CLSI Dy AUTOMATED BROTH MICRODILUTION, CLSI COMMATED BROTH MICRODILUTION, CLSI Dy AUTOMATED BROTH MICRODILUTION, CLSI COMMATED BROTH MICRODILUTION, CLSI DY AUTOMATED BROTH MICRODILUTION, CLSI COMMATED BROTH MICRODILUTION, CLSI Concentration: 2 LIG/MIC PIPERACILLIN+TAZOBACTUM SENSITIVE SENSITIVE DY AUTOMATED BROTH MICRODILUTION, CLSI COMMATED BR	
Test Name Value Unit Biological Refer CEFIXINE by AUTOMATED BROTH MICRODILUTION, CLSI COncentration: 8 µg/mL INTERMEDIATE INTERMEDIATE CEFTAZIDIME by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 8 µg/mL RESISTANT RESISTANT CEFTAZIDIME by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 4 µg/mL SENSITIVE by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 64 µg/mL SENSITIVE SENSITIVE LEVOFLOXACIN by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 2 µg/mL SENSITIVE SENSITIVE SENSITIVE SENSITIVE ILEVOFLOXACIN by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 2 µg/mL SENSITIVE SENSITIVE SENSITIVE ILEVOFLOXACIN by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 8 µg/mL SENSITIVE SENSITIVE SENSITIVE ILEVOFLOXACIN by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 8 µg/mL SENSITIVE SENSITIVE ILEVOFLOXACIN by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 8 µg/mL SENSITIVE SENSITIVE ILEVOFLOXACIN by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 8 µg/mL SENSITIVE SENSITIVE	
CEFIXIME by AUTOMATED BROTH MICRODILUTION, CLSI INTERMEDIATE by AUTOMATED BROTH MICRODILUTION, CLSI CEFOXITIN by AUTOMATED BROTH MICRODILUTION, CLSI RESISTANT COncentration: 8 µg/mL SENSITIVE CEFTRIAXONE by AUTOMATED BROTH MICRODILUTION, CLSI SENSITIVE SENSITIVE COncentration: 4 µg/mL SENSITIVE SENSITIVE CEFTRIAXONE by AUTOMATED BROTH MICRODILUTION, CLSI SENSITIVE SENSITIVE COncentration: 64 µg/mL SENSITIVE SENSITIVE EEVOFLOXACIN by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 2 µg/mL SENSITIVE SENSITIVE NETLIMICIN SULPHATE by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 8 µg/mL SENSITIVE SENSITIVE SENSITIVE PIPERACILLIN+TAZOBACTUM by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 16/4 µg/mL SENSITIVE SENSITIVE PIPERACILLIN+TAZOBACTUM by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 16/4 µg/mL SENSITIVE TICARCILLIN+CLAVULANIC ACID SENSITIVE	
by AUTOMATED BROTH MICRODILUTION, CLSI CEFOXITIN by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 8 µg/mL CEFTAZIDIME by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 4 µg/mL CEFTRIAXONE by AUTOMATED BROTH MICRODILUTION, CLSI COncentration: 4 µg/mL CEFTRIAXONE by AUTOMATED BROTH MICRODILUTION, CLSI FOSFOMYCIN by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 64 µg/mL LEVOFLOXACIN by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 2 µg/mL NETLIMICIN SULPHATE by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 8 µg/mL PIPERACILLIN+TAZOBACTUM by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 8 µg/mL SENSITIVE by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 16/4 µg/mL SENSITIVE by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 16/4 µg/mL SENSITIVE by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 16/4 µg/mL SENSITIVE <td>erence interval</td>	erence interval
CEFOXITIN by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 8 µg/mL RESISTANT CEFTAZIDIME by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 4 µg/mL SENSITIVE SENSITIVE SENSITIVE SENSITIVE SENSITIVE SENSITIVE SENSITIVE SENSITIVE SENSITIVE SUPPRATED BROTH MICRODILUTION, CLSI Concentration: 64 µg/mL LEVOFLOXACIN by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 64 µg/mL SENSITIVE SENSITIVE SENSITIVE SENSITIVE SENSITIVE LEVOFLOXACIN by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 2 µg/mL SENSITIVE SENSITIVE SENSITIVE NETLIMICIN SULPHATE by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 8 µg/mL SENSITIVE SENSITIVE SENSITIVE PIPERACILLIN+TAZOBACTUM by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 16/4 µg/mL SENSITIVE SENSITIVE TICARCILLIN+CLAVULANIC ACID SENSITIVE	
by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 4 µg/mL CEFTRIAXONE by AUTOMATED BROTH MICRODILUTION, CLSI FOSFOMYCIN by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 64 µg/mL LEVOFLOXACIN by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 2 µg/mL NETLIMICIN SULPHATE by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 8 µg/mL PIPERACILLIN+TAZOBACTUM by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 16/4 µg/mL TICARCILLIN+CLAVULANIC ACID	
by AUTOMATED BROTH MICRODILUTION, CLSI FOSFOMYCIN by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 64 µg/mL LEVOFLOXACIN by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 2 µg/mL NETLIMICIN SULPHATE by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 8 µg/mL PIPERACILLIN+TAZOBACTUM by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 16/4 µg/mL TICARCILLIN+CLAVULANIC ACID SENSITIVE	
FOSFOMYCIN by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 64 μg/mL SENSITIVE LEVOFLOXACIN by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 2 μg/mL SENSITIVE NETLIMICIN SULPHATE by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 8 μg/mL SENSITIVE PIPERACILLIN+TAZOBACTUM by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 16/4 μg/mL SENSITIVE TICARCILLIN+CLAVULANIC ACID SENSITIVE	
by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 2 µg/mL NETLIMICIN SULPHATE by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 8 µg/mL PIPERACILLIN+TAZOBACTUM by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 16/4 µg/mL TICARCILLIN+CLAVULANIC ACID	
by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 8 µg/mL PIPERACILLIN+TAZOBACTUM by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 16/4 µg/mL TICARCILLIN+CLAVULANIC ACID SENSITIVE	
by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 16/4 µg/mL TICARCILLIN+CLAVULANIC ACID SENSITIVE	
Concentration: 16/2 µg/mL	
TRIMETHOPRIM+SULPHAMETHAZOLE SENSITIVE by AUTOMATED BROTH MICRODILUTION, CLSI SENSITIVE Concentration: 2/38 μg/mL Concentration	
CEFIPIME SENSITIVE by Automated Broth Microdilution, CLSI	
An groups	

DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS , MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST



NAME	: Mrs. BALWINDER KAUR			
AGE/ GENDER	: 66 YRS/FEMALE	PATIENT ID	: 1549115	
COLLECTED BY	:	REG. NO./LAB NO.	: 122407150001	
REFERRED BY	:	REGISTRATION DATE	: 15/Jul/2024 08:47 AM	
BARCODE NO.	: 12503595	COLLECTION DATE	: 15/Jul/2024 09:12AM	
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	REPORTING DATE	: 17/Jul/2024 11:17AM	
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA			
Test Name	Value	Unit	Biological Reference interval	
Concentration: 2 ug/	nl			

A PIONEER DIAGNOSTIC CENTRE ↓ 0171-2532620, 8222896961 ☑ pkrjainhealthcare@gmail.com

Concentration: 2 µg/mL SENSITIVE DORIPFNFM by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 1 µg/mL **INTERMEDIATE** IMIPINEM by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 1 µg/mL MEROPENEM SENSITIVE by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 1 µg/mL COLISTIN SENSITIVE by AUTOMATED BROTH MICRODILUTION, CLSI Concentration: 0.06 µg/mL

INTERPRETATION:

In urine culture and sensitivity, presence of more than 100,000 organism per mL in midstream sample of urine is considered clinically significant. However in symptomatic patients, a smaller number of bacteria (100 to 10000/mL) may signify infection.
 Colony count of 100 to 10000/ mL indicate infection, if isolate from specimen obtained by suprapubic aspiration or "in-and-out" catheterization or from patients with indwelling catheters.
 SUSCEPTIBILITY:

 A test interpreted as SENSTITIVE implies that infection due to isolate may be appropriately treated with the dosage of an antimicrobial agent recommended for that type of infection and infecting species, unless otherwise indicated..
 A test interpreted as INTERMEDIATE implies that the" Infection due to the isolate may be appropriately treated in body sites where the drugs are

A test interpreted as **INTERMEDIATE** implies that the "Infection due to the isolate may be appropriately treated in body sites where the drugs are physiologically concentrated or when a high dosage of drug can be used".
 A test interpreted as **RESISTANT** implies that the "isolates are not inhibited by the usually achievable concentration of the agents with normal and the dot in the dot is a second sec

3.A test interpreted as **RESISTANT** implies that the "isolates" are not inhibited by the usually achievable concentration of the agents with normal dosage, schedule and/or fall in the range where specific microbial resistance mechanism are likely (e.g. beta-lactamases), and clinical efficacy has not been reliable in treatment studies.

CAUTION:

Conditions which can cause a false Negative culture:

1. Patient is on antibiotics. Please repeat culture post therapy.

2. Anaerobic bacterial infection.

- 3. Fastidious aerobic bacteria which are not able to grow on routine culture media.
- 4. Besides all these factors, at least in 25-40 % of cases there is no direct correlation between in vivo clinical picture.
- 5. Renal tuberculosis to be confirmed by AFB studies.





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

*** End Of Report ***

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)



