



PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

REPORTING DATE

: 19/Jul/2024 03:12PM

NAME : Mr. C R ARORA

AGE/ GENDER : 82 YRS/MALE **PATIENT ID** : 1544171

COLLECTED BY REG. NO./LAB NO. : 122407190019

REFERRED BY **REGISTRATION DATE** : 19/Jul/2024 01:38 PM BARCODE NO. : 12503688 **COLLECTION DATE** : 19/Jul/2024 01:46PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

Value Unit **Biological Reference interval** Test Name

CLINICAL CHEMISTRY/BIOCHEMISTRY **GLUCOSE RANDOM (R)**

96.93 GLUCOSE RANDOM (R): PLASMA mg/dL NORMAL: < 140.00

by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD) PREDIABETIC: 140.0 - 200.0 DIABETIC: > OR = 200.0

CLIENT CODE.

IN ACCORDANCE WITH AMERICAN DIABETES ASSOCIATION GUIDELINES:

1. A random plasma glucose level below 140 mg/dl is considered normal.

2. A random glucose level between 140 - 200 mg/dl is considered as glucose intolerant or prediabetic. A fasting and post-prnadial blood test (after consumption of 75 gms of glucose) is recommended for all such patients.

3. A random glucose level of above 200 mg/dl is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for all such patients. A fasting plasma glucose level in excess of 125 mg/dl on both occasions is confirmatory for diabetic state.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



CLIENT CODE.



PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

REPORTING DATE

: 19/Jul/2024 03:12PM

NAME : Mr. C R ARORA

by URICASE - OXIDASE PEROXIDASE

AGE/ GENDER : 82 YRS/MALE **PATIENT ID** : 1544171

COLLECTED BY : 122407190019 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 19/Jul/2024 01:38 PM BARCODE NO. : 12503688 **COLLECTION DATE** : 19/Jul/2024 01:46PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

Test Name	Value	Unit	Biological Reference interval
	KIDNEY FUNCTION	I TEST (BASIC)	
UREA: SERUM by UREASE - GLUTAMATE DEHYDROGENASE (GLDH)	33.33	mg/dL	10.00 - 50.00
CREATININE: SERUM by ENZYMATIC, SPECTROPHOTOMETERY	0.94	mg/dL	0.40 - 1.40
BLOOD UREA NITROGEN (BUN): SERUM by CALCULATED, SPECTROPHOTOMETERY	15.57	mg/dL	7.0 - 25.0
BLOOD UREA NITROGEN (BUN)/CREATININE RATIO: SERUM by CALCULATED, SPECTROPHOTOMETERY	16.56	RATIO	10.0 - 20.0
UREA/CREATININE RATIO: SERUM by CALCULATED, SPECTROPHOTOMETERY	35.46	RATIO	
URIC ACID: SERUM	5.33	mg/dL	3.60 - 7.70



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)





PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

REPORTING DATE

: 19/Jul/2024 03:12PM

NAME : Mr. C R ARORA

AGE/ GENDER : 82 YRS/MALE **PATIENT ID** : 1544171

COLLECTED BY REG. NO./LAB NO. : 122407190019

REFERRED BY **REGISTRATION DATE** : 19/Jul/2024 01:38 PM BARCODE NO. **COLLECTION DATE** : 19/Jul/2024 01:46PM : 12503688

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

Test Name Value Unit **Biological Reference interval**

INTERPRETATION:

CLIENT CODE.

Normal range for a healthy person on normal diet: 12 - 20

To Differentiate between pre- and postrenal azotemia. INCREASED RATIO (>20:1) WITH NORMAL CREATININE:

1. Prerenal azotemia (BUN rises without increase in creatinine) e.g. heart failure, salt depletion, dehydration, blood loss) due to decreased glomerular filtration rate.

Ž.Catabolic states with increased tissue breakdown.

3.GI hemorrhage.

4. High protein intake.

5. Impaired renal function plus.

6. Excess protein intake or production or tissue breakdown (e.g. infection, GI bleeding, thyrotoxicosis, Cushings syndrome, high protein diet,

burns, surgery, cachexia, high fever)

7. Urine reabsorption (e.g. ureterocolostomy)
8. Reduced muscle mass (subnormal creatinine production)
9. Certain drugs (e.g. tetracycline, glucocorticoids)
INCREASED RATIO (>20:1) WITH ELEVATED CREATININE LEVELS:

1. Postrenal azotemia (BUN rises disproportionately more than creatinine) (e.g. obstructive uropathy).

2. Prerenal azotemia superimposed on renal disease.

DECREASED RATIO (<10:1) WITH DECREASED BUN:

1.Acute tubular necrosis.

2.Low protein diet and starvation.

3. Severe liver disease.

4. Other causes of decreased urea synthesis.

5. Repeated dialysis (urea rather than creatinine diffuses out of extracellular fluid).

6.Inherited hyperammonemias (urea is virtually absent in blood)

7.SIADH (syndrome of inappropiate antidiuretic harmone) due to tubular secretion of urea.

8. Pregnancy

DECREASED RATIO (<10:1) WITH INCREASED CREATININE:

- 1. Phenacimide therapy (accelerates conversion of creatine to creatinine).
- 2. Rhabdomyolysis (releases muscle creatinine).
- 3. Muscular patients who develop renal failure

INAPPROPIATE RATIO

1. Diabetic ketoacidosis (acetoacetate causes false increase in creatinine with certain methodologies, resulting in normal ratio when dehydration should produce an increased BUN/creatinine ratio).

2. Cephalosporin therapy (interferes with creatinine measurement).

*** End Of Report ***

DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)