

A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

NAME : Mrs. HARPREET

AGE/ GENDER : 50 YRS/FEMALE **PATIENT ID** : 1554971

COLLECTED BY REG. NO./LAB NO. : 122407200009

REFERRED BY **REGISTRATION DATE** : 20/Jul/2024 09:16 AM BARCODE NO. **COLLECTION DATE** : 20/Jul/2024 09:24AM : 12503699 CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 20/Jul/2024 12:32PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Value Unit Test Name **Biological Reference interval**

CLINICAL CHEMISTRY/BIOCHEMISTRY

LIPID PROFILE: BASIC

CHOLESTEROL TOTAL: SERUM mg/dL **OPTIMAL:** < 200.0 215.75H

by CHOLESTEROL OXIDASE PAP **BORDERLINE HIGH: 200.0 - 239.0** HIGH CHOLESTEROL: > OR = 240.0

TRIGLYCERIDES: SERUM 152.86H mg/dL **OPTIMAL:** < 150.0

by GLYCEROL PHOSPHATE OXIDASE (ENZYMATIC) **BORDERLINE HIGH: 150.0 - 199.0**

HIGH: 200.0 - 499.0 VERY HIGH: > OR = 500.0

HDL CHOLESTEROL (DIRECT): SERUM LOW HDL: < 30.0 53.05 mg/dL

by SELECTIVE INHIBITION BORDERLINE HIGH HDL: 30.0 -

60.0

 $HIGH\ HDL: > OR = 60.0$ LDL CHOLESTEROL: SERUM **OPTIMAL:** < 100.0 132.13^H mg/dL

by CALCULATED, SPECTROPHOTOMETRY ABOVE OPTIMAL: 100.0 - 129.0 **BORDERLINE HIGH: 130.0 - 159.0**

HIGH: 160.0 - 189.0 **VERY HIGH: > OR = 190.0**

NON HDL CHOLESTEROL: SERUM 162.7H **OPTIMAL: < 130.0** mg/dL by CALCULATED, SPECTROPHOTOMETRY ABOVE OPTIMAL: 130.0 - 159.0

BORDERLINE HIGH: 160.0 - 189.0 HIGH: 190.0 - 219.0

VERY HIGH: > OR = 220.0 VLDL CHOLESTEROL: SERUM 30.57 0.00 - 45.00mg/dL

by CALCULATED, SPECTROPHOTOMETRY TOTAL LIPIDS: SERUM 584.36 mg/dL 350.00 - 700.00

by CALCULATED, SPECTROPHOTOMETRY CHOLESTEROL/HDL RATIO: SERUM **RATIO** LOW RISK: 3.30 - 4.40 4.07

by CALCULATED, SPECTROPHOTOMETRY AVERAGE RISK: 4.50 - 7.0 MODERATE RISK: 7.10 - 11.0

HIGH RISK: > 11.0 LDL/HDL RATIO: SERUM 2.49 **RATIO** LOW RISK: 0.50 - 3.0



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)







A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

REPORTING DATE

NAME : Mrs. HARPREET

AGE/ GENDER : 50 YRS/FEMALE **PATIENT ID** : 1554971

COLLECTED BY REG. NO./LAB NO. : 122407200009

REFERRED BY **REGISTRATION DATE** : 20/Jul/2024 09:16 AM BARCODE NO. **COLLECTION DATE** : 20/Jul/2024 09:24AM : 12503699

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

Value Unit **Biological Reference interval Test Name**

by CALCULATED, SPECTROPHOTOMETRY MODERATE RISK: 3.10 - 6.0

HIGH RISK: > 6.0

: 20/Jul/2024 12:32PM

RATIO 3.00 - 5.00TRIGLYCERIDES/HDL RATIO: SERUM 2.88^L

by CALCULATED. SPECTROPHOTOMETRY INTERPRETATION:

CLIENT CODE.

1. Measurements in the same patient can show physiological analytical variations. Three serial samples 1 week apart are recommended for

Total Cholesterol, Triglycerides, HDL & LDL Cholesterol.

2. As per NLA-2014 guidelines, all adults above the age of 20 years should be screened for lipid status. Selective screening of children above the age of 2 years with a family history of premature cardiovascular disease or those with at least one parent with high total cholesterol is

recommended. 3. Low HDL levels are associated with increased risk for Atherosclerotic Cardiovascular disease (ASCVD) due to insufficient HDL being available

to participate in reverse cholesterol transport, the process by which cholesterol is eliminated from peripheral tissues.

4. NLA-2014 identifies Non HDL Cholesterol (an indicator of all atherogeniclipoproteins such as LDL, VLDL, IDL, Lpa, Chylomicron remnants) along with LDL-cholesterol as co-primary target for cholesterol lowering therapy. Note that major risk factors can modify treatment goals for LDL &Non

5. Additional testing for Apolipoprotein B, hsCRP,Lp(a) & LP-PLA2 should be considered among patients with moderate risk for ASCVD for risk refinement



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)





A PIONEER DIAGNOSTIC CENTRE

REPORTING DATE

: 20/Jul/2024 01:47PM

NAME : Mrs. HARPREET

AGE/ GENDER : 50 YRS/FEMALE **PATIENT ID** : 1554971

COLLECTED BY : 122407200009 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 20/Jul/2024 09:16 AM BARCODE NO. : 12503699 **COLLECTION DATE** : 20/Jul/2024 09:24AM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

Test Name Value Unit **Biological Reference interval**

ENDOCRINOLOGY

THYROID FUNCTION TEST: TOTAL

TRIIODOTHYRONINE (T3): SERUM 1.244 ng/mL 0.35 - 1.93

by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

THYROXINE (T4): SERUM 8.54 4.87 - 12.60 μgm/dL

by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

THYROID STIMULATING HORMONE (TSH): SERUM 3.044 μIU/mL 0.35 - 5.50

by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

3rd GENERATION, ULTRASENSITIVE

INTERPRETATION:

CLIENT CODE.

TSH levels are subject to circadian variation, reaching peak levels between 2-4 a.m and at a minimum between 6-10 pm. The variation is of the order of 50%. Hence time of the day has influence on the measured serum TSH concentrations. TSH stimulates the production and secretion of the metabolically active hormones, thyroxine (T4) and trilodothyronine (T3). Failure at any level of regulation of the hypothalamic-pituitary-thyroid axis will result in either underproduction (hypothyroidism) or overproduction(hyperthyroidism) of T4 and/or T3.

CLINICAL CONDITION	Т3	T4	TSH	
Primary Hypothyroidism:	Reduced	Reduced	Increased (Significantly)	
Subclinical Hypothyroidism:	Normal or Low Normal	Normal or Low Normal	High	
Primary Hyperthyroidism:	Increased	Increased	Reduced (at times undetectable)	
Subclinical Hyperthyroidism:	Normal or High Normal	Normal or High Normal	Reduced	

- 1. T3 and T4 circulates in reversibly bound form with Thyroid binding globulins (TBG), and to a lesser extent albumin and Thyroid binding Pre Albumin so conditions in which TBG and protein levels alter such as pregnancy, excess estrogens, androgens, anabolic steroids and glucocorticoids may falsely affect the T3 and T4 levels and may cause false thyroid values for thyroid function tests.
- 2. Normal levels of T4 can also be seen in Hyperthyroid patients with :T3 Thyrotoxicosis, Decreased binding capacity due to hypoproteinemia or ingestion of certain drugs
- 3. Serum T4 levies in neonates and infants are higher than values in the normal adult, due to the increased concentration of TBG in neonate serum.
- 4. TSH may be normal in central hypothyroidism, recent rapid correction of hyperthyroidism or hypothroidism, pregnancy, phenytoin therapy.

TRIIODOTHYRONINE (T3)		THYROXINE (T4)		THYROID STIMULATING HORMONE (TSH)		
Age	Refferance Range (ng/mL)	Age	Refferance Range (µg/dL)	Age	Reference Range (μΙU/mL)	
0 - 7 Days	0.20 - 2.65	0 - 7 Days	5.90 - 18.58	0 - 7 Days	2.43 - 24.3	
7 Days - 3 Months	0.36 - 2.59	7 Days - 3 Months	6.39 - 17.66	7 Days - 3 Months	0.58 - 11.00	
3 - 6 Months	0.51 - 2.52	3 - 6 Months	6.75 – 17.04	3 Days – 6 Months	0.70 - 8.40	



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)





A PIONEER DIAGNOSTIC CENTRE

NAME : Mrs. HARPREET

AGE/ GENDER : 50 YRS/FEMALE **PATIENT ID** : 1554971

COLLECTED BY : 122407200009 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 20/Jul/2024 09:16 AM BARCODE NO. : 12503699 **COLLECTION DATE** : 20/Jul/2024 09:24AM

CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 20/Jul/2024 01:47PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name		Value	Unit		Biological Reference interval		
6 - 12 Months	0.74 - 2.40	6 - 12 Months	7.10 – 16.16	6 – 12 Months	0.70 - 7.00		
1 - 10 Years	0.92 - 2.28	1 - 10 Years	6.00 - 13.80	1 – 10 Years	0.60 - 5.50		
11- 19 Years	0.35 - 1.93	11 - 19 Years	4.87- 13.20	11 - 19 Years	0.50 - 5.50		
> 20 years (Adults)	0.35 - 1.93	> 20 Years (Adults)	4.87 - 12.60	> 20 Years (Adults)	0.35- 5.50		
	RECOM	MENDATIONS OF TSH LE	VELS DURING PREC	NANCY (μIU/mL)			
1st Trimester			0.10 – 2.50				1
2nd Trimester			0.20 - 3.00				1
3rd Trimester			0.30 - 4.10				

INCREASED TSH LEVELS:

- 1. Primary or untreated hypothyroidism may vary from 3 times to more than 100 times normal depending upon degree of hypofunction.
- 2. Hypothyroid patients receiving insufficient thyroid replacement therapy.
- 3. Hashimotos thyroiditis
- 4.DRUGS: Amphetamines, idonie containing agents & dopamine antagonist.
- 5. Neonatal period, increase in 1st 2-3 days of life due to post-natal surge

DECREASED TSH LEVELS:

- 1.Toxic multi-nodular goitre & Thyroiditis.
- 2. Over replacement of thyroid harmone in treatment of hypothyroidism.
- 3. Autonomously functioning Thyroid adenoma
- 4. Secondary pituatary or hypothalmic hypothyroidism
- 5. Acute psychiatric illness
- 6. Severe dehydration.
- 7.DRUGS: Glucocorticoids, Dopamine, Levodopa, T4 replacement therapy, Anti-thyroid drugs for thyrotoxicosis.

8. Pregnancy: 1st and 2nd Trimester

*** End Of Report ***



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)