

PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

💟 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. GITA ARORA					
AGE/ GENDER	: 38 YRS/FEMALE		PATIENT ID	: 1360289		
COLLECTED BY	:		REG. NO./LAB NO.	: 122407270013		
REFERRED BY	:		REGISTRATION DATE	: 27/Jul/2024 10:54 AM		
BARCODE NO.	: 12503840		COLLECTION DATE	: 27/Jul/2024 11:29AM		
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITU	JTE	REPORTING DATE	: 27/Jul/2024 05:02PM		
CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA						
Test Name		Value	Unit	Biological Reference interval		
		ENDOCI	RINOLOGY			
	710/					
	IHYI	ROID FUNC	TION TEST: TOTAL			
TRIIODOTHYRONINI	E (T3): SERUM vescent microparticle immunoassay	1.047	ng/mL	0.35 - 1.93		
THYROXINE (T4): SERUM 7.56 by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)			μgm/dL	4.87 - 12.60		
	ING HORMONE (TSH): SERUM NESCENT MICROPARTICLE IMMUNOASSAY	2.423	µIU/mL	0.35 - 5.50		

INTERPRETATION:

TSH levels are subject to circadian variation, reaching peak levels between 2-4 a.m and at a minimum between 6-10 pm. The variation is of the order of 50%. Hence time of the day has influence on the measured serum TSH concentrations. TSH stimulates the production and secretion of the metabolically active hormones, thyroxine (T4) and trilodothyronine (T3). Failure at any level of regulation of the hypothalamic-pituitary-thyroid axis will result in either underproduction (hypothyroidism) or overproduction(hyperthyroidism) of T4 and/or T3.

CLINICAL CONDITION	T3	T4	TSH	
Primary Hypothyroidism:	Reduced	Reduced	Increased (Significantly)	
Subclinical Hypothyroidism:	Normal or Low Normal	Normal or Low Normal	High	
Primary Hyperthyroidism:	Increased	Increased	Reduced (at times undetectable)	
Subclinical Hyperthyroidism:	Normal or High Normal	Normal or High Normal	Reduced	

LIMITATIONS:-

1. T3 and T4 circulates in reversibly bound form with Thyroid binding globulins (TBG), and to a lesser extent albumin and Thyroid binding Pre Albumin so conditions in which TBG and protein levels alter such as pregnancy, excess estrogens, androgens, anabolic steroids and glucocorticoids may falsely affect the T3 and T4 levels and may cause false thyroid values for thyroid function tests.

2. Normal levels of T4 can also be seen in Hyperthyroid patients with :T3 Thyrotoxicosis, Decreased binding capacity due to hypoproteinemia or ingestion of certain drugs (eg: phenytoin , salicylates).

3. Serum T4 levles in neonates and infants are higher than values in the normal adult , due to the increased concentration of TBG in neonate serum.

4. TSH may be normal in central hypothyroidism, recent rapid correction of hyperthyroidism or hypothroidism, pregnancy, phenytoin therapy.

TRIIODOTHYRONINE (T3)		THYROXINE (T4)		THYROID STIMULATING HORMONE (TSH)		
Age	Refferance Range (ng/mL)	Age	Refferance Range (µg/dL)	Age	Reference Range (μIU/mL)	
0-7 Days	0.20 - 2.65	0 - 7 Days	5.90 - 18.58	0 - 7 Days	2.43 - 24.3	
7 Days - 3 Months	0.36 - 2.59	7 Days - 3 Months	6.39 - 17.66	7 Days - 3 Months	0.58 - 11.00	
3 - 6 Months	0.51 - 2.52	3 - 6 Months	6.75 - 17.04	3 Days – 6 Months	0.70 - 8.40	





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)

440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)



PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana) A PIONEER DIAGNOSTIC CENTRE

└ 0171-2532620, 8222896961 **□** pkrjainhealthcare@gmail.com

NAME	: Mrs. GITA ARORA			
AGE/ GENDER	: 38 YRS/FEMALE	PATIENT ID	: 1360289	
COLLECTED BY	:	REG. NO./LAB NO.	: 122407270013	
REFERRED BY	:	REGISTRATION DATE	: 27/Jul/2024 10:54 AM	
BARCODE NO.	: 12503840	COLLECTION DATE	: 27/Jul/2024 11:29AM	
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	REPORTING DATE	: 27/Jul/2024 05:02PM	
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA			

Test Name		Value	Value Unit		Biological Reference interval		
6 - 12 Months	0.74 - 2.40	6 - 12 Months	7.10 - 16.16	6 – 12 Months	0.70 - 7.00		
1 - 10 Years	0.92 - 2.28	1 - 10 Years	6.00 - 13.80	1 – 10 Years	0.60 - 5.50		
11- 19 Years	0.35 - 1.93	11 - 19 Years	4.87-13.20	11 – 19 Years	0.50 - 5.50		
> 20 years (Adults)	0.35 - 1.93	> 20 Years (Adults)	4.87 - 12.60	> 20 Years (Adults)	0.35- 5.50		
	RECOM	MENDATIONS OF TSH LE	VELS DURING PREC	GNANCY (µIU/mL)			
1st Trimester			0.10 - 2.50				
2nd Trimester			0.20 - 3.00				
3rd Trimester			0.30 - 4.10				

INCREASED TSH LEVELS:

1.Primary or untreated hypothyroidism may vary from 3 times to more than 100 times normal depending upon degree of hypofunction.

2.Hypothyroid patients receiving insufficient thyroid replacement therapy.

3.Hashimotos thyroiditis

4.DRUGS: Amphetamines, idonie containing agents & dopamine antagonist.

5.Neonatal period, increase in 1st 2-3 days of life due to post-natal surge

DECREASED TSH LEVELS:

1.Toxic multi-nodular goitre & Thyroiditis.

2. Over replacement of thyroid harmone in treatment of hypothyroidism.

3. Autonomously functioning Thyroid adenoma

4.Secondary pituatary or hypothalmic hypothyroidism

5. Acute psychiatric illness

6.Severe dehydration.

7.DRUGS: Glucocorticoids, Dopamine, Levodopa, T4 replacement therapy, Anti-thyroid drugs for thyrotoxicosis.

8. Pregnancy: 1st and 2nd Trimester

*** End Of Report ***





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)

440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600, REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)

