**PKR JAIN HEALTHCARE INSTITUTE** NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

【 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. NITIKA					
AGE/ GENDER	: 23 YRS/FEMALE	PAT	IENT ID	: 1568540		
COLLECTED BY	:	REG.	NO./LAB NO.	: 122408020013		
REFERRED BY	RRED BY : REGISTRATION DAT		STRATION DATE	E : 02/Aug/2024 02:27 PM		
BARCODE NO.	: 12503954	COLL	LECTION DATE	: 02/Aug/2024 02:38PM		
CLIENT CODE.	: P.K.R JAIN HEALTHCARE	INSTITUTE <b>REP</b>	DRTING DATE	: 02/Aug/2024 04:45PM		
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD	), AMBALA CITY - HARYAN	A			
Test Name		Value	Unit	Biological Reference interval		
	CL	INICAL CHEMISTRY	/BIOCHEMISTR	Y		
		URIC AC	ID			
URIC ACID: SERUM		6.07	mg/dL	2.50 - 6.80		
1.GOUT occurs when 2.Uric Acid is the end ntestinal tract by mi <b>NCREASED:-</b> (A).DUE TO INCREASEI 1.Idiopathic primary pu 2.Excessive dietary pu 3.Cytolytic treatment 4.Polycythemai vera	high levels of Uric Acid in th product of purine metabolis crobial degradation. D <b>PRODUCTION:-</b>	m . Uric acid is excreted to	orm & accumulate ard a large degree by the	ound a joint. kidneys and to a smaller degree in the		
INTERPRETATION:- 1.GOUT occurs when 2.Uric Acid is the end Intestinal tract by mi- INCREASED:- (A).DUE TO INCREASED 1.Idiopathic primary pu 3.Cytolytic treatment 4.Polycythemai vera 5.Psoriasis. 6.Sickle cell anaemia (B).DUE TO DECREASE 1.Alcohol ingestion. 2.Thiazide diuretics. 3.Lactic acidosis. 4.Aspirin ingestion (lef 5.Diabetic ketoacidosi	high levels of Uric Acid in the product of purine metabolisi crobial degradation. <b>D PRODUCTION:-</b> gout. urines (organ meats, legumes, of malignancies especially le & myeloid metaplasia. etc. <b>D EXCREATION (BY KIDNEYS)</b> ess than 2 grams per day ). sis or starvation.	m . Uric acid is excreted to	orm & accumulate ard a large degree by the	bund a joint. I kidneys and to a smaller degree in the		
INTERPRETATION:- 1. GOUT occurs when 2. Uric Acid is the end Intestinal tract by mi- INCREASED:- (A).DUE TO INCREASED 1. Idiopathic primary full 2. Excessive dietary pull 3. Cytolytic treatment 4. Polycythemai vera for 5. Sickle cell anaemia (B).DUE TO DECREASE 1. Alcohol ingestion. 2. Thiazide diuretics. 3. Lactic acidosis. 4. Aspirin ingestion (lef 5. Diabetic ketoacidos 5. Renal failure due to DECREASED:- (A).DUE TO DIETARY D	high levels of Uric Acid in the product of purine metabolist crobial degradation. <b>D PRODUCTION:-</b> gout. of malignancies especially le & myeloid metaplasia. etc. <b>D EXCREATION (BY KIDNEYS)</b> ess than 2 grams per day ). sis or starvation. any cause etc. <b>DEFICIENCY</b>	m . Uric acid is excreted to ,anchovies, etc). eukemais & lymphomas.	orm & accumulate ard a large degree by the	bund a joint. Ridneys and to a smaller degree in the		
INTERPRETATION:- 1. GOUT occurs when 2. Uric Acid is the end intestinal tract by mi- INCREASED:- (A).DUE TO INCREASED 1. Idiopathic primary fu 3. Cytolytic treatment 4. Polycythemai vera 5. Psoriasis. 6. Sickle cell anaemia (B).DUE TO DECREASE 1. Alcohol ingestion. 2. Thiazide diuretics. 3. Lactic acidosis. 4. Aspirin ingestion (lef 5. Diabetic ketoacidos 6. Renal failure due to DECREASED:- (A).DUE TO DIETARY D 1. Dietary deficiency of 2. Fanconi syndrome 3. Multiple sclerosis.	high levels of Uric Acid in the product of purine metabolisi crobial degradation. DPRODUCTION:- gout. Irines (organ meats,legumes, of malignancies especially le & myeloid metaplasia. etc. DEXCREATION (BY KIDNEYS) ess than 2 grams per day ). sis or starvation. any cause etc. EFICIENCY of Zinc, Iron and molybdenum & Wilsons disease.	m . Uric acid is excreted to ,anchovies, etc). eukemais & lymphomas.	a large degree by the	bund a joint. It kidneys and to a smaller degree in the		
INTERPRETATION:- 1.GOUT occurs when 2.Uric Acid is the end intestinal tract by mi- INCREASED:- (A).DUE TO INCREASE 1.Idiopathic primary 9 2.Excessive dietary pu 3.Cytolytic treatment 4.Polycythemai vera 5.Psoriasis. 6.Sickle cell anaemia (B).DUE TO DECREASE 1.Alcohol ingestion. 2.Thiazide diuretics. 3.Lactic acidosis. 4.Aspirin ingestion (lef 5.Diabetic ketoacidos 6.Renal failure due to DECREASED:- (A).DUE TO DIETARY D 1.Dietary deficiency 0 2.Fanconi syndrome 3.Multiple sclerosis. 4.Syndrome of inappr (B).DUE TO INCREASEI	high levels of Uric Acid in the product of purine metabolisi crobial degradation. DPRODUCTION:- gout. urines (organ meats,legumes, of malignancies especially le & myeloid metaplasia. etc. DEXCREATION (BY KIDNEYS) ess than 2 grams per day ). sis or starvation. any cause etc. EFICIENCY of Zinc, Iron and molybdenum & Wilsons disease. opriate antidiuretic hormone DEXCREATION	m . Uric acid is excreted to ,anchovies, etc). eukemais & lymphomas. n. e (SIADH) secretion & low p	a large degree by the	bund a joint. I kidneys and to a smaller degree in the ds and ACTH, anti-coagulants and estrogens e		



TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT

**DR.VINAY CHOPRA** CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS , MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST

**NOT VALID FOR MEDICO LEGAL PURPOSE** 

440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. **REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)** 





# PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana) A PIONEER DIAGNOSTIC CENTRE

🔽 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. NITIKA			
AGE/ GENDER	: 23 YRS/FEMALE	PATIENT	ID	: 1568540
<b>COLLECTED BY</b>	:	REG. NO.	/LAB NO.	: 122408020013
<b>REFERRED BY</b>	:	REGISTR	ATION DATE	: 02/Aug/2024 02:27 PM
BARCODE NO.	: 12503954	COLLECT	ION DATE	: 02/Aug/2024 02:38PM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUT	TE <b>Report</b>	ING DATE	: 02/Aug/2024 03:42PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBAL	A CITY - HARYANA		
			11	Dialogical Deference interval
Test Name		Value	Unit	Biological Reference interval
Test Name		ENDOCRINOLO		
Test Name	THYR		ΟGY	
TRIIODOTHYRONINE	(T3): SERUM	ENDOCRINOLO COID FUNCTION TE 1.321	ΟGY	0.35 - 1.93
TRIIODOTHYRONINE by CMIA (CHEMILUMIN THYROXINE (T4): SEF	(T3): SERUM ESCENT MICROPARTICLE IMMUNOASSAY)	ENDOCRINOLO COID FUNCTION TE 1.321 7.71	OGY ST: TOTAL	
TRIIODOTHYRONINE by CMIA (CHEMILUMIN THYROXINE (T4): SEF by CMIA (CHEMILUMIN THYROID STIMULATI	(T3): SERUM ESCENT MICROPARTICLE IMMUNOASSAY) RUM ESCENT MICROPARTICLE IMMUNOASSAY) NG HORMONE (TSH): SERUM ESCENT MICROPARTICLE IMMUNOASSAY)	<b>ENDOCRINOLO</b> <b>COID FUNCTION TE</b> 1.321 7.71 1.554	DGY ST: TOTAL ng/mL	0.35 - 1.93

TSH levels are subject to circadian variation, reaching peak levels between 2-4 a.m and at a minimum between 6-10 pm. The variation is of the order of 50%. Hence time of the day has influence on the measured serum TSH concentrations. TSH stimulates the production and secretion of the metabolically active hormones, thyroxine (T4) and trilodothyronine (T3). Failure at any level of regulation of the hypothalamic-pituitary-thyroid axis will result in either underproduction (hypothyroidism) or overproduction(hyperthyroidism) of T4 and/or T3.

CLINICAL CONDITION	T3	T4	TSH
Primary Hypothyroidism:	Reduced	Reduced	Increased (Significantly)
Subclinical Hypothyroidism:	Normal or Low Normal	Normal or Low Normal	High
Primary Hyperthyroidism:	Increased	Increased	Reduced (at times undetectable)
Subclinical Hyperthyroidism:	Normal or High Normal	Normal or High Normal	Reduced

#### LIMITATIONS:-

1. T3 and T4 circulates in reversibly bound form with Thyroid binding globulins (TBG), and to a lesser extent albumin and Thyroid binding Pre Albumin so conditions in which TBG and protein levels alter such as pregnancy, excess estrogens, androgens, anabolic steroids and glucocorticoids may falsely affect the T3 and T4 levels and may cause false thyroid values for thyroid function tests.

2. Normal levels of T4 can also be seen in Hyperthyroid patients with :T3 Thyrotoxicosis, Decreased binding capacity due to hypoproteinemia or ingestion of certain drugs (eg: phenytoin , salicylates).

3. Serum T4 levies in neonates and infants are higher than values in the normal adult , due to the increased concentration of TBG in neonate serum.

4. TSH may be normal in central hypothyroidism, recent rapid correction of hyperthyroidism or hypothroidism, pregnancy, phenytoin therapy.

TRIIODOTHYRONINE (T3)		THYROXINE (T4)		THYROID STIMULATING HORMONE (TSH)	
Age	Refferance Range (ng/mL)	Age	Refferance Range (µg/dL)	Age	Reference Range ( μIU/mL)
0 - 7 Days	0.20 - 2.65	0 - 7 Days	5.90 - 18.58	0 - 7 Days	2.43 - 24.3
7 Days - 3 Months	0.36 - 2.59	7 Days - 3 Months	6.39 - 17.66	7 Days - 3 Months	0.58 - 11.00
3 - 6 Months	0.51 - 2.52	3 - 6 Months	6.75 - 17.04	3 Days – 6 Months	0.70 - 8.40





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)

440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. **REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)** 





### PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana) A PIONEER DIAGNOSTIC CENTRE

🖸 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. NITIKA		
AGE/ GENDER	: 23 YRS/FEMALE	PATIENT ID	: 1568540
COLLECTED BY	:	<b>REG. NO./LAB NO.</b>	: 122408020013
<b>REFERRED BY</b>	:	<b>REGISTRATION DATE</b>	: 02/Aug/2024 02:27 PM
BARCODE NO.	: 12503954	<b>COLLECTION DATE</b>	: 02/Aug/2024 02:38PM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	<b>REPORTING DATE</b>	: 02/Aug/2024 03:42PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - H	ARYANA	

Test Name	Name		Value		Unit		Biological Reference interva	
6 - 12 Months	0.74 - 2.40	6 - 12 Months	7.10 - 16.16	6 – 12 Months	0.70 - 7.00			
1 - 10 Years	0.92 - 2.28	1 - 10 Years	6.00 - 13.80	1 – 10 Years	0.60 - 5.50			
11- 19 Years	0.35 - 1.93	11 - 19 Years	4.87- 13.20	11 – 19 Years	0.50 - 5.50			
> 20 years (Adults)	0.35 - 1.93	> 20 Years (Adults)	4.87 - 12.60	> 20 Years (Adults)	0.35-5.50			
	RECO	MMENDATIONS OF TSH L	EVELS DURING PRE	GNANCY ( µIU/mL)				
1st Trimester			0.10 - 2.50					
	2nd Trimester		0.20 - 3.00					
	3rd Trimester			0.30 - 4.10				

### INCREASED TSH LEVELS:

1.Primary or untreated hypothyroidism may vary from 3 times to more than 100 times normal depending upon degree of hypofunction.

2.Hypothyroid patients receiving insufficient thyroid replacement therapy.

3.Hashimotos thyroiditis

4.DRUGS: Amphetamines, idonie containing agents & dopamine antagonist.

5.Neonatal period, increase in 1st 2-3 days of life due to post-natal surge

DECREASED TSH LEVELS:

1.Toxic multi-nodular goitre & Thyroiditis.

2. Over replacement of thyroid harmone in treatment of hypothyroidism.

3. Autonomously functioning Thyroid adenoma

4. Secondary pituatary or hypothalmic hypothyroidism

5. Acute psychiatric illness

6.Severe dehydration.

7.DRUGS: Glucocorticoids, Dopamine, Levodopa, T4 replacement therapy, Anti-thyroid drugs for thyrotoxicosis.

8. Pregnancy: 1st and 2nd Trimester





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)

440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600, REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)





## PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

🕻 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. NITIKA			
AGE/ GENDER	: 23 YRS/FEMALE	PATI	ENT ID	: 1568540
<b>COLLECTED BY</b>	:	REG.	NO./LAB NO.	: 122408020013
<b>REFERRED BY</b>	:	REGI	STRATION DATE	: 02/Aug/2024 02:27 PM
BARCODE NO.	: 12503954	COLI	ECTION DATE	: 02/Aug/2024 02:38PM
CLIENT CODE.	: P.K.R JAIN HEALTHCAR	E INSTITUTE <b>REP</b> (	DRTING DATE	: 02/Aug/2024 03:13PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROA	AD, AMBALA CITY - HARYAN	A	
Test Name		Value	Unit	Biological Reference interval
		IMMUNOPATHOLO	GY/SEROLOGY	
		WIDAL SLIDE AGGLU	TINATION TEST	
SALMONELLA TYPHI	0	1 : 80	TITRE	1 : 80
by SLIDE AGGLUTINA		1.40	TITOF	1.1/0
SALMONELLA TYPHI H 1 : 40 by SLIDE AGGLUTINATION		1:40	TITRE	1 : 160
SALMONELLA PARATYPHI AH		1:20	TITRE	1 : 160
by SLIDE AGGLUTINATION				
SALMONELLA PARA	ГҮРНІ ВН	1:20	TITRE	1:160

### by SLIDE AGGLUTINATION

### **INTERPRETATION:**

1. Titres of 1:80 or more for "O" agglutinin is considered significant.

2. Titres of 1:160 or more for "H" agglutinin is considered significant.

### LIMITATIONS:

1.Agglutinins usually appear by 5th to 6th day of illness of enteric fever, hence a negative result in early stage is inconclusive. The titre then rises till 3rd or 4th week, after which it declines gradually.

2.Lower titres may be found in normal individuals.

3.A single positive result has less significance than the rising agglutination titre, since demonstration of rising titre four or more in 1st and 3rd week is considered as a definite evidence of infection.

4.A simultaneous rise in H agglutinins is suggestive of paratyphoid infection.

### NOTE:

1. Individuals with prior infection or immunization with TAB vaccine may develop an ANAMNESTIC RESPONSE (False-Positive) during an unrelated fever *i.e* High titres of antibodies to various antigens. This may be differentiated by repitition of the test after a week.

2. The anamnestic response shows only a transient rise, while in enteric fever rise is sustained.

3.H agglutinins tend to persist for many months after vaccination but O agglutinins tend to disappear sooner i.e within 6 months. Therefore rise in Oagglutinins indicate recent infection.

\*\*\* End Of Report \*\*\*





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)

440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)

