



# P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

**A PIONEER DIAGNOSTIC CENTRE**

☎ 0171-2532620, 8222896961

✉ pkrjainhealthcare@gmail.com

NAME	: Mrs. SHEENA BAJAJ	PATIENT ID	: 1574335
AGE/ GENDER	: 32 YRS/FEMALE	REG. NO./LAB NO.	: 122408080004
COLLECTED BY	:	REGISTRATION DATE	: 08/Aug/2024 09:22 AM
REFERRED BY	:	COLLECTION DATE	: 08/Aug/2024 09:49AM
BARCODE NO.	: 12504044	REPORTING DATE	: 08/Aug/2024 10:39AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE		
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

## IMMUNOPATHOLOGY/SEROLOGY HEPATITIS C VIRUS (HCV) ANTIBODIES SCREENING

HEPATITIS C ANTIBODY (HCV) TOTAL  
RESULT

NON - REACTIVE

by IMMUNOCHROMATOGRAPHY

### INTERPRETATION:

1.Anti HCV total antibody assay identifies presence IgG antibodies in the serum . It is a useful screening test with a specificity of nearly 99%.  
2.It becomes positive approximately 24 weeks after exposure. The test can not isolate an active ongoing HCV infection from an old infection that has been cleared. All positive results must be confirmed for active disease by an HCV PCR test .

### FALSE NEGATIVE RESULTS SEEN IN:

- 1.Window period
- 2.Immunocompromised states.



DR.VINAY CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS , MD (PATHOLOGY)





# P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

**A PIONEER DIAGNOSTIC CENTRE**

☎ 0171-2532620, 8222896961 ✉ pkrjainhealthcare@gmail.com

<b>NAME</b>	: Mrs. SHEENA BAJAJ	<b>PATIENT ID</b>	: 1574335
<b>AGE/ GENDER</b>	: 32 YRS/FEMALE	<b>REG. NO./LAB NO.</b>	: 122408080004
<b>COLLECTED BY</b>	:	<b>REGISTRATION DATE</b>	: 08/Aug/2024 09:22 AM
<b>REFERRED BY</b>	:	<b>COLLECTION DATE</b>	: 08/Aug/2024 09:49AM
<b>BARCODE NO.</b>	: 12504044	<b>REPORTING DATE</b>	: 08/Aug/2024 10:39AM
<b>CLIENT CODE.</b>	: P.K.R JAIN HEALTHCARE INSTITUTE		
<b>CLIENT ADDRESS</b>	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

## ANTI HUMAN IMMUNODEFICIENCY VIRUS (HIV) ANTIBODIES HIV (1 & 2) SCREENING

HIV 1/2 AND P24 ANTIGEN RESULT NON - REACTIVE  
by IMMUNOCHROMATOGRAPHY

### INTERPRETATION:-

- 1.AIDS is caused by at least 2 known types of HIV viruses, HIV-1 and HIV HIV-2.
- 2.This NACO approved immuno-chromatographic solid phase ELISA assay detects antibodies against both HIV-1 and HIV-2 viruses.
- 3.The test is used for routine serologic screening of patients at risk for HIV-1 or HIV-2 infection.
- 4.All screening ELISA assays for HIV antibody detection have high sensitivity but have low specificity.
- 5.At this laboratory, all positive samples are cross checked for positivity with two alternate assays prior to reporting.

### NOTE:-


- 1.Confirmatory testing by Western blot is recommended for patients who are reactive for HIV by this assay.
- 2.Antibodies against HIV-1 and HIV-2 are usually not detectable until 6 to 12 weeks following exposure (window period) and are almost always detectable by 12 months.
- 3.The test is not recommended for children born to HIV infected mothers till the child turns two years old (as HIV antibodies may be transmitted passively to the child trans-placentally).

### FALSE NEGATIVE RESULT SEEN IN:

- 1.Window period
- 2.Severe immuno-suppression including advanced AIDS.



  
DR.VINAY CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

  
DR.YUGAM CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY)





# P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

**A PIONEER DIAGNOSTIC CENTRE**

☎ 0171-2532620, 8222896961 ✉ [pkrajainhealthcare@gmail.com](mailto:pkrajainhealthcare@gmail.com)

TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT.

<b>NAME</b>	: Mrs. SHEENA BAJAJ	<b>PATIENT ID</b>	: 1574335
<b>AGE/ GENDER</b>	: 32 YRS/FEMALE	<b>REG. NO./LAB NO.</b>	: 122408080004
<b>COLLECTED BY</b>	:	<b>REGISTRATION DATE</b>	: 08/Aug/2024 09:22 AM
<b>REFERRED BY</b>	:	<b>COLLECTION DATE</b>	: 08/Aug/2024 09:49AM
<b>BARCODE NO.</b>	: 12504044	<b>REPORTING DATE</b>	: 08/Aug/2024 10:39AM
<b>CLIENT CODE.</b>	: P.K.R JAIN HEALTHCARE INSTITUTE		
<b>CLIENT ADDRESS</b>	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

## HEPATITIS B SURFACE ANTIGEN (HBsAg) SCREENING

HEPATITIS B SURFACE ANTIGEN (HBsAg) NON - REACTIVE  
RESULT

by IMMUNOCHROMATOGRAPHY

### INTERPRETATION:-

1.HBsAG is the first serological marker of HBV infection to appear in the blood (approximately 30-60 days after infection and prior to the onset of clinical disease). It is also the last viral protein to disappear from blood and usually disappears by three months after infection in self limiting acute Hepatitis B viral infection.

2.Persistence of HBsAg in blood for more than six months implies chronic infection. It is the most common marker used for diagnosis of an acute Hepatitis B infection but has very limited role in assessing patients suffering from chronic hepatitis.

### **FALSE NEGATIVE RESULT SEEN IN:**

- 1.Window period.
- 2.Infection with HBsAg mutant strains
- 3.Hepatitis B Surface antigen (HBsAg) is the earliest indicator of HBV infection. Usually it appears in 27 - 41 days (as early as 14 days).
- 4.Appears 7 - 26 days before biochemical abnormalities. Peaks as ALT rises. Persists during the acute illness. Usually disappears 12- 20 weeks after the onset of symptoms / laboratory abnormalities in 90% of cases.
- 5.Is the most reliable serologic marker of HBV infection. Persistence > 6 months defines carrier state. May also be found in chronic infection.Hepatitis B vaccination does not cause a positive HBsAg. Titers are not of clinical value.

### **NOTE:-**

- 1.All reactive HBsAG Should be reconfirmed with neutralization test(HBsAg confirmatory test).
- 2.Anti - HAV IgM appears at the same time as symptoms in > 99% of cases, peaks within the first month, becomes nondetectable in 12 months (usually 6 months). Presence confirms diagnosis of recent acute infection.

\*\*\* End Of Report \*\*\*



DR.VINAY CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY)

