



# P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

**A PIONEER DIAGNOSTIC CENTRE**

☎ 0171-2532620, 8222896961 ✉ [pkrajainhealthcare@gmail.com](mailto:pkrajainhealthcare@gmail.com)

|                |  |                   |                        |
|----------------|--|-------------------|------------------------|
| NAME           | : Master. AARAV KHANNA                         | PATIENT ID        | : 1574906              |
| AGE/ GENDER    | : 11 YRS/MALE                                  | REG. NO./LAB NO.  | : 122408080020         |
| COLLECTED BY   | :  | REGISTRATION DATE | : 08/Aug/2024 03:24 PM |
| REFERRED BY    | :  | COLLECTION DATE   | : 08/Aug/2024 09:06PM  |
| BARCODE NO.    | : 12504060                                     | REPORTING DATE    | : 08/Aug/2024 05:01PM  |
| CLIENT CODE.   | : P.K.R JAIN HEALTHCARE INSTITUTE              |                   |                        |
| CLIENT ADDRESS | : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA |                   |                        |

| Test Name | Value | Unit | Biological Reference interval |
|-----------|-------|------|-------------------------------|
|-----------|-------|------|-------------------------------|

## CLINICAL CHEMISTRY/BIOCHEMISTRY

### CALCIUM

|                                    |       |       |              |
|------------------------------------|-------|-------|--------------|
| CALCIUM: SERUM                     | 10.43 | mg/dL | 8.50 - 10.60 |
| by ARSENAZO III, SPECTROPHOTOMETRY |       |       |              |

#### INTERPRETATION:-

1. Serum calcium (total) estimation is used for the diagnosis and monitoring of a wide range of disorders including diseases of bone, kidney, parathyroid gland, or gastrointestinal tract.
2. Calcium levels may also reflect abnormal vitamin D or protein levels.
3. The calcium content of an adult is somewhat over 1 kg (about 2% of the body weight). Of this, 99% is present as calcium hydroxyapatite in bones and <1% is present in the extra-osseous intracellular space or extracellular space (ECS).
4. In serum, calcium is bound to a considerable extent to proteins (approximately 40%), 10% is in the form of inorganic complexes, and 50% is present as free or ionized calcium.

**NOTE:-** Calcium ions affect the contractility of the heart and the skeletal musculature, and are essential for the function of the nervous system. In addition, calcium ions play an important role in blood clotting and bone mineralization.

#### HYPOCALCEMIA (LOW CALCIUM LEVELS) CAUSES :-

1. Due to the absence or impaired function of the parathyroid glands or impaired vitamin-D synthesis.
2. Chronic renal failure is also frequently associated with hypocalcemia due to decreased vitamin-D synthesis as well as hyperphosphatemia and skeletal resistance to the action of parathyroid hormone (PTH).
3. **NOTE:-** A characteristic symptom of hypocalcemia is latent or manifest tetany and osteomalacia.

#### HYPERCALCEMIA (INCREASE CALCIUM LEVELS) CAUSES:-

1. Increased mobilization of calcium from the skeletal system or increased intestinal absorption.
  2. Primary hyperparathyroidism (pHPT)
  3. Bone metastasis of carcinoma of the breast, prostate, thyroid gland, or lung.
- NOTE:-** Severe hypercalcemia may result in cardiac arrhythmia.



  
DR. VINAY CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

  
DR. YUGAM CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY)





# P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

**A PIONEER DIAGNOSTIC CENTRE**

☎ 0171-2532620, 8222896961 ✉ pkrajainhealthcare@gmail.com

**NAME** : Master. AARAV KHANNA  
**AGE/ GENDER** : 11 YRS/MALE **PATIENT ID** : 1574906  
**COLLECTED BY** : **REG. NO./LAB NO.** : 122408080020  
**REFERRED BY** : **REGISTRATION DATE** : 08/Aug/2024 03:24 PM  
**BARCODE NO.** : 12504060 **COLLECTION DATE** : 08/Aug/2024 09:06PM  
**CLIENT CODE.** : P.K.R JAIN HEALTHCARE INSTITUTE **REPORTING DATE** : 08/Aug/2024 05:01PM  
**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

| Test Name | Value | Unit | Biological Reference interval |
|-----------|-------|------|-------------------------------|
|-----------|-------|------|-------------------------------|

## VITAMINS

### VITAMIN D/25 HYDROXY VITAMIN D3

**VITAMIN D (25-HYDROXY VITAMIN D3): SERUM**  
by CLIA (CHEMILUMINESCENCE IMMUNOASSAY)

9.527<sup>L</sup>

ng/mL

**DEFICIENCY:** < 20.0  
**INSUFFICIENCY:** 20.0 - 30.0  
**SUFFICIENCY:** 30.0 - 100.0  
**TOXICITY:** > 100.0

#### INTERPRETATION:

|                  |          |       |
|------------------|----------|-------|
| DEFICIENT:       | < 20     | ng/mL |
| INSUFFICIENT:    | 21 - 29  | ng/mL |
| PREFERRED RANGE: | 30 - 100 | ng/mL |
| INTOXICATION:    | > 100    | ng/mL |

- Vitamin D compounds are derived from dietary ergocalciferol (from plants, Vitamin D2), or cholecalciferol (from animals, Vitamin D3), or by conversion of 7- dihydrocholecalciferol to Vitamin D3 in the skin upon Ultraviolet exposure.
- 25-OH--Vitamin D represents the main body reservoir and transport form of Vitamin D and transport form of Vitamin D, being stored in adipose tissue and tightly bound by a transport protein while in circulation.
- Vitamin D plays a primary role in the maintenance of calcium homeostasis. It promotes calcium absorption, renal calcium absorption and phosphate reabsorption, skeletal calcium deposition, calcium mobilization, mainly regulated by parathyroid hormone (PTH).
- Severe deficiency may lead to failure to mineralize newly formed osteoid in bone, resulting in rickets in children and osteomalacia in adults.

#### DECREASED:

- Lack of sunshine exposure.
- Inadequate intake, malabsorption (celiac disease)
- Depressed Hepatic Vitamin D 25- hydroxylase activity
- Secondary to advanced Liver disease
- Osteoporosis and Secondary Hyperparathyroidism (Mild to Moderate deficiency)
- Enzyme Inducing drugs: anti-epileptic drugs like phenytoin, phenobarbital and carbamazepine, that increases Vitamin D metabolism.

#### INCREASED:

- Hypervitaminosis D is Rare, and is seen only after prolonged exposure to extremely high doses of Vitamin D. When it occurs, it can result in severe hypercalcemia and hyperphosphatemia.

**CAUTION:** Replacement therapy in deficient individuals must be monitored by periodic assessment of Vitamin D levels in order to prevent hypervitaminosis D

**NOTE:-** Dark coloured individuals as compare to whites, is at higher risk of developing Vitamin D deficiency due to excess of melanin pigment which interfere with Vitamin D absorption.



DR.VINAY CHOPRA

CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA

CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY)





# P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

**A PIONEER DIAGNOSTIC CENTRE**

☎ 0171-2532620, 8222896961

✉ pkrjainhealthcare@gmail.com

**NAME** : Master. AARAV KHANNA  
**AGE/ GENDER** : 11 YRS/MALE  
**COLLECTED BY** :  
**REFERRED BY** :  
**BARCODE NO.** : 12504060  
**CLIENT CODE.** : P.K.R JAIN HEALTHCARE INSTITUTE  
**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**PATIENT ID** : 1574906  
**REG. NO./LAB NO.** : 122408080020  
**REGISTRATION DATE** : 08/Aug/2024 03:24 PM  
**COLLECTION DATE** : 08/Aug/2024 09:06PM  
**REPORTING DATE** : 08/Aug/2024 10:44PM

| Test Name | Value | Unit | Biological Reference interval |
|-----------|-------|------|-------------------------------|
|-----------|-------|------|-------------------------------|

## VITAMIN B12/COBALAMIN

VITAMIN B12/COBALAMIN: SERUM 242.7 pg/mL 200.0 - 1100.0

by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

### INTERPRETATION:-


| INCREASED VITAMIN B12         | DECREASED VITAMIN B12                         |
|-------------------------------|---|
| 1.Ingestion of Vitamin C      | 1.Pregnancy                                   |
| 2.Ingestion of Estrogen       | 2.DRUGS:Aspirin, Anti-convulsants, Colchicine |
| 3.Ingestion of Vitamin A      | 3.Ethanol lgestion                            |
| 4.Hepatocellular injury       | 4. Contraceptive Harmones                     |
| 5.Myeloproliferative disorder | 5.Haemodialysis                               |
| 6.Uremia                      | 6. Multiple Myeloma                           |

1.Vitamin B12 (cobalamin) is necessary for hematopoiesis and normal neuronal function.  
2.In humans, it is obtained only from animal proteins and requires intrinsic factor (IF) for absorption.  
3.The body uses its vitamin B12 stores very economically, reabsorbing vitamin B12 from the ileum and returning it to the liver; very little is excreted.  
4.Vitamin B12 deficiency may be due to lack of IF secretion by gastric mucosa (eg, gastrectomy, gastric atrophy) or intestinal malabsorption (eg, ileal resection, small intestinal diseases).  
5.Vitamin B12 deficiency frequently causes macrocytic anemia, glossitis, peripheral neuropathy, weakness, hyperreflexia, ataxia, loss of proprioception, poor coordination, and affective behavioral changes. These manifestations may occur in any combination; many patients have the neurologic defects without macrocytic anemia.  
6.Serum methylmalonic acid and homocysteine levels are also elevated in vitamin B12 deficiency states.  
7.Follow-up testing for antibodies to intrinsic factor (IF) is recommended to identify this potential cause of vitamin B12 malabsorption.  
**NOTE:**A normal serum concentration of vitamin B12 does not rule out tissue deficiency of vitamin B12. The most sensitive test for vitamin B12 deficiency at the cellular level is the assay for MMA. If clinical symptoms suggest deficiency, measurement of MMA and homocysteine should be considered, even if serum vitamin B12 concentrations are normal.

\*\*\* End Of Report \*\*\*



  
DR.VINAY CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

  
DR.YUGAM CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY)

