



P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

☎ 0171-2532620, 8222896961 ✉ pkrjainhealthcare@gmail.com

TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT.

NAME	: Mr. SACHIN	PATIENT ID	: 1583171
AGE/ GENDER	: 43 YRS/MALE	REG. NO./LAB NO.	: 122408170017
COLLECTED BY	:	REGISTRATION DATE	: 17/Aug/2024 12:08 PM
REFERRED BY	:	COLLECTION DATE	: 17/Aug/2024 12:13PM
BARCODE NO.	: 12504198	REPORTING DATE	: 17/Aug/2024 01:51PM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE		
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

CLINICAL CHEMISTRY/BIOCHEMISTRY

CALCIUM

CALCIUM: SERUM	9.65	mg/dL	8.50 - 10.60
by ARSENAZO III, SPECTROPHOTOMETRY			

INTERPRETATION:-

1. Serum calcium (total) estimation is used for the diagnosis and monitoring of a wide range of disorders including diseases of bone, kidney, parathyroid gland, or gastrointestinal tract.
2. Calcium levels may also reflect abnormal vitamin D or protein levels.
3. The calcium content of an adult is somewhat over 1 kg (about 2% of the body weight). Of this, 99% is present as calcium hydroxyapatite in bones and <1% is present in the extra-osseous intracellular space or extracellular space (ECS).
4. In serum, calcium is bound to a considerable extent to proteins (approximately 40%), 10% is in the form of inorganic complexes, and 50% is present as free or ionized calcium.

NOTE:- Calcium ions affect the contractility of the heart and the skeletal musculature, and are essential for the function of the nervous system. In addition, calcium ions play an important role in blood clotting and bone mineralization.


HYPOCALCEMIA (LOW CALCIUM LEVELS) CAUSES :-

1. Due to the absence or impaired function of the parathyroid glands or impaired vitamin-D synthesis.
2. Chronic renal failure is also frequently associated with hypocalcemia due to decreased vitamin-D synthesis as well as hyperphosphatemia and skeletal resistance to the action of parathyroid hormone (PTH).
3. **NOTE:-** A characteristic symptom of hypocalcemia is latent or manifest tetany and osteomalacia.

HYPERCALCEMIA (INCREASE CALCIUM LEVELS) CAUSES:-

1. Increased mobilization of calcium from the skeletal system or increased intestinal absorption.
 2. Primary hyperparathyroidism (pHPT)
 3. Bone metastasis of carcinoma of the breast, prostate, thyroid gland, or lung.
- NOTE:-** Severe hypercalcemia may result in cardiac arrhythmia.




DR. VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)


DR. YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY)





P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

☎ 0171-2532620, 8222896961 ✉ pkrjainhealthcare@gmail.com

NAME : Mr. SACHIN
AGE/ GENDER : 43 YRS/MALE **PATIENT ID** : 1583171
COLLECTED BY : **REG. NO./LAB NO.** : 122408170017
REFERRED BY : **REGISTRATION DATE** : 17/Aug/2024 12:08 PM
BARCODE NO. : 12504198 **COLLECTION DATE** : 17/Aug/2024 12:13PM
CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE **REPORTING DATE** : 17/Aug/2024 07:43PM
CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

ENDOCRINOLOGY

INTACT PARATHYROID HORMONE (PTH)

INTACT PARATHROID HORMONE (PTH): SERUM by CLIA (CHEMILUMINESCENCE IMMUNOASSAY)	140.3 ^H	pg/mL	9.5 - 75.0
---	--------------------	-------	------------

Intrepretation:-

Parathyroid hormone (PTH) is produced and secreted by the parathyroid glands, which are located along the posterior aspect of the thyroid gland. The serum calcium level regulates PTH secretion via negative feedback through the parathyroid calcium sensing receptor (CASR). Decreased calcium levels stimulate PTH release. Secreted PTH interacts with its specific type II G-protein receptor, causing rapid increases in renal tubular reabsorption of calcium and decreased phosphorus reabsorption. It also participates in long-term calciostatic functions by enhancing mobilization of calcium from bone and increasing renal synthesis of 1,25-dihydroxy vitamin D, which, in turn, increases intestinal calcium absorption.

The assay is useful for:

- Differential diagnosis of hypercalcemia
- Diagnosis of primary, secondary, and tertiary hyperparathyroidism
- Diagnosis of hypoparathyroidism
- Monitoring end-stage renal failure patients for possible renal osteodystrophy

Interpretation of results:

- An (appropriately) low PTH level and high phosphorus level in a hypercalcemic patient suggests that the hypercalcemia is not caused by PTH or PTH-like substances.
- An (appropriately) low PTH level with a low phosphorus level in a hypercalcemic patient suggests the diagnosis of paraneoplastic hypercalcemia.
- A low or normal PTH in a patient with hypocalcemia suggests hypoparathyroidism.


Low serum calcium and high PTH levels in a patient with normal renal function suggest resistance to PTH action (pseudohypoparathyroidism type 1a, 1b, 1c, or 2) or, very rarely, bio-ineffective PTH.

Elevated PTH value with a normal serum calcium in many cases in India is due to secondary hyperparathyroidism, primary cause being Vitamin D deficiency.

*** End Of Report ***




DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)


DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY)

