CLIENT CODE.



PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

: 30/Aug/2024 01:02PM

NAME : Mr. SUNIL KUMAR

AGE/ GENDER : 59 YRS/MALE **PATIENT ID** : 1595986

COLLECTED BY : 122408300007 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 30/Aug/2024 09:45 AM BARCODE NO. : 12504379 **COLLECTION DATE** : 30/Aug/2024 10:01AM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

Test Name Value Unit **Biological Reference interval**

CLINICAL CHEMISTRY/BIOCHEMISTRY

REPORTING DATE

UREA

38.06 10.00 - 50.00 **UREA: SERUM** mg/dL

by UREASE - GLUTAMATE DEHYDROGENASE (GLDH)



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)





PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

REPORTING DATE

: 30/Aug/2024 01:02PM

NAME : Mr. SUNIL KUMAR

AGE/ GENDER : 59 YRS/MALE **PATIENT ID** : 1595986

COLLECTED BY : 122408300007 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 30/Aug/2024 09:45 AM BARCODE NO. : 12504379 **COLLECTION DATE** : 30/Aug/2024 10:01AM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

Test Name Value Unit **Biological Reference interval**

URIC ACID

URIC ACID: SERUM 5.2 mg/dL 3.60 - 7.70

by URICASE - OXIDASE PEROXIDASE

INTERPRETATION:-

CLIENT CODE.

1.GOUT occurs when high levels of Uric Acid in the blood cause crystals to form & accumulate around a joint.

2.Uric Acid is the end product of purine metabolism. Uric acid is excreted to a large degree by the kidneys and to a smaller degree in the intestinal tract by microbial degradation.

INCREASED:-

(A).DUE TO INCREASED PRODUCTION:-

1. Idiopathic primary gout.

2. Excessive dietary purines (organ meats, legumes, anchovies, etc).

3. Cytolytic treatment of malignancies especially leukemais & lymphomas.

4. Polycythemai vera & myeloid metaplasia.

5.Psoriasis.

6. Sickle cell anaemia etc.

(B).DUE TO DECREASED EXCREATION (BY KIDNEYS)

1. Alcohol ingestion.

2. Thiazide diuretics

3.Lactic acidosis.

- 4. Aspirin ingestion (less than 2 grams per day).
- 5. Diabetic ketoacidosis or starvation.
- 6.Renal failure due to any cause etc.

DECREASED:-

(A).DUE TO DIETARY DEFICIENCY

- 1. Dietary deficiency of Zinc, Iron and molybdenum.
- 2. Fanconi syndrome & Wilsons disease.
- 3. Multiple sclerosis.
- 4. Syndrome of inappropriate antidiuretic hormone (SIADH) secretion & low purine diet etc.

(B).DUE TO INCREASED EXCREATION

1.Drugs:-Probenecid, sulphinpyrazone, aspirin doses (more than 4 grams per day), corticosterroids and ACTH, anti-coagulants and estrogens etc.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)





PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

NAME : Mr. SUNIL KUMAR

AGE/ GENDER : 59 YRS/MALE **PATIENT ID** : 1595986

COLLECTED BY : 122408300007 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 30/Aug/2024 09:45 AM BARCODE NO. : 12504379 **COLLECTION DATE** : 30/Aug/2024 10:01AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 30/Aug/2024 04:11PM

3.91

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name Value Unit **Biological Reference interval**

TUMOUR MARKER

PROSTATE SPECIFIC ANTIGEN (PSA) - TOTAL

PROSTATE SPECIFIC ANTIGEN (PSA) - TOTAL:

ng/mL

0.0 - 4.0

by CLIA (CHEMILUMINESCENCE IMMUNOASSAY)

INTERPRETATION:-

Expected Values for the PSA	
Smokers	< 4 ng/ml
Non-smokers	< 4 ng/ml

- 1. Prostate-specific antigen (PSA) is a glycoprotein that is produced by the prostate gland, the lining of the urethra, and the bulbourethral gland.
- 2. Normally, very little PSA is secreted in the blood.

INCREASED:-

- 1.Increased in glandular size and tissue damage caused by benign prostatic hypertrophy.
- 2. Prostatitis.
- 3. Prostate cancer may increase circulating PSA levels.
- 4.In patients with previously diagnosed prostate cance, PSA testing is advocated as an early indicator of tumor recurrence and as an indicator of response to therapy.

The test is also useful for initial screening for prostate cancer:-

- 1.Total PSA levels < 2 ng/ml almost rule out the possibility of prostatic malignancy.
- 2.Total PSA levels between 2 and 10 ng/ml lie in the grey zone. Such values may be obtained in prostatitis, benign hyperplasia and malignancy. Further testing including a free PSA/PSA ratio and prostate biopsy is recommended for these patients for confirmation of the diagnosis. 3. Total PSA values >10 ng/ml are highly suspicious for prostate cancer but further testing, such as prostate biopsy, is needed to diagnose the exact pathology.

*** End Of Report ***



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)