



# P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

**A PIONEER DIAGNOSTIC CENTRE**

☎ 0171-2532620, 8222896961

✉ pkrjainhealthcare@gmail.com

**NAME** : Mrs. HEENA DEVI  
**AGE/ GENDER** : 30 YRS/FEMALE  
**COLLECTED BY** :  
**REFERRED BY** :  
**BARCODE NO.** : 12504522  
**CLIENT CODE.** : P.K.R JAIN HEALTHCARE INSTITUTE  
**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**PATIENT ID** : 1603799  
**REG. NO./LAB NO.** : 122409060005  
**REGISTRATION DATE** : 06/Sep/2024 09:15 AM  
**COLLECTION DATE** : 06/Sep/2024 09:31AM  
**REPORTING DATE** : 07/Sep/2024 05:53AM

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

## HAEMATOLOGY

### HAEMOGLOBIN - HIGH PERFORMANCE LIQUID CHROMATOGRAPHY (HB-HPLC)

#### HAEMOGLOBIN VARIANTS

HAEMOGLOBIN A0 (ADULT) by HPLC (HIGH PERFORMANCE LIQUID CHROMATOGRAPHY)	85.1	%	83.00 - 90.00
HAEMOGLOBIN F (FOETAL) by HPLC (HIGH PERFORMANCE LIQUID CHROMATOGRAPHY)	<0.8	%	0.00 - 2.0
HAEMOGLOBIN A2 by HPLC (HIGH PERFORMANCE LIQUID CHROMATOGRAPHY)	2.7	%	1.50 - 3.70
PEAK 3 by HPLC (HIGH PERFORMANCE LIQUID CHROMATOGRAPHY)	5.3	%	< 10.0
OTHERS-NON SPECIFIC by HPLC (HIGH PERFORMANCE LIQUID CHROMATOGRAPHY)	ABSENT	%	ABSENT
HAEMOGLOBIN S by HPLC (HIGH PERFORMANCE LIQUID CHROMATOGRAPHY)	NOT DETECTED	%	< 0.02
HAEMOGLOBIN D (PUNJAB) by HPLC (HIGH PERFORMANCE LIQUID CHROMATOGRAPHY)	NOT DETECTED	%	< 0.02
HAEMOGLOBIN E by HPLC (HIGH PERFORMANCE LIQUID CHROMATOGRAPHY)	NOT DETECTED	%	< 0.02
HAEMOGLOBIN C by HPLC (HIGH PERFORMANCE LIQUID CHROMATOGRAPHY)	NOT DETECTED	%	< 0.02
UNKNOWN UNIDENTIFIED VARIANTS by HPLC (HIGH PERFORMANCE LIQUID CHROMATOGRAPHY)	NOT DETECTED	%	< 0.02
GLYCOSYLATED HAEMOGLOBIN (HbA1c): WHOLE BLOOD by HPLC (HIGH PERFORMANCE LIQUID CHROMATOGRAPHY)	5.7	%	4.0 - 6.4

#### RED BLOOD CELLS (RBCS) COUNT AND INDICES

HAEMOGLOBIN (HB) by AUTOMATED HEMATOLOGY ANALYZER	12	gm/dL	12.0 - 16.0
RED BLOOD CELL (RBC) COUNT by AUTOMATED HEMATOLOGY ANALYZER	4.32	Millions/cmm	3.50 - 5.00
PACKED CELL VOLUME (PCV) by AUTOMATED HEMATOLOGY ANALYZER	36.8 <sup>L</sup>	%	37.0 - 50.0
MEAN CORPUSCULAR VOLUME (MCV)	85.1	fL	80.0 - 100.0



DR.VINAY CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY)





# P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

**A PIONEER DIAGNOSTIC CENTRE**

☎ 0171-2532620, 8222896961 ✉ pkrjainhealthcare@gmail.com

**NAME** : Mrs. HEENA DEVI  
**AGE/ GENDER** : 30 YRS/FEMALE  
**COLLECTED BY** :  
**REFERRED BY** :  
**BARCODE NO.** : 12504522  
**CLIENT CODE.** : P.K.R JAIN HEALTHCARE INSTITUTE  
**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**PATIENT ID** : 1603799  
**REG. NO./LAB NO.** : 122409060005  
**REGISTRATION DATE** : 06/Sep/2024 09:15 AM  
**COLLECTION DATE** : 06/Sep/2024 09:31AM  
**REPORTING DATE** : 07/Sep/2024 05:53AM

Test Name	Value	Unit	Biological Reference interval
by AUTOMATED HEMATOLOGY ANALYZER			
MEAN CORPUSCULAR HAEMOGLOBIN (MCH)	27.7	pg	27.0 - 34.0
by AUTOMATED HEMATOLOGY ANALYZER			
MEAN CORPUSCULAR HEMOGLOBIN CONC. (MCHC)	32.6	g/dL	32.0 - 36.0
by AUTOMATED HEMATOLOGY ANALYZER			
RED CELL DISTRIBUTION WIDTH (RDW-CV)	13.2	%	11.00 - 16.00
by AUTOMATED HEMATOLOGY ANALYZER			
RED CELL DISTRIBUTION WIDTH (RDW-SD)	41.7	fL	35.0 - 56.0
by AUTOMATED HEMATOLOGY ANALYZER			
<b>OTHERS</b>			
NAKED EYE SINGLE TUBE RED CELL OSMOTIC FRAGILITY TEST	NEGATIVE (-ve)		NEGATIVE (-ve)
by SINGLE RED CELL OSMOTIC FRAGILITY			
MENTZERS INDEX	19.7	RATIO	BETA THALASSEMIA TRAIT: < 13.0 IRON DEFICIENCY ANEMIA: >13.0
by CALCULATED			
<b>INTERPRETATION</b>	THE ABOVE FINDINGS ARE SUGGESTIVE OF NORMAL HAEMOGLOBIN CHROMATOGRAPHIC PATTERN		

#### INTERPRETATION:

The Thalassemia syndromes, considered the most common genetic disorder worldwide, are a heterogenous group of mandelian disorders, all characterized by a lack of/or decreased synthesis of either the alpha-globin chains (alpha thalassemia) or the beta-globin chains (beta thalassemia) of haemoglobin.

#### **HIGH PERFORMANCE LIQUID CHROMATOGRAPHY (HPLC):**

1.HAEMOGLOBIN VARIANT ANALYSIS, BLOOD- High Performance liquid chromatography (HPLC) is a fast & accurate method for determining the presence and for quatitation of various types of normal haemoglobin and common abnormal hb variants, including but not limited to Hb S, C, E, D and Beta -thalassemia.

2.The diagnosis of these abnormal haemoglobin should be confirmed by DNA analysis.

3.The method use has a limited role in the diagnosis of alpha thalassemia.

4.Slight elevation in haemoglobin A2 may also occur in hyperthyroidism or when there is deficiency of vitamin b12 or folate and this should be istinguished from inherited elevation of HbA2 in Beta- thalassemia trait.

#### **NAKED EYE SINGLE TUBE RED CELL OSMOTIC FRAGILITY TEST (NESTROFT):**

1.It is a screening test to distinguish beta thalassemia trait. Also called as Naked Eye Single Tube Red Cell Osmotic Fragility Test.

2.The test showed a sensitivity of 100%, specificity of 85.47%, a positive predictive value of 66% and a negative predictive value of 100%.

3.A high negative predictive value can reasonably rule out beta thalassemia trait cases. So, it should be adopted as a screening test for beta thalassemia trait, as it is not practical or feasible to employ HbA2 in every case of anemia in childhood.

#### **MENTZERS INDEX:**

1.The Mentzer index, helpful in differentiating iron deficiency anemia from beta thalassemia. If a CBC indicates microcytic anemia, the Mentzer index is said to be a method of distinguishing between them.

2.If the index is less than 13, thalassemia is said to be more likely. If the result is greater than 13, then iron-deficiency anemia is said to be more



DR.VINAY CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY)





# P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

**A PIONEER DIAGNOSTIC CENTRE**

☎ 0171-2532620, 8222896961 ✉ [pkrajainhealthcare@gmail.com](mailto:pkrajainhealthcare@gmail.com)


<b>NAME</b>	: Mrs. HEENA DEVI	<b>PATIENT ID</b>	: 1603799
<b>AGE/ GENDER</b>	: 30 YRS/FEMALE	<b>REG. NO./LAB NO.</b>	: 122409060005
<b>COLLECTED BY</b>	:	<b>REGISTRATION DATE</b>	: 06/Sep/2024 09:15 AM
<b>REFERRED BY</b>	:	<b>COLLECTION DATE</b>	: 06/Sep/2024 09:31AM
<b>BARCODE NO.</b>	: 12504522	<b>REPORTING DATE</b>	: 07/Sep/2024 05:53AM
<b>CLIENT CODE.</b>	: P.K.R JAIN HEALTHCARE INSTITUTE		
<b>CLIENT ADDRESS</b>	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

likely.

3.The principle involved is as follows: In iron deficiency, the marrow cannot produce as many RBCs and they are small (microcytic), so the RBC count and the MCV will both be low, and as a result, the index will be greater than 13. Conversely, in thalassemia, which is a disorder of globin synthesis, the number of RBC's produced is normal, but the cells are smaller and more fragile. Therefore, the RBC count is normal, but the MCV is low, so the index will be less than 13.

**NOTE:** In practice, the Mentzer index is not a reliable indicator and should not, by itself, be used to differentiate. In addition, it would be possible for a patient with a microcytic anemia to have both iron deficiency and thalassemia, in which case the index would only suggest iron deficiency.



  
DR.VINAY CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

  
DR.YUGAM CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY)





# P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

**A PIONEER DIAGNOSTIC CENTRE**

☎ 0171-2532620, 8222896961 ✉ [pkrajainhealthcare@gmail.com](mailto:pkrajainhealthcare@gmail.com)

TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT.

<b>NAME</b>	: Mrs. HEENA DEVI	<b>PATIENT ID</b>	: 1603799
<b>AGE/ GENDER</b>	: 30 YRS/FEMALE	<b>REG. NO./LAB NO.</b>	: 122409060005
<b>COLLECTED BY</b>	:	<b>REGISTRATION DATE</b>	: 06/Sep/2024 09:15 AM
<b>REFERRED BY</b>	:	<b>COLLECTION DATE</b>	: 06/Sep/2024 09:31AM
<b>BARCODE NO.</b>	: 12504522	<b>REPORTING DATE</b>	: 06/Sep/2024 01:26PM
<b>CLIENT CODE.</b>	: P.K.R JAIN HEALTHCARE INSTITUTE		
<b>CLIENT ADDRESS</b>	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

## CLINICAL CHEMISTRY/BIOCHEMISTRY

### GLUCOSE RANDOM (R)

GLUCOSE RANDOM (R): PLASMA

74.27

mg/dL

by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD)

NORMAL: < 140.00

PREDIABETIC: 140.0 - 200.0

DIABETIC: > OR = 200.0

#### INTERPRETATION

##### IN ACCORDANCE WITH AMERICAN DIABETES ASSOCIATION GUIDELINES:

1. A random plasma glucose level below 140 mg/dl is considered normal.
2. A random glucose level between 140 - 200 mg/dl is considered as glucose intolerant or prediabetic. A fasting and post-prandial blood test (after consumption of 75 gms of glucose) is recommended for all such patients.
3. A random glucose level of above 200 mg/dl is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for all such patients. A fasting plasma glucose level in excess of 125 mg/dl on both occasions is confirmatory for diabetic state.



DR.VINAY CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY)







# P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

**A PIONEER DIAGNOSTIC CENTRE**

☎ 0171-2532620, 8222896961 ✉ pkrjainhealthcare@gmail.com

TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT.

**NAME** : Mrs. HEENA DEVI  
**AGE/ GENDER** : 30 YRS/FEMALE  
**COLLECTED BY** :  
**REFERRED BY** :  
**BARCODE NO.** : 12504522  
**CLIENT CODE.** : P.K.R JAIN HEALTHCARE INSTITUTE  
**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**PATIENT ID** : 1603799  
**REG. NO./LAB NO.** : 122409060005  
**REGISTRATION DATE** : 06/Sep/2024 09:15 AM  
**COLLECTION DATE** : 06/Sep/2024 09:31AM  
**REPORTING DATE** : 06/Sep/2024 01:26PM

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

## ENDOCRINOLOGY

### THYROID FUNCTION TEST: TOTAL

TRIIODOTHYRONINE (T3): SERUM by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)	1.44	ng/mL	0.35 - 1.93
THYROXINE (T4): SERUM by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)	8.36	µgm/dL	4.87 - 12.60
THYROID STIMULATING HORMONE (TSH): SERUM by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)	1.39	µIU/mL	0.35 - 5.50

3rd GENERATION, ULTRA SENSITIVE

#### INTERPRETATION:

TSH levels are subject to circadian variation, reaching peak levels between 2-4 a.m and at a minimum between 6-10 pm. The variation is of the order of 50%. Hence time of the day has influence on the measured serum TSH concentrations. TSH stimulates the production and secretion of the metabolically active hormones, thyroxine (T4) and triiodothyronine (T3). Failure at any level of regulation of the hypothalamic-pituitary-thyroid axis will result in either underproduction (hypothyroidism) or overproduction (hyperthyroidism) of T4 and/or T3.

CLINICAL CONDITION	T3	T4	TSH
Primary Hypothyroidism:	Reduced	Reduced	Increased (Significantly)
Subclinical Hypothyroidism:	Normal or Low Normal	Normal or Low Normal	High
Primary Hyperthyroidism:	Increased	Increased	Reduced (at times undetectable)
Subclinical Hyperthyroidism:	Normal or High Normal	Normal or High Normal	Reduced

#### LIMITATIONS:-

- T3 and T4 circulates in reversibly bound form with Thyroid binding globulins (TBG), and to a lesser extent albumin and Thyroid binding Pre Albumin so conditions in which TBG and protein levels alter such as pregnancy, excess estrogens, androgens, anabolic steroids and glucocorticoids may falsely affect the T3 and T4 levels and may cause false thyroid values for thyroid function tests.
- Normal levels of T4 can also be seen in Hyperthyroid patients with :T3 Thyrotoxicosis, Decreased binding capacity due to hypoproteinemia or ingestion of certain drugs (eg: phenytoin, salicylates).
- Serum T4 levels in neonates and infants are higher than values in the normal adult, due to the increased concentration of TBG in neonate serum.
- TSH may be normal in central hypothyroidism, recent rapid correction of hyperthyroidism or hypothyroidism, pregnancy, phenytoin therapy.

TRIIODOTHYRONINE (T3)		THYROXINE (T4)		THYROID STIMULATING HORMONE (TSH)	
Age	Refferance Range (ng/mL)	Age	Refferance Range (µg/dL)	Age	Reference Range (µIU/mL)
0 - 7 Days	0.20 - 2.65	0 - 7 Days	5.90 - 18.58	0 - 7 Days	2.43 - 24.3
7 Days - 3 Months	0.36 - 2.59	7 Days - 3 Months	6.39 - 17.66	7 Days - 3 Months	0.58 - 11.00
3 - 6 Months	0.51 - 2.52	3 - 6 Months	6.75 - 17.04	3 Days - 6 Months	0.70 - 8.40



  
DR. VINAY CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

  
DR. YUGAM CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY)





# P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

**A PIONEER DIAGNOSTIC CENTRE**

☎ 0171-2532620, 8222896961 ✉ pkrjainhealthcare@gmail.com

**NAME** : Mrs. HEENA DEVI  
**AGE/ GENDER** : 30 YRS/FEMALE  
**COLLECTED BY** :  
**REFERRED BY** :  
**BARCODE NO.** : 12504522  
**CLIENT CODE.** : P.K.R JAIN HEALTHCARE INSTITUTE  
**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**PATIENT ID** : 1603799  
**REG. NO./LAB NO.** : 122409060005  
**REGISTRATION DATE** : 06/Sep/2024 09:15 AM  
**COLLECTION DATE** : 06/Sep/2024 09:31AM  
**REPORTING DATE** : 06/Sep/2024 01:26PM

Test Name		Value		Unit	Biological Reference interval
6 - 12 Months	0.74 - 2.40	6 - 12 Months	7.10 - 16.16	6 - 12 Months	0.70 - 7.00
1 - 10 Years	0.92 - 2.28	1 - 10 Years	6.00 - 13.80	1 - 10 Years	0.60 - 5.50
11- 19 Years	0.35 - 1.93	11 - 19 Years	4.87- 13.20	11 - 19 Years	0.50 - 5.50
> 20 years (Adults)	0.35 - 1.93	> 20 Years (Adults)	4.87 - 12.60	> 20 Years (Adults)	0.35- 5.50
RECOMMENDATIONS OF TSH LEVELS DURING PREGNANCY ( $\mu$ U/mL)					
1st Trimester		0.10 - 2.50			
2nd Trimester		0.20 - 3.00			
3rd Trimester		0.30 - 4.10			

#### INCREASED TSH LEVELS:


- 1.Primary or untreated hypothyroidism may vary from 3 times to more than 100 times normal depending upon degree of hypofunction.
- 2.Hypothyroid patients receiving insufficient thyroid replacement therapy.
- 3.Hashimotos thyroiditis
- 4.DRUGS: Amphetamines, idonie containing agents & dopamine antagonist.
- 5.Neonatal period, increase in 1st 2-3 days of life due to post-natal surge

#### DECREASED TSH LEVELS:

- 1.Toxic multi-nodular goitre & Thyroiditis.
- 2.Over replacement of thyroid hormone in treatment of hypothyroidism.
- 3.Autonomously functioning Thyroid adenoma
- 4.Secondary pituitary or hypothalamic hypothyroidism
- 5.Acute psychiatric illness
- 6.Severe dehydration.
- 7.DRUGS: Glucocorticoids, Dopamine, Levodopa, T4 replacement therapy, Anti-thyroid drugs for thyrotoxicosis.
- 8.Pregnancy: 1st and 2nd Trimester



  
DR.VINAY CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

  
DR.YUGAM CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY)





# P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

**A PIONEER DIAGNOSTIC CENTRE**

☎ 0171-2532620, 8222896961

✉ pkrjainhealthcare@gmail.com

NAME	: Mrs. HEENA DEVI	PATIENT ID	: 1603799
AGE/ GENDER	: 30 YRS/FEMALE	REG. NO./LAB NO.	: 122409060005
COLLECTED BY	:	REGISTRATION DATE	: 06/Sep/2024 09:15 AM
REFERRED BY	:	COLLECTION DATE	: 06/Sep/2024 09:31AM
BARCODE NO.	: 12504522	REPORTING DATE	: 06/Sep/2024 01:26PM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE		
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

## IMMUNOPATHOLOGY/SEROLOGY HEPATITIS C VIRUS (HCV) ANTIBODIES SCREENING

HEPATITIS C ANTIBODY (HCV) TOTAL  
RESULT

NON - REACTIVE

by IMMUNOCHROMATOGRAPHY

### INTERPRETATION:

1.Anti HCV total antibody assay identifies presence IgG antibodies in the serum . It is a useful screening test with a specificity of nearly 99%.  
2.It becomes positive approximately 24 weeks after exposure. The test can not isolate an active ongoing HCV infection from an old infection that has been cleared. All positive results must be confirmed for active disease by an HCV PCR test .

### FALSE NEGATIVE RESULTS SEEN IN:

- 1.Window period
- 2.Immunocompromised states.

\*\*\* End Of Report \*\*\*



DR.VINAY CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY & MICROBIOLOGY)


DR.YUGAM CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS , MD (PATHOLOGY)



# Patient report

Bio-Rad  
D-10  
S/N: #DJ6F040603  
Sample ID:  
Injection date  
Injection #: 9  
Rack #: ---

DATE: 09/06/2024  
TIME: 05:53 PM  
Software version: 4.30-2  
12504522  
09/06/2024 04:40 PM  
Method: HbA2/F  
Rack position: 9



Peak table - ID: 12504522

Peak	R.time	Height	Area	Area %
A1a	0.20	4124	19930	1.1
A1b	0.29	4484	17127	0.9
F	0.46	1466	10634	< 0.8 *
LA1c/CHb-1	0.75	1781	15661	0.9
A1c	0.95	6448	68964	5.7
P3	1.54	12884	98156	5.3
A0	1.74	351020	1564914	85.1
A2	3.33	3029	44258	2.7
Total Area:		1839644		

Concentration:	%
F	< 0.8 *
A1c	5.7
A2	2.7