



P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

☎ 0171-2532620, 8222896961 ✉ pkrajainhealthcare@gmail.com

NAME : Mrs. KAWALJEET KAUR
AGE/ GENDER : 30 YRS/FEMALE
COLLECTED BY :
REFERRED BY :
BARCODE NO. : 12504549
CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE
CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

PATIENT ID : 1562189
REG. NO./LAB NO. : 122409070016
REGISTRATION DATE : 07/Sep/2024 11:43 AM
COLLECTION DATE : 07/Sep/2024 11:44 AM
REPORTING DATE : 07/Sep/2024 01:06 PM

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

HAEMATOLOGY

ERYTHROCYTE SEDIMENTATION RATE (ESR)

ERYTHROCYTE SEDIMENTATION RATE (ESR) by MODIFIED WESTERGREN AUTOMATED METHOD	53 ^H	mm/1st hr	0 - 20
---	-----------------	-----------	--------

INTERPRETATION:

1. ESR is a non-specific test because an elevated result often indicates the presence of inflammation associated with infection, cancer and autoimmune disease, but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it.
2. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other test such as C-reactive protein
3. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as some others, such as systemic lupus erythematosus

CONDITION WITH LOW ESR

A low ESR can be seen with conditions that inhibit the normal sedimentation of red blood cells, such as a high red blood cell count (polycythaemia), significantly high white blood cell count (leucocytosis), and some protein abnormalities. Some changes in red cell shape (such as sickle cells in sickle cell anaemia) also lower the ESR.

NOTE:

1. ESR and C - reactive protein (C-RP) are both markers of inflammation.
2. Generally, ESR does not change as rapidly as does CRP, either at the start of inflammation or as it resolves.
3. **CRP is not affected by as many other factors as is ESR, making it a better marker of inflammation.**
4. If the ESR is elevated, it is typically a result of two types of proteins, globulins or fibrinogen.
5. Women tend to have a higher ESR, and menstruation and pregnancy can cause temporary elevations.
6. Drugs such as dextran, methyldopa, oral contraceptives, penicillamine procainamide, theophylline, and vitamin A can increase ESR, while aspirin, cortisone, and quinine may decrease it



DR.VINAY CHOPRA

CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA

CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY)





P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

☎ 0171-2532620, 8222896961 ✉ pkrajainhealthcare@gmail.com

NAME : Mrs. KAWALJEET KAUR
AGE/ GENDER : 30 YRS/FEMALE
COLLECTED BY :
REFERRED BY :
BARCODE NO. : 12504549
CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE
CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

PATIENT ID : 1562189
REG. NO./LAB NO. : 122409070016
REGISTRATION DATE : 07/Sep/2024 11:43 AM
COLLECTION DATE : 07/Sep/2024 11:44AM
REPORTING DATE : 09/Sep/2024 07:52PM

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

SPECIAL INVESTIGATIONS

ANTI NUCLEAR ANTIBODY/FACTOR (ANA/ANF) - WITH REFLEX TO TITRES: IFA (HEP-2)

ANTI NUCLEAR ANTIBODY (ANA) - IFA, HEP2
by IFA (IMMUNO FLUORESCENT ASSAY)

NEGATIVE (-ve) NEGATIVE (-ve)

INTERPRETATION:

- 1.Anti Nuclear antibody (ANA) in dilutions is recommended for all positive results and follow up
- 2.Immunofluorescence microscopy using human cellular extracts like HEP-2 cells is a sensitive test for detection of serum antibodies that react specifically with various cellular proteins and nucleic acids
- 3.Test conducted on Serum

INTERPRETATION GUIDELINES : (Sample screening Dilution - 1:100):

Negative : No Immunofluorescence
+ : Weak Positive (1:100)
++ : Moderate Positive (1:320)
+++ : Strong Positive (1:1000)
++++ : Very strong Positive (1:3200)

COMMENTS:

Anti Nuclear antibody (ANA / ANF) is a group of autoantibodies directed against constituents of cell nuclei including DNA, RNA & various nuclear proteins. These autoantibodies are found with high frequency in patients with connective tissue disorders specially SLE. Since positive ANA results have been reported in healthy individuals, these reactivities are not by themselves diagnostic but must be correlated with other laboratory and clinical findings.

PATTERN	DISEASE ASSOCIATION
NUCLEAR	
Homogenous	SLE & other connective tissue disorders, Drug induced SLE
Peripheral	SLE & other connective tissue disorders
Speckled Coarse	Mixed connective Tissue Disorders (MCTD), Scleroderma-Polymyositis Overlap Syndrome, Raynauds Phenomenon, Psoriasis, Sjogrens Syndrome, Systemic Sclerosis.




DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)


DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY)





P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

☎ 0171-2532620, 8222896961 ✉ pkrjainhealthcare@gmail.com

NAME : Mrs. KAWALJEET KAUR
AGE/ GENDER : 30 YRS/FEMALE
COLLECTED BY :
REFERRED BY :
BARCODE NO. : 12504549
CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE
CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

PATIENT ID : 1562189
REG. NO./LAB NO. : 122409070016
REGISTRATION DATE : 07/Sep/2024 11:43 AM
COLLECTION DATE : 07/Sep/2024 11:44AM
REPORTING DATE : 09/Sep/2024 07:52PM

Test Name	Value	Unit	Biological Reference interval
Speckled Fine	SLE,Sjogrens syndrome,Scleroderma,Myositis,MCTD		
NUCLEAR DOTS			
Few	Auto-immune & Viral disease- Primary Biliay Cirrhosis & Chronic Active Hepatitis, Rarely Collagen Vascular disease		
Multiple	Primary Biliary Cirrhosis (>30%)		
Centromere	CREST syndrome, Progresive Systemic Sclerosis		
NUCLEOLAR			
Homogeneous	Scleroderma, Myositis, Raynauds Phenomena, SLE & Rheumatoid arthritis		
Clumpy	Systemic sclerosis & Scleroderma		
CYTOPLASMIC			
Mitochondrial	Primary Biliary Cirrhosis,Scleroderma & Overlap syndrome		
Ribosomal	SLE (10-20%)		

*** End Of Report ***



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY)

