

PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

NAME : Mr. VIKAS

AGE/ GENDER : 44 YRS/MALE **PATIENT ID** : 1449906

COLLECTED BY : 122409080003 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 08/Sep/2024 11:16 AM BARCODE NO. : 12504556 **COLLECTION DATE** : 08/Sep/2024 11:26AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 08/Sep/2024 12:26PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Value Unit **Biological Reference interval** Test Name

CLINICAL CHEMISTRY/BIOCHEMISTRY **GLUCOSE FASTING (F)**

GLUCOSE FASTING (F): PLASMA 102.33^H mg/dL NORMAL: < 100.0

by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD) PREDIABETIC: 100.0 - 125.0 **DIABETIC:** > **0R** = **126.0**

IN ACCORDANCE WITH AMERICAN DIABETES ASSOCIATION GUIDELINES:

1. A fasting plasma glucose level below 100 mg/dl is considered normal.

2. A fasting plasma glucose level between 100 - 125 mg/dl is considered as glucose intolerant or prediabetic. A fasting and post-prandial blood test (after consumption of 75 gms of glucose) is recommended for all such patients.

3. A fasting plasma glucose level of above 125 mg/dl is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for all such patients. A fasting plasma glucose level in excess of 125 mg/dl on both occasions is confirmatory for diabetic state.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)





PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

REPORTING DATE

:08/Sep/2024 12:26PM

NAME : Mr. VIKAS

AGE/ GENDER : 44 YRS/MALE **PATIENT ID** : 1449906

COLLECTED BY : 122409080003 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 08/Sep/2024 11:16 AM BARCODE NO. : 12504556 **COLLECTION DATE** : 08/Sep/2024 11:26AM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

Test Name Value Unit **Biological Reference interval**

URIC ACID

URIC ACID: SERUM 5.76 mg/dL 3.60 - 7.70

by URICASE - OXIDASE PEROXIDASE

INTERPRETATION:-

CLIENT CODE.

1.GOUT occurs when high levels of Uric Acid in the blood cause crystals to form & accumulate around a joint.

2.Uric Acid is the end product of purine metabolism. Uric acid is excreted to a large degree by the kidneys and to a smaller degree in the intestinal tract by microbial degradation.

INCREASED:

(A).DUE TO INCREASED PRODUCTION:-

1. Idiopathic primary gout.

2. Excessive dietary purines (organ meats, legumes, anchovies, etc).

3. Cytolytic treatment of malignancies especially leukemais & lymphomas.

4. Polycythemai vera & myeloid metaplasia.

5.Psoriasis.

6. Sickle cell anaemia etc.

(B).DUE TO DECREASED EXCREATION (BY KIDNEYS)

- 1. Alcohol ingestion.
- 2. Thiazide diuretics
- 3.Lactic acidosis.
- 4. Aspirin ingestion (less than 2 grams per day).
- 5. Diabetic ketoacidosis or starvation.
- 6.Renal failure due to any cause etc.

DECREASED:-

(A).DUE TO DIETARY DEFICIENCY

- 1. Dietary deficiency of Zinc, Iron and molybdenum.
- 2. Fanconi syndrome & Wilsons disease.
- 3. Multiple sclerosis.
- 4. Syndrome of inappropriate antidiuretic hormone (SIADH) secretion & low purine diet etc.

(B).DUE TO INCREASED EXCREATION

1.Drugs:-Probenecid, sulphinpyrazone, aspirin doses (more than 4 grams per day), corticosterroids and ACTH, anti-coagulants and estrogens etc.

*** End Of Report ***



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)

