



# P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

**A PIONEER DIAGNOSTIC CENTRE**

☎ 0171-2532620, 8222896961

✉ pkrjainhealthcare@gmail.com

**NAME** : Mr. GURSEWAK  
**AGE/ GENDER** : 44 YRS/MALE  
**COLLECTED BY** :  
**REFERRED BY** :  
**BARCODE NO.** : 12504559  
**CLIENT CODE.** : P.K.R JAIN HEALTHCARE INSTITUTE  
**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**PATIENT ID** : 1606513  
**REG. NO./LAB NO.** : 122409090002  
**REGISTRATION DATE** : 09/Sep/2024 08:21 AM  
**COLLECTION DATE** : 09/Sep/2024 08:23AM  
**REPORTING DATE** : 09/Sep/2024 01:25PM

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

## CLINICAL CHEMISTRY/BIOCHEMISTRY

### TRIGLYCERIDES

**TRIGLYCERIDES: SERUM**  
by GLYCEROL PHOSPHATE OXIDASE (ENZYMATIC)

324.43<sup>H</sup>

mg/dL

**OPTIMAL:** < 150.0  
**BORDERLINE HIGH:** 150.0 - 199.0  
**HIGH:** 200.0 - 499.0  
**VERY HIGH:** > OR = 500.0

#### INTERPRETATION:

NCEP RECOMMENDATIONS	TRIGLYCERIDES IN ADULTS (mg/dL)
DESIRABLE	< 150.0
BORDERLINE HIGH	150.0 – 199.0
HIGH	200.0 – 499.0
VERY HIGH	>OR = 500.0

#### NOTE

- Measurements in the same patient can show physiological variations. Three serial samples 1 week apart are recommended to establish basal triglyceride levels.
- Certain conditions such as acute illness, stress, pregnancy, dietary changes especially changes in intake of saturated fatty acids, lipid lowering drugs, alcohol or prednisone may cause variation in lipid levels.

#### COMMENTS

National Lipid association - 2014 identifies elevated Triglycerides as an independent risk factor for Coronary Heart Disease (CHD).

\*\*\* End Of Report \*\*\*



DR.VINAY CHOPRA

CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA

CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY)

