CLIENT CODE.



PKR JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

REPORTING DATE

: 14/Sep/2024 03:54PM

NAME : Mr. SHANTANU

AGE/ GENDER : 30 YRS/MALE **PATIENT ID** : 1612778

COLLECTED BY : 122409140028 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 14/Sep/2024 11:37 AM BARCODE NO. : 12504705 **COLLECTION DATE** : 14/Sep/2024 11:40AM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

Test Name Value Unit **Biological Reference interval**

SWASTHYA WELLNESS PANEL: 1.5 COMPLETE BLOOD COUNT (CBC)

RED BLOOD CELLS (RBCS) COUNT AND INDICES

HAEMOGLOBIN (HB) by CALORIMETRIC	14.1	gm/dL	12.0 - 17.0
RED BLOOD CELL (RBC) COUNT by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	4.53	Millions/cmm	3.50 - 5.00
PACKED CELL VOLUME (PCV) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	40.7	%	40.0 - 54.0
MEAN CORPUSCULAR VOLUME (MCV) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	90	fL	80.0 - 100.0
MEAN CORPUSCULAR HAEMOGLOBIN (MCH) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	31.2	pg	27.0 - 34.0
MEAN CORPUSCULAR HEMOGLOBIN CONC. (MCHC) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	34.7	g/dL	32.0 - 36.0
RED CELL DISTRIBUTION WIDTH (RDW-CV) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	12	%	11.00 - 16.00
RED CELL DISTRIBUTION WIDTH (RDW-SD) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	40.9	fL	35.0 - 56.0
MENTZERS INDEX by CALCULATED	19.87	RATIO	BETA THALASSEMIA TRAIT: < 13.0 IRON DEFICIENCY ANEMIA: >13.0
GREEN & KING INDEX by CALCULATED	23.9	RATIO	BETA THALASSEMIA TRAIT:<= 65.0 IRON DEFICIENCY ANEMIA: > 65.0
WHITE BLOOD CELLS (WBCS)			
TOTAL LEUCOCYTE COUNT (TLC) by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY DIFFERENTIAL LEUCOCYTE COUNT (DLC)	7160	/cmm	4000 - 11000
NEUTROPHILS by Flow cytometry by SF cube & microscopy	53	%	50 - 70
LYMPHOCYTES by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	33	%	20 - 40
EOSINOPHILS by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	6 ^H	%	1 - 6



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)







A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

NAME : Mr. SHANTANU

AGE/ GENDER : 30 YRS/MALE **PATIENT ID** : 1612778

COLLECTED BY : 122409140028 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 14/Sep/2024 11:37 AM BARCODE NO. : 12504705 **COLLECTION DATE** : 14/Sep/2024 11:40AM

CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 14/Sep/2024 03:54PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name	Value	Unit	Biological Reference interval
MONOCYTES by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	8	%	2 - 12
BASOPHILS by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY ABSOLUTE LEUKOCYTES (WBC) COUNT	0	%	0 - 1
ABSOLUTE NEUTROPHIL COUNT by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	3795	/cmm	2000 - 7500
ABSOLUTE LYMPHOCYTE COUNT by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	2363 ^L	/cmm	800 - 4900
ABSOLUTE EOSINOPHIL COUNT by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	430 ^H	/cmm	40 - 440
ABSOLUTE MONOCYTE COUNT by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	573	/cmm	80 - 880
ABSOLUTE BASOPHIL COUNT by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	0	/cmm	0 - 110
PLATELETS AND OTHER PLATELET PREDICTIVE MARKI	ERS.		
PLATELET COUNT (PLT) by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	262000	/cmm	150000 - 450000
PLATELETCRIT (PCT) by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	0.28	%	0.10 - 0.36
MEAN PLATELET VOLUME (MPV) by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	11	fL	6.50 - 12.0
PLATELET LARGE CELL COUNT (P-LCC) by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	83000	/cmm	30000 - 90000
PLATELET LARGE CELL RATIO (P-LCR) by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	31.7	%	11.0 - 45.0
PLATELET DISTRIBUTION WIDTH (PDW) by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE NOTE: TEST CONDUCTED ON EDTA WHOLE BLOOD	16.3	%	15.0 - 17.0



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS, MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST





A PIONEER DIAGNOSTIC CENTRE

: 14/Sep/2024 05:39PM

NAME : Mr. SHANTANU

AGE/ GENDER : 30 YRS/MALE **PATIENT ID** : 1612778

COLLECTED BY : 122409140028 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 14/Sep/2024 11:37 AM BARCODE NO. : 12504705 **COLLECTION DATE** : 14/Sep/2024 11:40AM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

Test Name Value Unit **Biological Reference interval**

REPORTING DATE

GLYCOSYLATED HAEMOGLOBIN (HBA1C)

5.6 GLYCOSYLATED HAEMOGLOBIN (HbA1c): 4.0 - 6.4

WHOLF BLOOD

CLIENT CODE.

by HPLC (HIGH PERFORMANCE LIQUID CHROMATOGRAPHY)

ESTIMATED AVERAGE PLASMA GLUCOSE 114.02 mg/dL 60.00 - 140.00

by HPLC (HIGH PERFORMANCE LIQUID CHROMATOGRAPHY)

INTERPRETATION:

AS PER AMERICAN DIABETES ASSOCIATION (ADA):			
REFERENCE GROUP	GLYCOSYLATED HEMOGLOGIB (HBAIC) in %		
Non diabetic Adults >= 18 years	<5.7		
At Risk (Prediabetes)	5.7 – 6.4		
Diagnosing Diabetes	>= 6.5		
	Age > 19 Years		
	Goals of Therapy:	< 7.0	
Therapeutic goals for glycemic control	Actions Suggested:	>8.0	
	Age < 19 Y	ears	
	Goal of therapy:	<7.5	

COMMENTS:

- 1.Glycosylated hemoglobin (HbA1c) test is three monthly monitoring done to assess compliace with therapeutic regimen in diabetic patients. 2. Since Hb1c reflects long term fluctuations in blood glucose concentration, a diabetic patient who has recently under good control may still have high concentration of HbAlc. Converse is true for a diabetic previously under good control but now poorly controlled.
- 3. Target goals of < 7.0 % may be beneficial in patients with short duration of diabetes, long life expectancy and no significant cardiovascular disease. In patients with significant complications of diabetes, limited life expectancy or extensive co-morbid conditions, targetting a goal of < 7.0% may not be
- 4.High HbA1c (>9.0 -9.5 %) is strongly associated with risk of development and rapid progression of microvascular and nerve complications 5. Any condition that shorten RBC life span like acute blood loss, hemolytic anemia falsely lower HbA1c results.
- 6.HbA1c results from patients with HbSS,HbSC and HbD must be interpreted with caution, given the pathological processes including anemia,increased red cell turnover, and transfusion requirement that adversely impact HbA1c as a marker of long-term gycemic control.
- 7. Specimens from patients with polycythemia or post-splenctomy may exhibit increse in HbA1c values due to a somewhat longer life span of the red cells

CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)







A PIONEER DIAGNOSTIC CENTRE

NAME : Mr. SHANTANU

AGE/ GENDER : 30 YRS/MALE **PATIENT ID** : 1612778

COLLECTED BY REG. NO./LAB NO. : 122409140028

REFERRED BY **REGISTRATION DATE** : 14/Sep/2024 11:37 AM BARCODE NO. **COLLECTION DATE** : 14/Sep/2024 11:40AM : 12504705

CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 14/Sep/2024 04:41PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name Value Unit **Biological Reference interval**

ERYTHROCYTE SEDIMENTATION RATE (ESR)

ERYTHROCYTE SEDIMENTATION RATE (ESR)

0 - 20

by MODIFIED WESTERGREN AUTOMATED METHOD

INTERPRETATION:

- 1. ESR is a non-specific test because an elevated result often indicates the presence of inflammation associated with infection, cancer and autoimmune disease, but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it.
- 2. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other test such as C-reactive protein
- 3. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as some others, such as systemic lupus erythematosus
 CONDITION WITH LOW ESR

A low ESR can be seen with conditions that inhibit the normal sedimentation of red blood cells, such as a high red blood cell count (polycythaemia), significantly high white blood cell count (leucocytosis), and some protein abnormalities. Some changes in red cell shape (such as sickle cells in sickle cell anaemia) also lower the ESR.

- ESR and C reactive protein (C-RP) are both markers of inflammation.
 Generally, ESR does not change as rapidly as does CRP, either at the start of inflammation or as it resolves.
 CRP is not affected by as many other factors as is ESR, making it a better marker of inflammation.
 If the ESR is elevated, it is typically a result of two types of proteins, globulins or fibrinogen.
- 5. Women tend to have a higher ESR, and menstruation and pregnancy can cause temporary elevations.
- 6. Drugs such as dextran, methyldopa, oral contraceptives, penicillamine procainamide, theophylline, and vitamin A can increase ESR, while aspirin, cortisone, and quinine may decrease it



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)







A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

REPORTING DATE

: 14/Sep/2024 03:54PM

NAME : Mr. SHANTANU

AGE/ GENDER : 30 YRS/MALE **PATIENT ID** : 1612778

COLLECTED BY : 122409140028 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 14/Sep/2024 11:37 AM BARCODE NO. : 12504705 **COLLECTION DATE** : 14/Sep/2024 11:40AM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

Test Name Value Unit **Biological Reference interval**

CLINICAL CHEMISTRY/BIOCHEMISTRY GLUCOSE FASTING (F)

87.47 GLUCOSE FASTING (F): PLASMA mg/dL NORMAL: < 100.0

by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD) PREDIABETIC: 100.0 - 125.0 DIABETIC: > 0R = 126.0

CLIENT CODE.

IN ACCORDANCE WITH AMERICAN DIABETES ASSOCIATION GUIDELINES:

1. A fasting plasma glucose level below 100 mg/dl is considered normal.

2. A fasting plasma glucose level between 100 - 125 mg/dl is considered as glucose intolerant or prediabetic. A fasting and post-prandial blood test (after consumption of 75 gms of glucose) is recommended for all such patients.

3. A fasting plasma glucose level of above 125 mg/dl is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for all such patients. A fasting plasma glucose level in excess of 125 mg/dl on both occasions is confirmatory for diabetic state.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)





A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

NAME : Mr. SHANTANU

AGE/ GENDER : 30 YRS/MALE **PATIENT ID** : 1612778

COLLECTED BY : 122409140028 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 14/Sep/2024 11:37 AM BARCODE NO. : 12504705 **COLLECTION DATE** : 14/Sep/2024 11:40AM

CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 14/Sep/2024 03:54PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name	Value	Unit	Biological Reference interval
	LIPID PROFILE	: BASIC	
CHOLESTEROL TOTAL: SERUM by CHOLESTEROL OXIDASE PAP	144.61	mg/dL	OPTIMAL: < 200.0 BORDERLINE HIGH: 200.0 - 239.0 HIGH CHOLESTEROL: > OR = 240.0
TRIGLYCERIDES: SERUM by GLYCEROL PHOSPHATE OXIDASE (ENZYMATIC)	114.84	mg/dL	OPTIMAL: < 150.0 BORDERLINE HIGH: 150.0 - 199.0 HIGH: 200.0 - 499.0 VERY HIGH: > OR = 500.0
HDL CHOLESTEROL (DIRECT): SERUM by SELECTIVE INHIBITION	40.21	mg/dL	LOW HDL: < 30.0 BORDERLINE HIGH HDL: 30.0 - 60.0 HIGH HDL: > OR = 60.0
LDL CHOLESTEROL: SERUM by CALCULATED, SPECTROPHOTOMETRY	81.43	mg/dL	OPTIMAL: < 100.0 ABOVE OPTIMAL: 100.0 - 129.0 BORDERLINE HIGH: 130.0 - 159.0 HIGH: 160.0 - 189.0 VERY HIGH: > OR = 190.0
NON HDL CHOLESTEROL: SERUM by CALCULATED, SPECTROPHOTOMETRY	104.4	mg/dL	OPTIMAL: < 130.0 ABOVE OPTIMAL: 130.0 - 159.0 BORDERLINE HIGH: 160.0 - 189.0 HIGH: 190.0 - 219.0 VERY HIGH: > OR = 220.0
VLDL CHOLESTEROL: SERUM by CALCULATED, SPECTROPHOTOMETRY	22.97	mg/dL	0.00 - 45.00
TOTAL LIPIDS: SERUM by CALCULATED, SPECTROPHOTOMETRY	404.06	mg/dL	350.00 - 700.00
CHOLESTEROL/HDL RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	3.6	RATIO	LOW RISK: 3.30 - 4.40 AVERAGE RISK: 4.50 - 7.0 MODERATE RISK: 7.10 - 11.0 HIGH RISK: > 11.0
LDL/HDL RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	2.03	RATIO	LOW RISK: 0.50 - 3.0 MODERATE RISK: 3.10 - 6.0 HIGH RISK: > 6.0



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)







A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

REPORTING DATE

: 14/Sep/2024 03:54PM

NAME : Mr. SHANTANU

AGE/ GENDER : 30 YRS/MALE **PATIENT ID** : 1612778

COLLECTED BY : 122409140028 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 14/Sep/2024 11:37 AM BARCODE NO. **COLLECTION DATE** : 14/Sep/2024 11:40AM : 12504705

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

Test Name Value Unit **Biological Reference interval**

TRIGLYCERIDES/HDL RATIO: SERUM **RATIO** 3.00 - 5.002.86^L by CALCULATED, SPECTROPHOTOMETRY

INTERPRETATION:

CLIENT CODE.

1. Measurements in the same patient can show physiological analytical variations. Three serial samples 1 week apart are recommended for Total Cholesterol, Triglycerides, HDL & LDL Cholesterol.

2. As per NLA-2014 guidelines, all adults above the age of 20 years should be screened for lipid status. Selective screening of children above the age of 2 years with a family history of premature cardiovascular disease or those with at least one parent with high total cholesterol is recommended.

3. Low HDL levels are associated with increased risk for Atherosclerotic Cardiovascular disease (ASCVD) due to insufficient HDL being available

to participate in reverse cholesterol transport, the process by which cholesterol is eliminated from peripheral tissues.

4. NLA-2014 identifies Non HDL Cholesterol (an indicator of all atherogeniclipoproteins such as LDL, VLDL, IDL, Lpa, Chylomicron remnants) along with LDL-cholesterol as co-primary target for cholesterol lowering therapy. Note that major risk factors can modify treatment goals for LDL &Non

5. Additional testing for Apolipoprotein B, hsCRP,Lp(a) & LP-PLA2 should be considered among patients with moderate risk for ASCVD for risk refinement



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)





A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

NAME : Mr. SHANTANU

AGE/ GENDER : 30 YRS/MALE **PATIENT ID** : 1612778

COLLECTED BY : 122409140028 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 14/Sep/2024 11:37 AM BARCODE NO. : 12504705 **COLLECTION DATE** : 14/Sep/2024 11:40AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 14/Sep/2024 03:54PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name Value Unit **Biological Reference interval**

LIVER FUNCTION TEST (COMPLETE)

BILIRUBIN TOTAL: SERUM by DIAZOTIZATION, SPECTROPHOTOMETRY	0.68	mg/dL	INFANT: 0.20 - 8.00 ADULT: 0.00 - 1.20
BILIRUBIN DIRECT (CONJUGATED): SERUM by DIAZO MODIFIED, SPECTROPHOTOMETRY	0.24	mg/dL	0.00 - 0.40
BILIRUBIN INDIRECT (UNCONJUGATED): SERUM by CALCULATED, SPECTROPHOTOMETRY	0.44	mg/dL	0.10 - 1.00
SGOT/AST: SERUM by IFCC, WITHOUT PYRIDOXAL PHOSPHATE	25.72	U/L	7.00 - 45.00
SGPT/ALT: SERUM by IFCC, WITHOUT PYRIDOXAL PHOSPHATE	40.31	U/L	0.00 - 49.00
AST/ALT RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	0.64	RATIO	0.00 - 46.00
ALKALINE PHOSPHATASE: SERUM by Para Nitrophenyl Phosphatase by Amino Methyl PROPANOL	75.75	U/L	40.0 - 130.0
GAMMA GLUTAMYL TRANSFERASE (GGT): SERUM by SZASZ, SPECTROPHTOMETRY	15.79	U/L	0.00 - 55.0
TOTAL PROTEINS: SERUM by BIURET, SPECTROPHOTOMETRY	6.61	gm/dL	6.20 - 8.00
ALBUMIN: SERUM by BROMOCRESOL GREEN	4.44	gm/dL	3.50 - 5.50
GLOBULIN: SERUM by CALCULATED, SPECTROPHOTOMETRY	2.17 ^L	gm/dL	2.30 - 3.50
A : G RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	2.05 ^H	RATIO	1.00 - 2.00

INTERPRETATION

NOTE:- To be correlated in individuals having SGOT and SGPT values higher than Normal Referance Range.

USE:- Differential diagnosis of diseases of hepatobiliary system and pancreas.

INCREASED:

DRUG HEPATOTOXICITY	> 2
ALCOHOLIC HEPATITIS	> 2 (Highly Suggestive)
CIRRHOSIS	1.4 - 2.0
INTRAHEPATIC CHOLESTATIS	> 1.5
HEPATOCELLULAR CARCINOMA & CHRONIC HEPATITIS	> 1.3 (Slightly Increased)



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)







A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

NAME : Mr. SHANTANU

AGE/ GENDER : 30 YRS/MALE **PATIENT ID** : 1612778

COLLECTED BY : 122409140028 REG. NO./LAB NO.

: 14/Sep/2024 11:37 AM REFERRED BY **REGISTRATION DATE** BARCODE NO. : 12504705 **COLLECTION DATE** : 14/Sep/2024 11:40AM

CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 14/Sep/2024 03:54PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name Value Unit **Biological Reference interval**

DECREASED:

1. Acute Hepatitis due to virus, drugs, toxins (with AST increased 3 to 10 times upper limit of normal)

2. Extra Hepatic cholestatis: 0.8 (normal or slightly decreased).

PROGNOSTIC SIGNIFICANCE:

NORMAL	< 0.65
GOOD PROGNOSTIC SIGN	0.3 - 0.6
POOR PROGNOSTIC SIGN	1.2 - 1.6



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)





A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

NAME : Mr. SHANTANU

AGE/ GENDER : 30 YRS/MALE **PATIENT ID** : 1612778

COLLECTED BY : 122409140028 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 14/Sep/2024 11:37 AM BARCODE NO. : 12504705 **COLLECTION DATE** : 14/Sep/2024 11:40AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 14/Sep/2024 05:28PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

KIDNEY FUNCTION TEST (COMPLETE)

	KIDITET TOROTION TEST	(OOMI ELIL)	
UREA: SERUM	25.66	mg/dL	10.00 - 50.00
by UREASE - GLUTAMATE DEHYDROGENASE	,		
CREATININE: SERUM	0.64	mg/dL	0.40 - 1.40
by ENZYMATIC, SPECTROPHOTOMETERY			
BLOOD UREA NITROGEN (BUN): SERUM	11.99	mg/dL	7.0 - 25.0
by CALCULATED, SPECTROPHOTOMETRY			
BLOOD UREA NITROGEN (BUN)/CREATIN	INE 18.73	RATIO	10.0 - 20.0
RATIO: SERUM			
by CALCULATED, SPECTROPHOTOMETRY			
UREA/CREATININE RATIO: SERUM	40.09	RATIO	
by CALCULATED, SPECTROPHOTOMETRY	10.07	10,1110	
URIC ACID: SERUM	5.03	mg/dL	3.60 - 7.70
by URICASE - OXIDASE PEROXIDASE	3.03	Trig/ UL	3.00 - 7.70
CALCIUM: SERUM	8.74	mg/dL	8.50 - 10.60
by ARSENAZO III, SPECTROPHOTOMETRY	0.74	Trig/ uL	8.50 - 10.00
	2.01	ma/dl	2.20 4.70
PHOSPHOROUS: SERUM	2.81	mg/dL	2.30 - 4.70
by PHOSPHOMOLYBDATE, SPECTROPHOTON	METRY		
<u>ELECTROLYTES</u>			
SODIUM: SERUM	141.1	mmol/L	135.0 - 150.0
by ISE (ION SELECTIVE ELECTRODE)			
POTASSIUM: SERUM	4.46	mmol/L	3.50 - 5.00
by ISE (ION SELECTIVE ELECTRODE)	1.10	THITION E	0.00 0.00
CHLORIDE: SERUM	105.82	mmol/L	90.0 - 110.0
by ISE (ION SELECTIVE ELECTRODE)	100.02	THITIOI/ L	70.0 110.0
ESTIMATED GLOMERULAR FILTERATION	DATE		
ESTIMATED GLOMERULAR FILTERATION	RATE 130.6		
(eGFR): SERLIM			

(eGFR): SERUM by CALCULATED

INTERPRETATION:

To differentiate between pre- and post renal azotemia.

INCREASED RATIO (>20:1) WITH NORMAL CREATININE:

- 1. Prerenal azotemia (BUN rises without increase in creatinine) e.g. heart failure, salt depletion, dehydration, blood loss) due to decreased glomerular filtration rate.
- 2. Catabolic states with increased tissue breakdown.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)





A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

NAME : Mr. SHANTANU

AGE/ GENDER : 30 YRS/MALE **PATIENT ID** : 1612778

COLLECTED BY : 122409140028 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 14/Sep/2024 11:37 AM

BARCODE NO. : 12504705 **COLLECTION DATE** : 14/Sep/2024 11:40AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 14/Sep/2024 05:28PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name Value Unit **Biological Reference interval**

3. GI haemorrhage.

4. High protein intake.

5. Impaired renal function plus

6. Excess protein intake or production or tissue breakdown (e.g. infection, GI bleeding, thyrotoxicosis, Cushing's syndrome, high protein diet, burns, surgery, cachexia, high fever).

7. Urine reabsorption (e.g. ureter colostomy)

8. Reduced muscle mass (subnormal creatinine production)

9. Certain drugs (e.g. tetracycline, glucocorticoids)

INCREASED RATIO (>20:1) WITH ELEVATED CREATININE LEVELS:

- 1. Postrenal azotemia (BUN rises disproportionately more than creatinine) (e.g. obstructive uropathy).
- 2. Prerenal azotemia superimposed on renal disease.

DECREASED RATIO (<10:1) WITH DECREASED BUN:

- 1. Acute tubular necrosis.
- 2. Low protein diet and starvation.
- 3. Severe liver disease.
- 4. Other causes of decreased urea synthesis.
- 5. Repeated dialysis (urea rather than creatinine diffuses out of extracellular fluid).
- 6. Inherited hyperammonemias (urea is virtually absent in blood).
- 7. SIADH (syndrome of inappropiate antidiuretic harmone) due to tubular secretion of urea.
- 8. Pregnancy.

DECREASED RATIO (<10:1) WITH INCREASED CREATININE:

- 1. Phenacimide therapy (accelerates conversion of creatine to creatinine).
- 2. Rhabdomyolysis (releases muscle creatinine).
- 3. Muscular patients who develop renal failure.

INAPPROPIATE RATIO:

- 1. Diabetic ketoacidosis (acetoacetate causes false increase in creatinine with certain methodologies, resulting in normal ratio when dehydration should produce an increased BUN/creatinine ratio).
- 2. Cephalosporin therapy (interferes with creatinine measurement). ESTIMATED GLOMERULAR FILTERATION RATE:

STIMATED GEOMEROEAR TETERATION RATE.				
CKD STAGE	DESCRIPTION	GFR (mL/min/1.73m2)	ASSOCIATED FINDINGS	
G1	Normal kidney function	>90	No proteinuria	
G2	Kidney damage with	>90	Presence of Protein,	
	normal or high GFR		Albumin or cast in urine	
G3a	Mild decrease in GFR	60 -89		
G3b	Moderate decrease in GFR	30-59		
G4	Severe decrease in GFR	15-29		
G5	Kidney failure	<15		



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)







A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

REPORTING DATE

: 14/Sep/2024 05:28PM

NAME : Mr. SHANTANU

AGE/ GENDER : 30 YRS/MALE **PATIENT ID** : 1612778

COLLECTED BY REG. NO./LAB NO. : 122409140028

REFERRED BY **REGISTRATION DATE** : 14/Sep/2024 11:37 AM BARCODE NO. **COLLECTION DATE** : 14/Sep/2024 11:40AM : 12504705

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

Test Name Value Unit **Biological Reference interval**

COMMENTS:

CLIENT CODE.

1. Estimated Glomerular filtration rate (eGFR) is the sum of filtration rates in all functioning nephrons and so an estimation of the GFR provides a measure of functioning nephrons of the kidney.

2. eGFR calculated using the 2009 CKD-EPI creatinine equation and GFR category reported as per KDIGO guideline 2012

3. In patients, with eGFR creatinine between 45-59 ml/min/1.73 m2 (G3) and without any marker of Kidney damage, It is recommended to measure eGFR with Cystatin C for confirmation of CKD

4. eGFR category G1 OR G2 does not fullfill the criteria for CKD, in the absence of evidence of Kidney Damage
5. In a suspected case of Acute Kidney Injury (AKI), measurement of eGFR should be done after 48-96 hours of any Intervention or procedure
6. eGFR calculated by Serum Creatinine may be less accurate due to certain factors like Race, Muscle Mass, Diet, Certain Drugs. In such cases, eGFR should be calculated using Serum Cystatin C
7. A decrease in eGFR implies either progressive renal disease, or a reversible process causing decreased nephron function (eg, severe dehydration).

KDIGO guideline, 2012 recommends Chronic Kidney Disease (CKD) should be classified based on cause, eGFR category and Albuminuria (ACR) category. GFR & ACR category combined together reflect risk of progression and helps Clinician to identify the individual who are progressing at more rapid rate than anticipated



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)





PKR JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

REPORTING DATE

: 14/Sep/2024 05:39PM

NAME : Mr. SHANTANU

AGE/ GENDER : 30 YRS/MALE **PATIENT ID** : 1612778

COLLECTED BY : 122409140028 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 14/Sep/2024 11:37 AM BARCODE NO. : 12504705 **COLLECTION DATE** : 14/Sep/2024 11:40AM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

Test Name	Value	Unit	Biological Reference interval
	IRON PROFILE		
	IKONTKOTILL		
IRON: SERUM	64.14	μg/dL	59.0 - 158.0
by FERROZINE, SPECTROPHOTOMETRY		. •	
UNSATURATED IRON BINDING CAPACITY (UIBC)	212.86	μg/dL	150.0 - 336.0
:SERUM	2.2.00	MO/ W=	
by FERROZINE, SPECTROPHOTOMETERY			
	277	/41	220 420
TOTAL IRON BINDING CAPACITY (TIBC)	277	μg/dL	230 - 430
:SERUM			
by SPECTROPHOTOMETERY			
%TRANSFERRIN SATURATION: SERUM	23.16	%	15.0 - 50.0
by CALCULATED, SPECTROPHOTOMETERY (FERENE)			
TRANSFERRIN: SERUM	196.67 ^L	mg/dL	200.0 - 350.0

by SPECTROPHOTOMETERY (FERENE) INTERPRETATION:-

CLIENT CODE.

MATERIA RESTAURANT			
VARIABLES	ANEMIA OF CHRONIC DISEASE	IRON DEFICIENCY ANEMIA	THALASSEMIA α/β TRAIT
SERUM IRON:	Normal to Reduced	Reduced	Normal
TOTAL IRON BINDING CAPACITY:	Decreased	Increased	Normal
% TRANSFERRIN SATURATION:	Decreased	Decreased < 12-15 %	Normal
SERUM FERRITIN:	Normal to Increased	Decreased	Normal or Increased

196.67L

IRON:

1. Serum iron studies is recommended for differential diagnosis of microcytic hypochromic anemia.i.e iron deficiency anemia, zinc deficiency

anemia, anemia of chronic disease and thalassemia syndromes.

2. It is essential to isolate iron deficiency anemia from Beta thalassemia syndromes because during iron replacement which is therapeutic for iron deficiency anemia, is severely contra-indicated in Thalassemia. TOTAL IRON BINDING CAPACITY (TIBC):

1. It is a direct measure of protein transferrin which transports iron from the gut to storage sites in the bone marrow.

% TRANSFERRIN SATURATION:

1.Occurs in idiopathic hemochromatosis and transfusional hemosiderosis where no unsaturated iron binding capacity is available for iron mobilization. Similar condition is seen in congenital deficiency of transferrin.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)







A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

NAME : Mr. SHANTANU

AGE/ GENDER : 30 YRS/MALE **PATIENT ID** : 1612778

COLLECTED BY : 122409140028 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 14/Sep/2024 11:37 AM BARCODE NO. : 12504705 **COLLECTION DATE** : 14/Sep/2024 11:40AM

CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 14/Sep/2024 03:54PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name Value Unit **Biological Reference interval**

ENDOCRINOLOGY

THYROID FUNCTION TEST: TOTAL

TRIIODOTHYRONINE (T3): SERUM 1.32 ng/mL 0.35 - 1.93by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

THYROXINE (T4): SERUM 4.87 - 12.60 6.71 μgm/dL

by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

THYROID STIMULATING HORMONE (TSH): SERUM 1.51 μIU/mL 0.35 - 5.50

by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

3rd GENERATION, ULTRASENSITIVE

INTERPRETATION:

TSH levels are subject to circadian variation, reaching peak levels between 2-4 a.m and at a minimum between 6-10 pm. The variation is of the order of 50%. Hence time of the day has influence on the measured serum TSH concentrations. TSH stimulates the production and secretion of the metabolically active hormones, thyroxine (T4) and trilodothyronine (T3). Failure at any level of regulation of the hypothalamic-pituitary-thyroid axis will result in either underproduction (hypothyroidism) or overproduction(hyperthyroidism) of T4 and/or T3.

CLINICAL CONDITION	Т3	T4	TSH
Primary Hypothyroidism:	Reduced	Reduced	Increased (Significantly)
Subclinical Hypothyroidism:	Normal or Low Normal	Normal or Low Normal	High
Primary Hyperthyroidism:	Increased	Increased	Reduced (at times undetectable)
Subclinical Hyperthyroidism:	Normal or High Normal	Normal or High Normal	Reduced

- 1. T3 and T4 circulates in reversibly bound form with Thyroid binding globulins (TBG), and to a lesser extent albumin and Thyroid binding Pre Albumin so conditions in which TBG and protein levels alter such as pregnancy, excess estrogens, androgens, anabolic steroids and glucocorticoids may falsely affect the T3 and T4 levels and may cause false thyroid values for thyroid function tests.
- 2. Normal levels of T4 can also be seen in Hyperthyroid patients with :T3 Thyrotoxicosis, Decreased binding capacity due to hypoproteinemia or ingestion of certain drugs
- 3. Serum T4 levies in neonates and infants are higher than values in the normal adult, due to the increased concentration of TBG in neonate serum.
- 4. TSH may be normal in central hypothyroidism, recent rapid correction of hyperthyroidism or hypothroidism, pregnancy, phenytoin therapy.

TRIIODOTHY	RONINE (T3)	THYROXINE (T4)		THYROID STIMULATING HORMONE (TSH)		
Age	Refferance Range (ng/mL)	Age	Refferance Range (μg/dL)	Age	Reference Range (μΙυ/mL)	
0-7 Days	0.20 - 2.65	0 - 7 Days	5.90 - 18.58	0 - 7 Days	2.43 - 24.3	
7 Days - 3 Months	0.36 - 2.59	7 Days - 3 Months	6.39 - 17.66	7 Days - 3 Months	0.58 - 11.00	
3 - 6 Months	0.51 - 2.52	3 - 6 Months	6.75 – 17.04	3 Days – 6 Months	0.70 - 8.40	



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)





A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

NAME : Mr. SHANTANU

AGE/ GENDER : 30 YRS/MALE **PATIENT ID** : 1612778

COLLECTED BY REG. NO./LAB NO. : 122409140028

REFERRED BY **REGISTRATION DATE** : 14/Sep/2024 11:37 AM BARCODE NO. : 12504705 **COLLECTION DATE** : 14/Sep/2024 11:40AM

CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 14/Sep/2024 03:54PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name			Value	Unit		Biologic	al Reference interval	
6 - 12 Months	0.74 - 2.40	6 - 12 Months	7.10 – 16.16	6 – 12 Months	0.70 - 7.00			
1 - 10 Years	0.92 - 2.28	1 - 10 Years	6.00 - 13.80	1 – 10 Years	0.60 - 5.50			
11- 19 Years	0.35 - 1.93	11 - 19 Years	4.87- 13.20	11 – 19 Years	0.50 - 5.50			
> 20 years (Adults)	0.35 - 1.93	> 20 Years (Adults)	4.87 - 12.60	> 20 Years (Adults)	0.35- 5.50			
RECOMMENDATIONS OF TSH LEVELS DURING PREGNANCY (µIU/mL)								
	1st Trimester		0.10 - 2.50			0.10 – 2.50		
	2nd Trimester		0.20 – 3.00					
	3rd Trimester		0.30 - 4.10					

INCREASED TSH LEVELS:

- 1. Primary or untreated hypothyroidism may vary from 3 times to more than 100 times normal depending upon degree of hypofunction.
- 2. Hypothyroid patients receiving insufficient thyroid replacement therapy.
- 3. Hashimotos thyroiditis
- 4.DRUGS: Amphetamines, idonie containing agents & dopamine antagonist.
- 5. Neonatal period, increase in 1st 2-3 days of life due to post-natal surge

DECREASED TSH LEVELS:

- 1.Toxic multi-nodular goitre & Thyroiditis.
- 2. Over replacement of thyroid harmone in treatment of hypothyroidism.
- 3. Autonomously functioning Thyroid adenoma
- 4. Secondary pituatary or hypothalmic hypothyroidism
- 5. Acute psychiatric illness
- 6. Severe dehydration.
- 7.DRUGS: Glucocorticoids, Dopamine, Levodopa, T4 replacement therapy, Anti-thyroid drugs for thyrotoxicosis.

8. Pregnancy: 1st and 2nd Trimester

DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)





A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

REPORTING DATE

ng/mL

NAME : Mr. SHANTANU

AGE/ GENDER : 30 YRS/MALE **PATIENT ID** : 1612778

COLLECTED BY : 122409140028 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 14/Sep/2024 11:37 AM BARCODE NO. : 12504705 **COLLECTION DATE** : 14/Sep/2024 11:40AM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

Test Name Value Unit **Biological Reference interval**

VITAMINS

VITAMIN D/25 HYDROXY VITAMIN D3

VITAMIN D (25-HYDROXY VITAMIN D3): SERUM by CLIA (CHEMILUMINESCENCE IMMUNOASSAY)

26.2L

DEFICIENCY: < 20.0

: 14/Sep/2024 06:59PM

INSUFFICIENCY: 20.0 - 30.0 SUFFICIENCY: 30.0 - 100.0

TOXICITY: > 100.0

INTERPRETATION:

CLIENT CODE.

DEFICIENT:	< 20	ng/mL
	21 22	<i>J</i> , ,
INSUFFICIENT:	21 - 29	ng/mL
PREFFERED RANGE:	30 - 100	ng/mL
INTOXICATION:	> 100	ng/mL

1. Vitamin D compounds are derived from dietary ergocalciferol (from plants, Vitamin D2), or cholecalciferol (from animals, Vitamin D3), or by conversion of 7- dihydrocholecalciferol to Vitamin D3 in the skin upon Ultraviolet exposure.

2.25-OH--Vitamin D represents the main body resevoir and transport form of Vitamin D and transport form of Vitamin D, being stored in adipose tissue and tightly bound by a transport protein while in circulation.

3. Vitamin D plays a primary role in the maintenance of calcium homeostatis. It promotes calcium absorption, renal calcium absorption and phosphate reabsorption, skeletal calcium deposition, calcium mobilization, mainly regulated by parathyroid harmone (PTH).

4. Severe deficiency may lead to failure to mineralize newly formed osteoid in bone, resulting in rickets in children and osteomalacia in adults.

DECREASED:

- 1.Lack of sunshine exposure
- 2.Inadequate intake, malabsorption (celiac disease)
 3.Depressed Hepatic Vitamin D 25- hydroxylase activity
- 4. Secondary to advanced Liver disease
- 5. Osteoporosis and Secondary Hyperparathroidism (Mild to Moderate deficiency)
- 6.Enzyme Inducing drugs: anti-epileptic drugs like phenytoin, phenobarbital and carbamazepine, that increases Vitamin D metabolism.

1. Hypervitaminosis D is Rare, and is seen only after prolonged exposure to extremely high doses of Vitamin D. When it occurs, it can result in severe hypercalcemia and hyperphophatemia.

CAUTION: Replacement therapy in deficient individuals must be monitored by periodic assessment of Vitamin D levels in order to prevent hypervitaminosis D

NOTE:-Dark coloured individuals as compare to whites, is at higher risk of developing Vitamin D deficiency due to excess of melanin pigment which interefere with Vitamin D absorption.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)





A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

NAME : Mr. SHANTANU

AGE/ GENDER : 30 YRS/MALE **PATIENT ID** : 1612778

COLLECTED BY : 122409140028 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 14/Sep/2024 11:37 AM BARCODE NO. : 12504705 **COLLECTION DATE** : 14/Sep/2024 11:40AM

CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 14/Sep/2024 03:54PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name Value Unit **Biological Reference interval**

VITAMIN B12/COBALAMIN

VITAMIN B12/COBALAMIN: SERUM 236.1 pg/mL 200.0 - 1100.0

by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

INTERPRETATION:-

INCREASED VITAMIN B12	DECREASED VITAMIN B12		
1.Ingestion of Vitamin C	1.Pregnancy		
2.Ingestion of Estrogen	2.DRUGS:Aspirin, Anti-convulsants, Colchicine		
3.Ingestion of Vitamin A	3.Ethanol Igestion		
4.Hepatocellular injury	4. Contraceptive Harmones		
5.Myeloproliferative disorder	5.Haemodialysis 5.Haemodialysi 5.Haemodialysis 5.Haemodialysis 5.Haemodialysis 5.Haemodialysis		
6.Uremia	6. Multiple Myeloma		

- 1. Vitamin B12 (cobalamin) is necessary for hematopoiesis and normal neuronal function.
- 2.In humans, it is obtained only from animal proteins and requires intrinsic factor (IF) for absorption.
- 3. The body uses its vitamin B12 stores very economically, reabsorbing vitamin B12 from the ileum and returning it to the liver; very little is excreted.
- 4.Vitamin B12 deficiency may be due to lack of IF secretion by gastric mucosa (eg, gastrectomy, gastric atrophy) or intestinal malabsorption (eg, ileal resection, small intestinal diseases).
- 5. Vitamin B12 deficiency frequently causes macrocytic anemia, glossitis, peripheral neuropathy, weakness, hyperreflexia, ataxia, loss of proprioception, poor coordination, and affective behavioral changes. These manifestations may occur in any combination; many patients have the neurologic defects without macrocytic anemia.
- 6.Serum methylmalonic acid and homocysteine levels are also elevated in vitamin B12 deficiency states.
- 7. Follow-up testing for antibodies to intrinsic factor (IF) is recommended to identify this potential cause of vitamin B12 malabsorption. NOTE:A normal serum concentration of vitamin B12 does not rule out tissue deficiency of vitamin B12. The most sensitive test for vitamin B12 deficiency at the cellular level is the assay for MMA. If clinical symptoms suggest deficiency, measurement of MMA and homocysteine should be considered, even if serum vitamin B12 concentrations are normal.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)





PKR JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

NAME : Mr. SHANTANU

AGE/ GENDER : 30 YRS/MALE **PATIENT ID** : 1612778

COLLECTED BY : 122409140028 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 14/Sep/2024 11:37 AM BARCODE NO. : 12504705 **COLLECTION DATE** : 14/Sep/2024 11:40AM

CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 14/Sep/2024 03:54PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Value Unit **Biological Reference interval** Test Name

CLINICAL PATHOLOGY URINE ROUTINE & MICROSCOPIC EXAMINATION

PHYSICAL EXAMINATION

QUANTITY RECIEVED ml by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

PALE YELLOW PALE YELLOW **COLOUR**

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY **TRANSPARANCY CLEAR CLEAR**

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY 1.02 1.002 - 1.030 SPECIFIC GRAVITY

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY.

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

CHEMICAL EXAMINATION

REACTION **ACIDIC**

PROTEIN NEGATIVE (-ve) NEGATIVE (-ve)

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY **SUGAR NEGATIVE** (-ve) **NEGATIVE** (-ve)

рΗ 5.5 5.0 - 7.5

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY **BILIRUBIN NEGATIVE** (-ve) **NEGATIVE** (-ve)

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

NITRITE NEGATIVE (-ve) **NEGATIVE** (-ve)

EU/dL **NOT DETECTED** UROBILINOGEN 0.2 - 1.0

KETONE BODIES **NEGATIVE (-ve) NEGATIVE (-ve)**

NEGATIVE (-ve) NEGATIVE (-ve) BLOOD

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

NEGATIVE (-ve) ASCORBIC ACID **NEGATIVE** (-ve) by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

MICROSCOPIC EXAMINATION



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)





A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

NAME : Mr. SHANTANU

AGE/ GENDER : 30 YRS/MALE **PATIENT ID** : 1612778

COLLECTED BY : 122409140028 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 14/Sep/2024 11:37 AM BARCODE NO. : 12504705 **COLLECTION DATE** : 14/Sep/2024 11:40AM

CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 14/Sep/2024 03:54PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name	Value	Unit	Biological Reference interval
RED BLOOD CELLS (RBCs) by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	NEGATIVE (-ve)	/HPF	0 - 3
PUS CELLS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	3-5	/HPF	0 - 5
EPITHELIAL CELLS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	2-4	/HPF	ABSENT
CRYSTALS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	NEGATIVE (-ve)		NEGATIVE (-ve)
CASTS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	NEGATIVE (-ve)		NEGATIVE (-ve)
BACTERIA by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	NEGATIVE (-ve)		NEGATIVE (-ve)
OTHERS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	NEGATIVE (-ve)		NEGATIVE (-ve)
TRICHOMONAS VAGINALIS (PROTOZOA) by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	ABSENT		ABSENT

*** End Of Report



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS, MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST

