TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT.

NAME	: Mr. SHUBHAM		
AGE/ GENDER	: 23 YRS/MALE	**PATIENT ID**	: 1642828
COLLECTED BY	:	**REG. NO./LAB NO.**	: **122410140021**
REFERRED BY	:	**REGISTRATION DATE**	: 14/Oct/2024 02:10 PM
BARCODE NO.	: 12505174	**COLLECTION DATE**	: 14/Oct/2024 03:42PM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	**REPORTING DATE**	: 14/Oct/2024 04:07PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA		

Test Name	Value	Unit	Biological Reference interval

HAEMATOLOGY
COMPLETE BLOOD COUNT (CBC)

RED BLOOD CELLS (RBCS) COUNT AND INDICES

Test Name	Value	Unit	Biological Reference interval
HAEMOGLOBIN (HB) *by CALORIMETRIC*	14.4	gm/dL	12.0 - 17.0
RED BLOOD CELL (RBC) COUNT *by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE*	4.95	Millions/cmm	3.50 - 5.00
PACKED CELL VOLUME (PCV) *by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER*	44.1	%	40.0 - 54.0
MEAN CORPUSCULAR VOLUME (MCV) *by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER*	89.2	fL	80.0 - 100.0
MEAN CORPUSCULAR HAEMOGLOBIN (MCH) *by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER*	29.1	pg	27.0 - 34.0
MEAN CORPUSCULAR HEMOGLOBIN CONC. (MCHC) *by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER*	32.6	g/dL	32.0 - 36.0
RED CELL DISTRIBUTION WIDTH (RDW-CV) *by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER*	13.7	%	11.00 - 16.00
RED CELL DISTRIBUTION WIDTH (RDW-SD) *by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER*	45.6	fL	35.0 - 56.0
MENTZERS INDEX *by CALCULATED*	18.02	RATIO	BETA THALASSEMIA TRAIT: < 13.0 IRON DEFICIENCY ANEMIA: >13.0
GREEN & KING INDEX *by CALCULATED*	24.7	RATIO	BETA THALASSEMIA TRAIT:<= 65.0 IRON DEFICIENCY ANEMIA: > 65.0

WHITE BLOOD CELLS (WBCS)

Test Name	Value	Unit	Biological Reference interval
TOTAL LEUCOCYTE COUNT (TLC) *by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY*	7810	/cmm	4000 - 11000
NUCLEATED RED BLOOD CELLS (nRBCS) *by AUTOMATED 6 PART HEMATOLOGY ANALYZER*	NIL		0.00 - 20.00
NUCLEATED RED BLOOD CELLS (nRBCS) % *by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER*	NIL	%	< 10 %

DIFFERENTIAL LEUCOCYTE COUNT (DLC)

Test Name	Value	Unit	Biological Reference interval
NEUTROPHILS *by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY*	74[H]	%	50 - 70

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)

NAME	: Mr. SHUBHAM		
AGE/ GENDER	: 23 YRS/MALE	**PATIENT ID**	: 1642828
COLLECTED BY	:	**REG. NO./LAB NO.**	**: 122410140021**
REFERRED BY	:	**REGISTRATION DATE**	: 14/Oct/2024 02:10 PM
BARCODE NO.	: 12505174	**COLLECTION DATE**	: 14/Oct/2024 03:42PM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	**REPORTING DATE**	: 14/Oct/2024 04:07PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA		

Test Name	Value	Unit	Biological Reference interval
LYMPHOCYTES	16^L	%	20 - 40
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY			
EOSINOPHILS	3	%	1 - 6
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY			
MONOCYTES	7	%	2 - 12
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY			
BASOPHILS	0	%	0 - 1
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY			
ABSOLUTE LEUKOCYTES (WBC) COUNT			
ABSOLUTE NEUTROPHIL COUNT	5779	/cmm	2000 - 7500
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY			
ABSOLUTE LYMPHOCYTE COUNT	1250	/cmm	800 - 4900
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY			
ABSOLUTE EOSINOPHIL COUNT	234	/cmm	40 - 440
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY			
ABSOLUTE MONOCYTE COUNT	547	/cmm	80 - 880
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY			
ABSOLUTE BASOPHIL COUNT	0	/cmm	0 - 110
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY			
PLATELETS AND OTHER PLATELET PREDICTIVE MARKERS.			
PLATELET COUNT (PLT)	180000	/cmm	150000 - 450000
by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE			
PLATELETCRIT (PCT)	0.22	%	0.10 - 0.36
by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE			
MEAN PLATELET VOLUME (MPV)	12^H	fL	6.50 - 12.0
by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE			
PLATELET LARGE CELL COUNT (P-LCC)	75000	/cmm	30000 - 90000
by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE			
PLATELET LARGE CELL RATIO (P-LCR)	41.6	%	11.0 - 45.0
by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE			
PLATELET DISTRIBUTION WIDTH (PDW)	16.1	%	15.0 - 17.0
by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE			

NOTE: TEST CONDUCTED ON EDTA WHOLE BLOOD

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)

NOT VALID FOR MEDICO LEGAL PURPOSE

440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600
REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)

Page 2 of 5

NAME	: Mr. SHUBHAM		
AGE/ GENDER	: 23 YRS/MALE	**PATIENT ID**	: 1642828
COLLECTED BY	:	**REG. NO./LAB NO.**	: **122410140021**
REFERRED BY	:	**REGISTRATION DATE**	: 14/Oct/2024 02:10 PM
BARCODE NO.	: 12505174	**COLLECTION DATE**	: 14/Oct/2024 03:42PM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	**REPORTING DATE**	: 14/Oct/2024 04:50PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA		

Test Name	Value	Unit	Biological Reference interval

IMMUNOPATHOLOGY/SEROLOGY

TYPHOID COMBO SCREEN (TYPHOID ANTIGEN, IgG AND IgM): SERUM

TYPHOID ANTIGEN - SERUM	NEGATIVE (-ve)		NEGATIVE (-ve)
by ICT (IMMUNOCHROMATOGRAPHY)			
TYPHI DOT ANTIBODY IgG	NEGATIVE (-ve)		NEGATIVE (-ve)
by ICT (IMMUNOCHROMATOGRAPHY)			
TYPHI DOT ANTIBODY IgM	NEGATIVE (-ve)		NEGATIVE (-ve)
by ICT (IMMUNOCHROMATOGRAPHY)			

INTE*RPRETATION:*

Typhoid fever is a life threatening illness caused by the bacterium Salmonella typhus.The infection is acquired typically by ingestion. On reaching the gut, the bacilli attach themselves to the epithelial cells of the intestinal villi and penetrate the lamina and submucosa. They are then phagocytosed there by polymorphs and mesenteric lymph nodes, where they multiply and, via the thoracic duct, enter the blood stream. A transient bacteremia follows, during which the bacilli are seeded in the liver, gall bladder, spleen, bone marrow, lymph nodes, and kidneys, where further multiplication takes place. Towards the end of the incubation period, there occurs a massive bacteremia from these sites, heralding the onset of the clinical symptoms.

The diagnosis of typhoid consists of isolation of the bacilli and the demonstration of antibodies. The isolation of the bacilli is very time consuming and antibody detection is not very specific. Other tests include the Widal reaction. The advantage of this test is that it takes only 10-20 minutes and requires only a small amount of stool/serum/plasma to perform. It is the easiest and most specific method for detecting S. typhi infection.

RELATIVE SENSTIVITY OF TYPHOID ANTIGEN DETECTION: 98.7%
RELATIVE SPECIFICITY OF TYPHOID ANTIGEN DETECTION: 97.4%

DETECTABLE IgM RESPONSE:

ONSET OF FEVER	PERCENT POSITIVE
4 - 6 DAYS	43.5
6 - 9 DAYS	92.9
> 9 DAYS	99.5

1.This is a solid phase, immunochromatographic ELISA assay that detects specific IgM and IgG Antibodies against the OUTER MEMBRAN PROTEIN(OMP) of the Salmonella species. IgM antibodies appear in the serum 2-3 days post infection and are indicative of a recent infection while the IgG antibodies appear later and are useful for presumptive diagnosis of Enteric fever if the patient presents more than a week after onset of symptoms.

2.This is a useful screening assay for the early detection of Enteric fever and has a high sensitivity. However the test has moderate specificity and false positive results may be obtained in the following situations:

· Antibodies against Salmonella may cross react with other antibodies.

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)

NOT VALID FOR MEDICO LEGAL PURPOSE

440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600
REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)

Page 3 of 5

NAME	: Mr. SHUBHAM		
AGE/ GENDER	: 23 YRS/MALE	PATIENT ID	: 1642828
COLLECTED BY	:	REG. NO./LAB NO.	: 122410140021
REFERRED BY	:	REGISTRATION DATE	: 14/Oct/2024 02:10 PM
BARCODE NO.	: 12505174	COLLECTION DATE	: 14/Oct/2024 03:42PM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	REPORTING DATE	: 14/Oct/2024 04:50PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA		

Test Name	Value	Unit	Biological Reference interval

· Unrelated infections may lead to production of specific Salmonella antibodies if the patient has previously been exposed to Salmonella infection **(ANAMNESTIC RESPONSE)**.

NOTE:-Rapid blood culture performed during 1^{st} week of infection is highly recommended for confirmation of all IgM positive results. In case the patient has presented after the first week of infection, a thorough clinical correlation and confirmatory Widal test must be performed to establish the diagnosis.

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)

NOT VALID FOR MEDICO LEGAL PURPOSE

440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600
REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)

Page 4 of 5

NAME	: Mr. SHUBHAM		
AGE/ GENDER	: 23 YRS/MALE	**PATIENT ID**	: 1642828
COLLECTED BY	:	**REG. NO./LAB NO.**	: **122410140021**
REFERRED BY	:	**REGISTRATION DATE**	: 14/Oct/2024 02:10 PM
BARCODE NO.	: 12505174	**COLLECTION DATE**	: 14/Oct/2024 03:42PM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	**REPORTING DATE**	: 15/Oct/2024 06:52AM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA		

Test Name	Value	Unit	Biological Reference interval

DENGUE FEVER ANTIGEN NS1 - ELISA (QUANTITATIVE)

Test Name	Value	Unit	Biological Reference interval
DENGUE NS1 ANTIGEN QUANTITATIVE *by ELISA (ENZYME LINKED IMMUNOSORBENT ASSAY)*	0.339	INDEX	NEGATIVE: < 0.90 BORDERLINE: 0.90 - 1.10 POSITIVE: >=1.10
DENGUE NS1 ANTIGEN RESULT *by ELISA (ENZYME LINKED IMMUNOSORBENT ASSAY)*	NEGATIVE (-ve)		NEGATIVE (-ve)

INTERPRETATION

DENGUE ANTIGEN NS1		
VALUE	**UNIT**	**RESULT**
< 0.90	INDEX	NEGATIVE (-ve)
0.90 - 1.10	INDEX	BORDERLINE
>=1.10	INDEX	POSITIVE (+ve)

1.The test becomes positive within 0-9 days of exposure to the virus (positive results are obtained within 24 hours of exposure in the overwhelming majority of patients) and generally remains positive till 15 days after exposure. The Dengue NS-1 antigen test is extremely useful in the early diagnosis of the disease thus helping in proper follow up and monitoring of the patients.

2.The IgM antibodies on the other hand take a minimum of 5-10 days in primary infection and 4-5 days in secondary infections to test positive and hence are suitable for the diagnosis of dengue fever only when the fever is approximately one week old.

*** End Of Report ***

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)

TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT.

NOT VALID FOR MEDICO LEGAL PURPOSE

440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600
REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)

Page 5 of 5