



# P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

**A PIONEER DIAGNOSTIC CENTRE**

☎ 0171-2532620, 8222896961 ✉ pkrjainhealthcare@gmail.com

**NAME** : Mr. CHARANJIT PRASHAR  
**AGE/ GENDER** : 53 YRS/MALE  
**COLLECTED BY** :  
**REFERRED BY** :  
**BARCODE NO.** : 12505348  
**CLIENT CODE.** : P.K.R JAIN HEALTHCARE INSTITUTE  
**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**PATIENT ID** : 1593968  
**REG. NO./LAB NO.** : 122410250015  
**REGISTRATION DATE** : 25/Oct/2024 01:13 PM  
**COLLECTION DATE** : 25/Oct/2024 01:14PM  
**REPORTING DATE** : 25/Oct/2024 03:14PM

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

## CLINICAL CHEMISTRY/BIOCHEMISTRY

### KIDNEY FUNCTION TEST (BASIC)

UREA: SERUM <i>by UREASE - GLUTAMATE DEHYDROGENASE (GLDH)</i>	24.83	mg/dL	10.00 - 50.00
CREATININE: SERUM <i>by ENZYMATIC, SPECTROPHOTOMETRY</i>	0.91	mg/dL	0.40 - 1.40
BLOOD UREA NITROGEN (BUN): SERUM <i>by CALCULATED, SPECTROPHOTOMETRY</i>	11.6	mg/dL	7.0 - 25.0
BLOOD UREA NITROGEN (BUN)/CREATININE RATIO: SERUM <i>by CALCULATED, SPECTROPHOTOMETRY</i>	12.75	RATIO	10.0 - 20.0
UREA/CREATININE RATIO: SERUM <i>by CALCULATED, SPECTROPHOTOMETRY</i>	27.29	RATIO	
URIC ACID: SERUM <i>by URICASE - OXIDASE PEROXIDASE</i>	6.74	mg/dL	3.60 - 7.70



DR.VINAY CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY)





# P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

**A PIONEER DIAGNOSTIC CENTRE**

☎ 0171-2532620, 8222896961 ✉ pkrjainhealthcare@gmail.com

NAME	: Mr. CHARANJIT PRASHAR	PATIENT ID	: 1593968
AGE/ GENDER	: 53 YRS/MALE	REG. NO./LAB NO.	: 122410250015
COLLECTED BY	:	REGISTRATION DATE	: 25/Oct/2024 01:13 PM
REFERRED BY	:	COLLECTION DATE	: 25/Oct/2024 01:14PM
BARCODE NO.	: 12505348	REPORTING DATE	: 25/Oct/2024 03:14PM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE		
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

#### INTERPRETATION:

Normal range for a healthy person on normal diet: 12 - 20

To Differentiate between pre- and postrenal azotemia.

#### **INCREASED RATIO (>20:1) WITH NORMAL CREATININE:**

- 1.Prerenal azotemia (BUN rises without increase in creatinine) e.g. heart failure, salt depletion,dehydration, blood loss) due to decreased glomerular filtration rate.
- 2.Catabolic states with increased tissue breakdown.
- 3.GI hemorrhage.
- 4.High protein intake.
- 5.Impaired renal function plus .
- 6.Excess protein intake or production or tissue breakdown (e.g. infection, GI bleeding, thyrotoxicosis, Cushings syndrome, high protein diet, burns,surgery, cachexia, high fever).
- 7.Urine reabsorption (e.g. ureterocolostomy)
- 8.Reduced muscle mass (subnormal creatinine production)
- 9.Certain drugs (e.g. tetracycline, glucocorticoids)

#### **INCREASED RATIO (>20:1) WITH ELEVATED CREATININE LEVELS:**

- 1.Postrenal azotemia (BUN rises disproportionately more than creatinine) (e.g. obstructive uropathy).
- 2.Prerenal azotemia superimposed on renal disease.

#### **DECREASED RATIO (<10:1) WITH DECREASED BUN :**

- 1.Acute tubular necrosis.
- 2.Low protein diet and starvation.
- 3.Severe liver disease.
- 4.Other causes of decreased urea synthesis.
- 5.Repeated dialysis (urea rather than creatinine diffuses out of extracellular fluid).
- 6.Inherited hyperammonemias (urea is virtually absent in blood).
- 7.SIADH (syndrome of inappropriate antidiuretic hormone) due to tubular secretion of urea.
- 8.Pregnancy.

#### **DECREASED RATIO (<10:1) WITH INCREASED CREATININE:**

- 1.Phenacimide therapy (accelerates conversion of creatine to creatinine).
- 2.Rhabdomyolysis (releases muscle creatinine).
- 3.Muscular patients who develop renal failure.

#### **INAPPROPRIATE RATIO:**

- 1.Diabetic ketoacidosis (acetoacetate causes false increase in creatinine with certain methodologies,resulting in normal ratio when dehydration should produce an increased BUN/creatinine ratio).
- 2.Cephalosporin therapy (interferes with creatinine measurement).

\*\*\* End Of Report \*\*\*



DR.VINAY CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS , MD (PATHOLOGY)

