PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana) A PIONEER DIAGNOSTIC CENTRE

【 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. MANILA VAID			
AGE/ GENDER	: 45 YRS/FEMALE		PATIENT ID	: 1677980
COLLECTED BY	:		REG. NO./LAB NO.	: 122411210013
REFERRED BY	:		REGISTRATION DATE	: 21/Nov/2024 11:36 AM
BARCODE NO.	: 12505766		COLLECTION DATE	: 21/Nov/2024 11:38AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITU	ΤЕ	REPORTING DATE	: 21/Nov/2024 01:30PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBAL	A CITY - H	ARYANA	
Test Name		Value	Unit	Biological Reference interval
	SWASTI	HYA WI	ELLNESS PANEL: 1.2	2
	СОМР	LETE BI	LOOD COUNT (CBC)	
RED BLOOD CELLS	(RBCS) COUNT AND INDICES			
HAEMOGLOBIN (HI	3)	11.2 ^L	gm/dL	12.0 - 16.0
RED BLOOD CELL () by HYDRO DYNAMIC F	RBC) COUNT OCUSING, ELECTRICAL IMPEDENCE	4.35	Millions/	cmm 3.50 - 5.00
PACKED CELL VOLU	JME (PCV) utomated hematology analyzer	32.4 ^L	%	37.0 - 50.0
MEAN CORPUSCULA	AR VOLUME (MCV) utomated hematology analyzer	74.4 ^L	KR fl	80.0 - 100.0
	AR HAEMOGLOBIN (MCH) utomated hematology analyzer	25.7 ^L	pg	27.0 - 34.0
	AR HEMOGLOBIN CONC. (MCHC) UTOMATED HEMATOLOGY ANALYZER	34.5	g/dL	32.0 - 36.0
	JTION WIDTH (RDW-CV) UTOMATED HEMATOLOGY ANALYZER	14.3	%	11.00 - 16.00
	JTION WIDTH (RDW-SD) utomated hematology analyzer	40.8	fL	35.0 - 56.0
MENTZERS INDEX by CALCULATED		17.1	RATIO	BETA THALASSEMIA TRAIT: < 13.0 IRON DEFICIENCY ANEMIA:
GREEN & KING IND by calculated	EX	24.41	RATIO	>13.0 BETA THALASSEMIA TRAIT:<= 65.0 IRON DEFICIENCY ANEMIA: > 65.0
WHITE BLOOD CEI	LLS (WBCS)			
TOTAL LEUCOCYTE	COUNT (TLC) by sf cube & microscopy	8220	/cmm	4000 - 11000
DIFFERENTIAL LE	<u>UCOCYTE COUNT (DLC)</u>			
NEUTROPHILS	BY SF CUBE & MICROSCOPY	60	%	50 - 70
by FLOW CYTOMETRY				

DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS , MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST

NOT VALID FOR MEDICO LEGAL PURPOSE



PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana) A PIONEER DIAGNOSTIC CENTRE

🔽 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. MANILA VAID			
AGE/ GENDER	: 45 YRS/FEMALE	P	ATIENT ID	: 1677980
COLLECTED BY	:	R	EG. NO./LAB NO.	: 122411210013
REFERRED BY	:	R	EGISTRATION DATE	: 21/Nov/2024 11:36 AM
BARCODE NO.	: 12505766	C	OLLECTION DATE	: 21/Nov/2024 11:38AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTIT	TUTE R	EPORTING DATE	: 21/Nov/2024 01:30PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBA	ALA CITY - HAR	YANA	
Test Name		Value	Unit	Biological Reference interval
	Y BY SF CUBE & MICROSCOPY			
EOSINOPHILS	Y BY SF CUBE & MICROSCOPY	3	%	1 - 6
MONOCYTES		6	%	2 - 12
	Y BY SF CUBE & MICROSCOPY			
BASOPHILS	Y BY SF CUBE & MICROSCOPY	0	%	0 - 1
	CYTES (WBC) COUNT			
ABSOLUTE NEUTR	OPHIL COUNT y by sf cube & microscopy	4932	/cmm	2000 - 7500
ABSOLUTE LYMPH		2548 ^L	/cmm	800 - 4900
ABSOLUTE EOSINO		247	/cmm	40 - 440
ABSOLUTE MONOC by FLOW CYTOMETR	CYTE COUNT Y by sf cube & microscopy	493	/cmm	80 - 880
ABSOLUTE BASOP	HIL COUNT y by sf cube & microscopy	0	/cmm	0 - 110
PLATELETS AND (DTHER PLATELET PREDICTIVE	MARKERS.		
PLATELET COUNT by hydro dynamic f	(PLT) FOCUSING, ELECTRICAL IMPEDENCE	367000	/cmm	150000 - 450000
PLATELETCRIT (PC by HYDRO DYNAMIC F	CT) FOCUSING, ELECTRICAL IMPEDENCE	0.32	%	0.10 - 0.36

9

70000

19.2

15.5



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS , MD (PATHOLOGY)

fL

%

%

/cmm

6.50 - 12.0

11.0 - 45.0

15.0 - 17.0

30000 - 90000

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST

MEAN PLATELET VOLUME (MPV)

PLATELET LARGE CELL COUNT (P-LCC)

PLATELET LARGE CELL RATIO (P-LCR)

PLATELET DISTRIBUTION WIDTH (PDW)

by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE

by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE

by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE

by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE NOTE: TEST CONDUCTED ON EDTA WHOLE BLOOD





A PIONEER DIAGNOSTIC CENTRE

【 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

AGE/ GENDER : 45 YRS/FEMALE PATIENT ID : 167798∪ COLLECTED BY : REG. NO./LAB NO. : 122411210013 REFERRED BY : REGISTRATION DATE : 21/Nov/2024 11:36 AM BARCODE NO. : 12505766 COLLECTION DATE : 21/Nov/2024 11:38AM CLIENT CODE : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 21/Nov/2024 01:30PM CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA : image: state st	
REFERRED BY :: REGISTRATION DATE : 21/Nov/2024 11:36 AM BARCODE NO. : 12505766 COLLECTION DATE : 21/Nov/2024 11:38AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 21/Nov/2024 01:30PM CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA Test Name Value Unit Biological Reference in ERYTHROCYTE SEDIMENTATION RATE (ESR) 40 ^H mm/1st hr 0 - 20 by RED CELL AGGREGATION BY CAPILLARY PHOTOMETRY 40 ^H mm/1st hr 0 - 20 VNERPRETATION: 1. ESR is a non-specific test because an elevated result often indicates the presence of inflammation associated with infection, cancer a immune disease, but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it. . . . 2. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other as C-reactive protein . . 3. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as some others, systemic Lupus erythematosus . . CONDITION WITH LO	
BARCODE NO. : 12505766 COLLECTION DATE : 21/Nov/2024 11:38AM CLIENT CODE. : P.K.R. JAIN HEALTHCARE INSTITUTE REPORTING DATE : 21/Nov/2024 01:30PM CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA : 21/Nov/2024 01:30PM Test Name Value Unit Biological Reference in ERYTHROCYTE SEDIMENTATION RATE (ESR) 40 ^H mm/1st hr 0 - 20 by RED CELL AGGREGATION BY CAPILLARY PHOTOMETRY MM 1 - 20 INTERPRETATION: 1. SSR is a non-specific test because an elevated result often indicates the presence of inflammation associated with infection, cancer a immune disease, but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it. 2. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other as C-reactive protein 3. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as some others, systemic lupus erythematosus CONDITION WITH LOW ESR CONDITION WITH LOW ESR 1 1 A low ESR can be seen with conditions that inhibit the normal sedimentation of red blood cells, such as a high red blood cell count (polycythaemia), significantly high white blood cell count (leucocytosis), and some protein abnormalities. Some changes in red cell sh as sickle cells in sickle cell anaemia) also lower the ESR.	
CLIENT CODE : P.K.R. JAIN HEALTHCARE INSTITUTE REPORTING DATE : 21/Nov/2024 01:30PM CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA Biological Reference in Test Name Value Unit Biological Reference in ERYTHROCYTE SEDIMENTATION RATE (ESR) 40^H mm/1st hr 0 - 20 <i>by RED CELL AGGREGATION BY CAPILLARY PHOTOMETRY</i> 40^H mm/1st hr 0 - 20 INTERPRETATION 1. SER is a non-specific test because an elevated result often indicates the presence of inflammation associated with infection, cancer a immune disease, but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it. 2. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other as C-reactive protein 3. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as some others, systemic lupus erythematosus Some changes in red cell some since cell some changes in red cell some sickle cell anaemia) also lower the ESR. MOTION WITH LOW ESR . . . 1. Sens and C - reactive protein (C-RP) are both markers of inflammation. . . 2. Generally, ESR does not change as rapidly as does CRP, either at the start of inflammation or as it resolves. . 3. CRP is not affected by as	
CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA Test Name Value Unit Biological Reference in ERYTHROCYTE SEDIMENTATION RATE (ESR) 40 ^H mm/1st hr 0 - 20 by ReD CELL AGGREGATION BY CAPILLARY PHOTOMETRY 40 ^H mm/1st hr 0 - 20 INTERPRETATION: 1.5SR is a non-specific test because an elevated result often indicates the presence of inflammation associated with infection, cancer a immune disease, but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it. 2. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other as C-reactive protein 3. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as some others, systemic lupus erythematosus CONDITION WITH LOW ESR A low ESR can be seen with conditions that inhibit the normal sedimentation of red blood cells, such as a high red blood cell count (polycythaemia), significantly high white blood cell count (leucocytosis), and some protein abnormalities. Some changes in red cell sh as sickle cells in sickle cell anaemia) also lower the ESR. NOTE: 1. SR is elevated, it is typically a result of two types of proteins, globulins or finflammation. 2. Generally, ESR does not change as rapidly as does CRP, either at the start of inflammation or as it resolves. 3. CRP is not affected by as many other factors as is ESR, makingit a b	
Test Name Value Unit Biological Reference in ERYTHROCYTE SEDIMENTATION RATE (ESR) by RED CELL AGGREGATION BY CAPILLARY PHOTOMETRY by RED CELL AGGREGATION BY CAPILLARY PHOTOMETRY INTERPRETATION: 1. ESR is a non-specific test because an elevated result often indicates the presence of inflammation associated with infection, cancer a immune disease, but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it. 2. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other as C-reactive protein 3. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as some others, systemic lupus erythematosus CONDITION WITH LOW ESR A low ESR can be seen with conditions that inhibit the normal sedimentation of red blood cells, such as a high red blood cell count (polycythaemia), significantly high white blood cell count (leucocytosis), and some protein abnormalities. Some changes in red cell sh as sickle cells in sickle cells on the ESR. NOTE: 1. ESR is devated, it is typically a result of two types of proteins, globulins or fibrinogen. 2. GRPS in the devated, it is typically a result of two types of proteins, globulins or fibrinogen. 3. Wome tend to have a higher ESR.	
ERYTHROCYTE SEDIMENTATION RATE (ESR) ERYTHROCYTE SEDIMENTATION RATE (ESR) by RED CELL AGGREGATION BY CAPILLARY PHOTOMETRY INTERPRETATION: 1. ESR is a non-specific test because an elevated result often indicates the presence of inflammation associated with infection, cancer a immune disease, but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it. 2. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other as C-reactive protein 3. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as some others, systemic lupus erythematosus CONDITION WITH LOW ESR A low ESR can be seen with conditions that inhibit the normal sedimentation of red blood cells, such as a high red blood cell count (polycythaemia), significantly high white blood cell count (leucocytosis), and some protein abnormalities. Some changes in red cell sh as sickle cells in sickle cell anaemia) also lower the ESR. NOTE: 1. ESR is a dee not change as rapidly as does CRP, either at the start of inflammation or as it resolves. 3. CPP is not affected by as many other factors as is ESR, making it a better marker of inflammation. Comparison of the above diseases as well as some changes in red cell sh as sickle cells in sickle cell anaemia) also lower the ESR.	
ERYTHROCYTE SEDIMENTATION RATE (ESR) 40 ^H mm/1st hr 0 - 20 by RED CELL AGGREGATION BY CAPILLARY PHOTOMETRY 0.4 ^H mm/1st hr 0 - 20 INTERPRETATION: 1. ESR is a non-specific test because an elevated result often indicates the presence of inflammation associated with infection, cancer a immune disease, but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it. 2. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other as C-reactive protein 3. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as some others, systemic lupus erythematosus CONDITION WITH LOW ESR A low ESR can be seen with conditions that inhibit the normal sedimentation of red blood cells, such as a high red blood cell count (polycythaemia), significantly high white blood cell count (leucocytosis), and some protein abnormalities. Some changes in red cell sh as sickle cells in sickle cell anaemia) also lower the ESR. NOTE: 1. ESR and C - reactive protein (C-RP) are both markers of inflammation. 2. Generally, ESR does not change as rapidly as does CRP, either at the start of inflammation or as it resolves. 3. Or Pis not affected by as many other factors as is ESR, making it a better marker of inflammation. 4. If the ESR is elevated, it is typically a result of two types of proteins, globulins or fibrinogen. 5. Women tend to have a higher ESR. and menstruation and pregnancy can caus	terval
ERYTHROCYTE SEDIMENTATION RATE (ESR) 40 ^H mm/1st hr 0 - 20 <i>by RED CELL AGGREGATION BY CAPILLARY PHOTOMETRY</i> 0.40 ^H mm/1st hr 0 - 20 INTERPRETATION: 1. ESR is a non-specific test because an elevated result often indicates the presence of inflammation associated with infection, cancer a immune disease, but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it. 2. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other as C-reactive protein 3. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as some others, systemic lupus erythematosus CONDITION WITH LOW ESR A low ESR can be seen with conditions that inhibit the normal sedimentation of red blood cells, such as a high red blood cell count (polycythaemia), significantly high white blood cell count (leucocytosis) , and some protein abnormalities. Some changes in red cell sh as sickle cells in sickle cell anaemia) also lower the ESR. NOTE: 1. ESR and C - reactive protein (C-RP) are both markers of inflammation. 2. Generally, ESR does not change as rapidly as does CRP, either at the start of inflammation or as it resolves. 3. Grep is not affected by as many other factors as is ESR, making it a better marker of inflammation. 4. If the ESR is elevated, it is typically a result of two types of proteins, globulins or fibrinogen. 5. Wome tend to have a higher ESR. and menstruation and pregnancy can	
by RED CELL AGGREGATION BY CAPILLARY PHOTOMETRY INTERPRETATION: 1. ESR is a non-specific test because an elevated result often indicates the presence of inflammation associated with infection, cancer a mmune disease, but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it. 2. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other as C-reactive protein 3. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as some others, systemic lupus erythematosus CONDITION WITH LOW ESR A low ESR can be seen with conditions that inhibit the normal sedimentation of red blood cells, such as a high red blood cell count (polycythaemia), significantly high white blood cell count (leucocytosis), and some protein abnormalities. Some changes in red cell sh as sickle cells in sickle cell anaemia) also lower the ESR. NOTE: 1. ESR and C - reactive protein (C-RP) are both markers of inflammation. 2. Generally, ESR does not change as rapidly as does CRP, either at the start of inflammation or as it resolves. 3. CRP is not affected by as many other factors as is ESR, making it a better marker of inflammation. 4. If the ESR is elevated, it is typically a result of two types of proteins, globulins or fibrinogen. 5. Women tend to have a higher ESR, and menstruation and pregnancy can cause temporary elevations.	
 1. ESR is a non-specific test because an elevated result often indicates the presence of inflammation associated with infection, cancer a immune disease, but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it. 2. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other as C-reactive protein 3. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as some others, systemic lupus erythematosus CONDITION WITH LOW ESR A low ESR can be seen with conditions that inhibit the normal sedimentation of red blood cells, such as a high red blood cell count (polycythaemia), significantly high white blood cell count (leucocytosis), and some protein abnormalities. Some changes in red cell sh as sickle cells in sickle cell anaemia) also lower the ESR. NOTE: 1. ESR and C - reactive protein (C-RP) are both markers of inflammation. 2. Generally, ESR does not change as rapidly as does CRP, either at the start of inflammation or as it resolves. 3. CRP is not affected by as many other factors as is ESR, making it a better marker of inflammation. 4. If the ESR is elevated, it is typically a result of two types of proteins, globulins or fibrinogen. 5. Women tend to have a higher ESR. and menstruation and pregnancy can cause temporary elevations. 	
 immune disease, but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it. 2. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other as C-reactive protein 3. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as some others, systemic lupus erythematosus CONDITION WITH LOW ESR A low ESR can be seen with conditions that inhibit the normal sedimentation of red blood cells, such as a high red blood cell count (polycythaemia), significantly high white blood cell count (leucocytosis), and some protein abnormalities. Some changes in red cell sh as sickle cells in sickle cell anaemia) also lower the ESR. NOTE: 1. ESR and C - reactive protein (C-RP) are both markers of inflammation. 2. Generally, ESR does not change as rapidly as does CRP, either at the start of inflammation or as it resolves. 3. CRP is not affected by as many other factors as is ESR, making it a better marker of inflammation. 4. If the ESR is elevated, it is typically a result of two types of proteins, globulins or fibrinogen. 5. Women tend to have a higher ESR. and menstruation and pregnancy can cause temporary elevations. 	ind aut
as C-reactive protein 3. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as some others, systemic lupus erythematosus CONDITION WITH LOW ESR A low ESR can be seen with conditions that inhibit the normal sedimentation of red blood cells, such as a high red blood cell count (polycythaemia), significantly high white blood cell count (leucocytosis), and some protein abnormalities. Some changes in red cell sh as sickle cells in sickle cell anaemia) also lower the ESR. NOTE: 1. ESR and C - reactive protein (C-RP) are both markers of inflammation. 2. Generally, ESR does not change as rapidly as does CRP, either at the start of inflammation or as it resolves. 3. CRP is not affected by as many other factors as is ESR, making it a better marker of inflammation. 4. If the ESR is elevated, it is typically a result of two types of proteins, globulins or fibrinogen. 5. Women tend to have a higher ESR. and menstruation and pregnancy can cause temporary elevations.	
 This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as some others, systemic lupus erythematosus CONDITION WITH LOW ESR A low ESR can be seen with conditions that inhibit the normal sedimentation of red blood cells, such as a high red blood cell count (polycythaemia), significantly high white blood cell count (leucocytosis), and some protein abnormalities. Some changes in red cell sh as sickle cells in sickle cell anaemia) also lower the ESR. NOTE: Generally, ESR does not change as rapidly as does CRP, either at the start of inflammation or as it resolves. GRP is not affected by as many other factors as is ESR, making it a better marker of inflammation. If the ESR is elevated, it is typically a result of two types of proteins, globulins or fibrinogen. Women tend to have a higher ESR. and menstruation and pregnancy can cause temporary elevations. 	test su
CONDITION WITH LOW ESR A low ESR can be seen with conditions that inhibit the normal sedimentation of red blood cells, such as a high red blood cell count (polycythaemia), significantly high white blood cell count (leucocytosis), and some protein abnormalities. Some changes in red cell sh as sickle cells in sickle cell anaemia) also lower the ESR. NOTE: 1. ESR and C - reactive protein (C-RP) are both markers of inflammation. 2. Generally, ESR does not change as rapidly as does CRP, either at the start of inflammation or as it resolves. 3. CRP is not affected by as many other factors as is ESR, making it a better marker of inflammation. 4. If the ESR is elevated, it is typically a result of two types of proteins, globulins or fibrinogen. 5. Women tend to have a higher ESR, and menstruation and pregnancy can cause temporary elevations.	such a
A low ESR can be seen with conditions that inhibit the normal sedimentation of red blood cells, such as a high red blood cell count (polycythaemia), significantly high white blood cell count (leucocytosis), and some protein abnormalities. Some changes in red cell sh as sickle cells in sickle cell anaemia) also lower the ESR. NOTE: 1. ESR and C - reactive protein (C-RP) are both markers of inflammation. 2. Generally, ESR does not change as rapidly as does CRP, either at the start of inflammation or as it resolves. 3. CRP is not affected by as many other factors as is ESR, making it a better marker of inflammation. 4. If the ESR is elevated, it is typically a result of two types of proteins, globulins or fibrinogen. 5. Women tend to have a higher ESR, and menstruation and pregnancy can cause temporary elevations.	
as sickle cells in sickle cell anaemia) also lower the ESR. NOTE: 1. ESR and C - reactive protein (C-RP) are both markers of inflammation. 2. Generally, ESR does not change as rapidly as does CRP, either at the start of inflammation or as it resolves. 3. CRP is not affected by as many other factors as is ESR, making it a better marker of inflammation. 4. If the ESR is elevated, it is typically a result of two types of proteins, globulins or fibrinogen. 5. Women tend to have a higher ESR, and menstruation and pregnancy can cause temporary elevations.	
NOTE: 1. ESR and C - reactive protein (C-RP) are both markers of inflammation. 2. Generally, ESR does not change as rapidly as does CRP, either at the start of inflammation or as it resolves. 3. CRP is not affected by as many other factors as is ESR, making it a better marker of inflammation. 4. If the ESR is elevated, it is typically a result of two types of proteins, globulins or fibrinogen. 5. Women tend to have a higher ESR, and menstruation and pregnancy can cause temporary elevations.	ape (su
 Generally, ESR does not change as rapidly as does CRP, either at the start of inflammation or as it resolves. CRP is not affected by as many other factors as is ESR, making it a better marker of inflammation. If the ESR is elevated, it is typically a result of two types of proteins, globulins or fibrinogen. Women tend to have a higher ESR, and menstruation and pregnancy can cause temporary elevations. 	
3. CRP is not affected by as many other factors as is ESR, making it a better marker of inflammation. 4. If the ESR is elevated, it is typically a result of two types of proteins, globulins or fibrinogen. 5. Women tend to have a higher ESR, and menstruation and pregnancy can cause temporary elevations.	
4. If the ESR is elevated, it is typically a result of two types of proteins, globulins or fibrinogen. 5. Women tend to have a higher ESR, and menstruation and pregnancy can cause temporary elevations.	
5. Women tend to have a higher ESR, and menstruation and pregnancy can cause temporary elevations. 6. Drugs such as devtran, methyldopa, oral contracentives, penicillamine procainamide, theophylline, and vitamin A can increase FSR.	
aspirin, cortisone, and quinine may decrease it	while
	while



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS , MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST

NOT VALID FOR MEDICO LEGAL PURPOSE



A PIONEER DIAGNOSTIC CENTRE

🔽 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. MANILA VAID				
AGE/ GENDER	: 45 YRS/FEMALE	PA	FIENT ID	: 16779	80
COLLECTED BY	:	REG	G. NO./LAB NO.	: 1224	11210013
REFERRED BY	:	REG	GISTRATION DATE	:21/No	ov/2024 11:36 AM
BARCODE NO.	: 12505766	CO	LECTION DATE	:21/No	ov/2024 11:38AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE II	NSTITUTE RE	PORTING DATE	:21/No	ov/2024 01:30PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD,	AMBALA CITY - HARYA	NA		
Test Name		Value	Unit		Biological Reference interva
	CLIN	ICAL CHEMISTR	Y/BIOCHEMIST	'RY	
		GLUCOSE FA	STING (F)		
GLUCOSE FASTING by GLUCOSE OXIDASE	(F): PLASMA - PEROXIDASE (GOD-POD)	81.56	mg/dL		NORMAL: < 100.0 PREDIABETIC: 100.0 - 125.0 DIABETIC: > 0R = 126.0
INTERPRETATION	I AMERICAN DIABETES ASSOCI	ATION GUIDELINES:			DIABETIC: > 0

A fasting plasma glucose level below 100 mg/dl is considered normal.
 A fasting plasma glucose level between 100 - 125 mg/dl is considered as glucose intolerant or prediabetic. A fasting and post-prandial blood test (after consumption of 75 gms of glucose) is recommended for all such patients.
 A fasting plasma glucose level of above 125 mg/dl is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for all such patients. A fasting plasma glucose level in excess of 125 mg/dl on both occasions is confirmatory for diabetic state.





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



A PIONEER DIAGNOSTIC CENTRE

【 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. MANILA VAID			
AGE/ GENDER	: 45 YRS/FEMALE		PATIENT ID	: 1677980
COLLECTED BY	:		REG. NO./LAB NO.	: 122411210013
REFERRED BY	:		REGISTRATION DATE	: 21/Nov/2024 11:36 AM
BARCODE NO.	: 12505766		COLLECTION DATE	: 21/Nov/2024 11:38AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INS	TITUTE	REPORTING DATE	: 21/Nov/2024 01:30PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, A	MBALA CITY - H	IARYANA	
Test Name		Value	Unit	Biological Reference interval
		LIPID PI	ROFILE : BASIC	
CHOLESTEROL TO	TAL: SERUM	164.4	mg/dL	OPTIMAL: < 200.0
by CHOLESTEROL O	KIDASE PAP		ů	BORDERLINE HIGH: 200.0 - 239.0 HIGH CHOLESTEROL: > OR =
TDICI VCEDIDEC. C	EDUM	70.01	TI / t	240.0
TRIGLYCERIDES: S by GLYCEROL PHOSE	ERUM PHATE OXIDASE (ENZYMATIC)	76.91	mg/dL	OPTIMAL: < 150.0 BORDERLINE HIGH: 150.0 - 199.0 HIGH: 200.0 - 499.0 VERY HIGH: > OR = 500.0
HDL CHOLESTERO by SELECTIVE INHIBIT	L (DIRECT): SERUM 70N	70.44	mg/dL	LOW HDL: < 30.0 BORDERLINE HIGH HDL: 30.0 60.0 HIGH HDL: > OR = 60.0
LDL CHOLESTERO	L: SERUM ECTROPHOTOMETRY	78.58	mg/dL	OPTIMAL: < 100.0 ABOVE OPTIMAL: 100.0 - 129. BORDERLINE HIGH: 130.0 - 159.0 HIGH: 160.0 - 189.0 VERY HIGH: > OR = 190.0
NON HDL CHOLES' by calculated, spe		93.96	mg/dL	OPTIMAL: < 130.0 ABOVE OPTIMAL: 130.0 - 159. BORDERLINE HIGH: 160.0 - 189.0 HIGH: 190.0 - 219.0 VERY HIGH: > OR = 220.0
VLDL CHOLESTER	DL: SERUM ECTROPHOTOMETRY	15.38	mg/dL	0.00 - 45.00
TOTAL LIPIDS: SEF		405.71	mg/dL	350.00 - 700.00
CHOLESTEROL/HI		2.33	RATIO	LOW RISK: 3.30 - 4.40 AVERAGE RISK: 4.50 - 7.0 MODERATE RISK: 7.10 - 11.0 HIGH RISK: > 11.0



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS , MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST



A PIONEER DIAGNOSTIC CENTRE

🔽 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. MANILA VAID		
AGE/ GENDER	: 45 YRS/FEMALE	PATIENT ID	: 1677980
COLLECTED BY	:	REG. NO./LAB NO.	: 122411210013
REFERRED BY	:	REGISTRATION DATE	: 21/Nov/2024 11:36 AM
BARCODE NO.	: 12505766	COLLECTION DATE	: 21/Nov/2024 11:38AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	REPORTING DATE	: 21/Nov/2024 01:30PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY -	HARYANA	

Test Name	Value	Unit	Biological Reference interval
LDL/HDL RATIO: SERUM by Calculated, SPECTROPHOTOMETRY	1.12	RATIO	LOW RISK: 0.50 - 3.0 MODERATE RISK: 3.10 - 6.0 HIGH RISK: > 6.0
TRIGLYCERIDES/HDL RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	1.09 ^L	RATIO	3.00 - 5.00

INTERPRETATION:

1. Measurements in the same patient can show physiological analytical variations. Three serial samples 1 week apart are recommended for Total Cholesterol, Triglycerides, HDL & LDL Cholesterol.

2. As per NLA-2014 guidelines, all adults above the age of 20 years should be screened for lipid status. Selective screening of children above the age of 2 years with a family history of premature cardiovascular disease or those with at least one parent with high total cholesterol is recommended.

3. Low HDL levels are associated with increased risk for Atherosclerotic Cardiovascular disease (ASCVD) due to insufficient HDL being available

 Low hole to consider a structure of the process by which cholesterol is eliminated from peripheral tissues.
 NLA-2014 identifies Non HDL Cholesterol (an indicator of all atherogeniclipoproteins such as LDL, VLDL, IDL, Lpa, Chylomicron remnants) along with LDL-cholesterol as co- primary target for cholesterol lowering therapy. Note that major risk factors can modify treatment goals for LDL & Non HDL

5. Additional testing for Apolipoprotein B, hsCRP,Lp(a) & LP-PLA2 should be considered among patients with moderate risk for ASCVD for risk refinement



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)





A PIONEER DIAGNOSTIC CENTRE

【 0171-2532620, 8222896961 ☑ pkrjainhealthcare@gmail.com

NAME	: Mrs. MANILA VAID			
AGE/ GENDER	: 45 YRS/FEMALE]	PATIENT ID	: 1677980
COLLECTED BY	:	1	REG. NO./LAB NO.	: 122411210013
REFERRED BY	:	1	REGISTRATION DATE	: 21/Nov/2024 11:36 AM
BARCODE NO.	: 12505766		COLLECTION DATE	: 21/Nov/2024 11:38AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITU	UTE	REPORTING DATE	: 21/Nov/2024 01:30PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBA	LA CITY - HAF	RYANA	
Test Name		Value	Unit	Biological Reference interva
	LIVER	FUNCTION	TEST (COMPLETE)	
BILIRUBIN TOTAL: by diazotization, sf		0.71	mg/dL	INFANT: 0.20 - 8.00 ADULT: 0.00 - 1.20
	C (CONJUGATED): SERUM	0.15	mg/dL	0.00 - 0.40
BILIRUBIN INDIRE by CALCULATED, SPE	CT (UNCONJUGATED): SERUM	0.56	mg/dL	0.10 - 1.00
SGOT/AST: SERUM by IFCC, WITHOUT PY	RIDOXAL PHOSPHATE	17.06	U/L	7.00 - 45.00
SGPT/ALT: SERUM by IFCC, WITHOUT PY	RIDOXAL PHOSPHATE	19.49	U/L	0.00 - 49.00
AST/ALT RATIO: SI by calculated, spe		0.88	RATIO	0.00 - 46.00
ALKALINE PHOSPH by Para NITROPHEN PROPANOL	IATASE: SERUM YL PHOSPHATASE BY AMINO METHYL	73.55	U/L	40.0 - 130.0
GAMMA GLUTAMY	L TRANSFERASE (GGT): SERUM	25.61	U/L	0.00 - 55.0

ALKALINE PHOSPHATASE: SERUM by PARA NITROPHENYL PHOSPHATASE BY AMINO METHYL PROPANOL	73.55	U/L
GAMMA GLUTAMYL TRANSFERASE (GGT): SERUM by szasz, spectrophtometry	25.61	U/L
TOTAL PROTEINS: SERUM by biuret, spectrophotometry	6.73	gm/dL
ALBUMIN: SERUM by BROMOCRESOL GREEN	4.49	gm/dL
GLOBULIN: SERUM by CALCULATED, SPECTROPHOTOMETRY	2.24 ^L	gm/dL
A : G RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	2	RATIO

INTERPRETATION

NOTE: - To be correlated in individuals having SGOT and SGPT values higher than Normal Referance Range. USE: - Differential diagnosis of diseases of hepatobiliary system and pancreas.

INCREASED:

DRUG HEPATOTOXICITY	> 2
ALCOHOLIC HEPATITIS	> 2 (Highly Suggestive)
CIRRHOSIS	1.4 - 2.0
INTRAHEPATIC CHOLESTATIS	> 1.5
HEPATOCELLULAR CARCINOMA & CHRONIC HEPATITIS	> 1.3 (Slightly Increased)





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)

440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600, REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)



6.20 - 8.00

3.50 - 5.50

2.30 - 3.50

1.00 - 2.00



A PIONEER DIAGNOSTIC CENTRE

【 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. MANILA VAID		
AGE/ GENDER	: 45 YRS/FEMALE	PATIENT ID	: 1677980
COLLECTED BY	:	REG. NO./LAB NO.	: 122411210013
REFERRED BY	:	REGISTRATION DATE	: 21/Nov/2024 11:36 AM
BARCODE NO.	: 12505766	COLLECTION DATE	: 21/Nov/2024 11:38AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	REPORTING DATE	: 21/Nov/2024 01:30PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY	- HARYANA	

|--|

DECREASED:

1. Acute Hepatitis due to virus, drugs, toxins (with AST increased 3 to 10 times upper limit of normal)

2. Extra Hepatic cholestatis: 0.8 (normal or slightly decreased).

NORMAL	< 0.65
GOOD PROGNOSTIC SIGN	0.3 - 0.6
POOR PROGNOSTIC SIGN	1.2 - 1.6



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS , MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST





A PIONEER DIAGNOSTIC CENTRE

🔽 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME :	Mrs. MANILA VAID				
AGE/ GENDER : 45 YRS/FEMALE			PATIENT ID	: 1677980	
COLLECTED BY :		REG. NO./LAB NO.		: 122411210013	
REFERRED BY :			REGISTRATION DATE	: 21/Nov/2024 11:36 AM	
BARCODE NO. :	DE NO. : 12505766 COLLE		COLLECTION DATE	: 21/Nov/2024 11:38AM	
CLIENT CODE.	P.K.R JAIN HEALTHCARE INSTI	ITUTE REPORTING DATE		: 21/Nov/2024 01:30PM	
CLIENT ADDRESS :	NASIRPUR, HISSAR ROAD, AME	BALA CITY - HA	RYANA		
Test Name		Value	Unit	Biological Reference interva	
	KIDNE	Y FUNCTIO	N TEST (COMPLETE)		
UREA: SERUM by UREASE - GLUTAMATE	DEHYDROGENASE (GLDH)	20.61	mg/dL	10.00 - 50.00	
CREATININE: SERUM by ENZYMATIC, SPECTRO	PHOTOMETERY	0.62	mg/dL	0.40 - 1.20	
BLOOD UREA NITROG by CALCULATED, SPECTE		9.63	mg/dL	7.0 - 25.0	
BLOOD UREA NITROG RATIO: SERUM by CALCULATED, SPECTF	EN (BUN)/CREATININE	15.53	RATIO	10.0 - 20.0	
UREA/CREATININE R. by CALCULATED, SPECTE		33.24	RATIO		
URIC ACID: SERUM by URICASE - OXIDASE PL	EROXIDASE	3.18	mg/dL	2.50 - 6.80	
CALCIUM: SERUM by ARSENAZO III, SPECTR	OPHOTOMETRY	9.7	mg/dL	8.50 - 10.60	
	M E, SPECTROPHOTOMETRY	2.76	mg/dL	2.30 - 4.70	
ELECTROLYTES SODIUM: SERUM		137.7	mmol/L	135.0 - 150.0	
by ISE (ION SELECTIVE EL POTASSIUM: SERUM by ISE (ION SELECTIVE EL		3.98	mmol/L	3.50 - 5.00	
CHLORIDE: SERUM by ISE (ION SELECTIVE EL		103.28	mmol/L	90.0 - 110.0	
	ULAR FILTERATION RATE				
ESTIMATED GLOMERI (eGFR): SERUM by CALCULATED INTERPRETATION:	ULAR FILTERATION RATE	111.8			

INTERPRETATION:

To differentiate between pre- and post renal azotemia.

INCREASED RATIO (>20:1) WITH NORMAL CREATININE:

1. Prerenal azotemia (BUN rises without increase in creatinine) e.g. heart failure, salt depletion, dehydration, blood loss) due to decreased glomerular filtration rate.

2. Catabolic states with increased tissue breakdown.

3. GI haemorrhage.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)



A PIONEER DIAGNOSTIC CENTRE

🔽 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. MANILA VAID		
AGE/ GENDER	: 45 YRS/FEMALE	PATIENT ID	: 1677980
COLLECTED BY	:	REG. NO./LAB NO.	: 122411210013
REFERRED BY	:	REGISTRATION DAT	TE : 21/Nov/2024 11:36 AM
BARCODE NO.	: 12505766	COLLECTION DATE	: 21/Nov/2024 11:38AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUT	TE REPORTING DATE	: 21/Nov/2024 01:30PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA	A CITY - HARYANA	
Test Name		Value Unit	Biological Reference interval
 Postrenal azotemia Prerenal azotemia DECREASED RATIO (< Acute tubular necessary 	20:1) WITH ELEVATED CREATININE LEVEN a (BUN rises disproportionately more the superimposed on renal disease. 10:1) WITH DECREASED BUN :		iropathy).
 Severe liver diseas Other causes of de Repeated dialysis Inherited hyperam 	nd starvation.	blood).	

3. Muscular patients who develop renal failure.

INAPPROPIATE RATIO:

1. Diabetic ketoacidosis (acetoacetate causes false increase in creatinine with certain methodologies, resulting in normal ratio when dehydration should produce an increased BUN/creatinine ratio).

Cephalosporin therapy (interferes with creatinine measurement).
 ESTIMATED GLOMERULAR FILTERATION RATE:

CKD STAGE	DESCRIPTION	GFR (mL/min/1.73m2)	ASSOCIATED FINDINGS No proteinuria	
G1	Normal kidney function	>90		
G2	Kidney damage with normal or high GFR	>90	Presence of Protein , Albumin or cast in urine	
G3a	Mild decrease in GFR	60 -89		
G3b	Moderate decrease in GFR	30-59		
G4	Severe decrease in GFR	15-29		
G5	Kidney failure	<15		



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)







A PIONEER DIAGNOSTIC CENTRE

0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. MANILA VAID		
AGE/ GENDER	: 45 YRS/FEMALE	PATIENT ID	: 1677980
COLLECTED BY	:	REG. NO./LAB NO.	: 122411210013
REFERRED BY	:	REGISTRATION DATE	: 21/Nov/2024 11:36 AM
BARCODE NO.	: 12505766	COLLECTION DATE	: 21/Nov/2024 11:38AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	REPORTING DATE	: 21/Nov/2024 01:30PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - H	IARYANA	

Test Name	Value	Unit	Biological Reference interval

COMMENTS:

1. Estimated Glomerular filtration rate (eGFR) is the sum of filtration rates in all functioning nephrons and so an estimation of the GFR provides a measure of functioning nephrons of the kidney. 2. eGFR calculated using the 2009 CKD-EPI creatinine equation and GFR category reported as per KDIGO guideline 2012

3. In patients, with eGFR creatinine between 45-59 ml/min/1.73 m2 (G3) and without any marker of Kidney damage, It is recommended to measure eGFR with Cystatin C for confirmation of CKD

4. eGFR category G1 OR G2 does not fullfill the criteria for CKD, in the absence of evidence of Kidney Damage 5. In a suspected case of Acute Kidney Injury (AKI), measurement of eGFR should be done after 48-96 hours of any Intervention or procedure 6. eGFR calculated by Serum Creatinine may be less accurate due to certain factors like Race, Muscle Mass, Diet, Certain Drugs. In such cases, eGFR should be calculated using Serum Cystatin C 7. A decrease in eGFR implies either progressive renal disease, or a reversible process causing decreased nephron function (eg, severe dehydration).

ADVICE:

KDIGO guideline, 2012 recommends Chronic Kidney Disease (CKD) should be classified based on cause, eGFR category and Albuminuria (ACR) category. GFR & ACR category combined together reflect risk of progression and helps Clinician to identify the individual who are progressing at more rapid rate than anticipated



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)





A PIONEER DIAGNOSTIC CENTRE

💟 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

45 YRS/FEMALE	РА	TIENT ID	1677000	
			: 1677980	
	RE	G. NO./LAB NO.	: 122411210013	
	RE	GISTRATION DATE	: 21/Nov/2024 11:36 AM	
12505766	CO	LLECTION DATE	: 21/Nov/2024 11:38AM	
P.K.R JAIN HEALTHCARE INSTITUT	TE RE	PORTING DATE	: 21/Nov/2024 01:30PM	
NASIRPUR, HISSAR ROAD, AMBALA	A CITY - HARYA	NA		
	Value	Unit	Biological Reference interval	
	ENDOCRI	NOLOGY		
THYRO	ID FUNCTIO	ON TEST: TOTAL		
	1.27	ng/mL	0.35 - 1.93	
THYROXINE (T4): SERUM by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)		µgm/dL	4.87 - 12.60	
	1.47	µIU/mL	0.35 - 5.50	
SENSITIVE				
	NASIRPUR, HISSAR ROAD, AMBALA THYRO (T3): SERUM CENT MICROPARTICLE IMMUNOASSAY) RUM	12505766 CO P.K.R JAIN HEALTHCARE INSTITUTE RE NASIRPUR, HISSAR ROAD, AMBALA CITY - HARVA Value ENDOCRI THYROD FUNCTION (T3): SERUM (T3): SERUM (T4):	P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA Value Unit ENDOCRINOLOGY THYROUD FUNCTION TEST: TOTAL (T3): SERUM 1.27 ng/mL CENT MICROPARTICLE IMMUNOASSAY) RUM 7.24 µgm/dL CENT MICROPARTICLE IMMUNOASSAY) NG HORMONE (TSH): SERUM 1.47 µIU/mL	

TSH levels are subject to circadian variation, reaching peak levels between 2-4 a.m and at a minimum between 6-10 pm. The variation is of the order of 50%. Hence time of the day has influence on the measured serum TSH concentrations. TSH stimulates the production and secretion of the metabolically active hormones, thyroxine (T4) and triiodothyronine (T3). Failure at any level of regulation of the hypothalamic-pituitary-thyroid axis will result in either underproduction (hypothyroidism) or overproduction(hyperthyroidism) of T4 and/or T3.

CLINICAL CONDITION	Т3	T4	TSH
Primary Hypothyroidism: Reduced		Reduced Increased (Significant	
Subclinical Hypothyroidism:	Normal or Low Normal	Normal or Low Normal High	
Primary Hyperthyroidism:	Increased	Increased	Reduced (at times undetectable)
Subclinical Hyperthyroidism:	Normal or High Normal	Normal or High Normal	Reduced

LIMITATIONS:-

1. T3 and T4 circulates in reversibly bound form with Thyroid binding globulins (TBG), and to a lesser extent albumin and Thyroid binding Pre Albumin so conditions in which TBG and protein levels alter such as pregnancy, excess estrogens, androgens, anabolic steroids and glucocorticoids may falsely affect the T3 and T4 levels and may cause false thyroid values for thyroid function tests.

2. Normal levels of T4 can also be seen in Hyperthyroid patients with :T3 Thyrotoxicosis, Decreased binding capacity due to hypoproteinemia or ingestion of certain drugs (e.g.: phenytoin , salicylates).

3. Serum T4 levels in neonates and infants are higher than values in the normal adult , due to the increased concentration of TBG in neonate serum.

4. TSH may be normal in central hypothyroidism, recent rapid correction of hyperthyroidism or hypothyroidism, pregnancy, phenytoin therapy.

TRIIODOTH	YRONINE (T3)	THYROXINE (T4)		THYROID STIMU	LATING HORMONE (TSH)
Age	Refferance Range (ng/mL)	Age	Refferance Range (µg/dL)	Age	Reference Range (μIU/mL)
0-7 Days	0.20 - 2.65	0 - 7 Days	5.90 - 18.58	0 - 7 Days	2.43 - 24.3
7 Days - 3 Months	0.36 - 2.59	7 Days - 3 Months	6.39 - 17.66	7 Days - 3 Months	0.58 - 11.00
3 - 6 Months	0.51 - 2.52	3 - 6 Months	6.75 - 17.04	3 Days – 6 Months	0.70 - 8.40
6 - 12 Months	0.74 - 2.40	6 - 12 Months	7.10 - 16.16	6 – 12 Months	0.70 - 7.00





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)





A PIONEER DIAGNOSTIC CENTRE © 0171-2532620, 8222896961 ⊠ pkrjainhealthcare@gmail.com

NAME	: Mrs. MANILA VAID			
AGE/ GENDER	: 45 YRS/FEMALE	PATIENT ID	: 1677980	
COLLECTED BY	:	REG. NO./LAB NO.	: 122411210013	
REFERRED BY	:	REGISTRATION DATE	: 21/Nov/2024 11:36 AM	
BARCODE NO.	: 12505766	COLLECTION DATE	: 21/Nov/2024 11:38AM	
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	REPORTING DATE	: 21/Nov/2024 01:30PM	
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA			

Test Name		Value Unit		t	Biological Reference interval	
1 - 10 Years	0.92 - 2.28	1 - 10 Years	6.00 - 13.80	1 – 10 Years	0.60 - 5.50	
11- 19 Years	0.35 - 1.93	11 - 19 Years	4.87- 13.20	11 – 19 Years	0.50 - 5.50	
> 20 years (Adults)	0.35 - 1.93	> 20 Years (Adults)	4.87 - 12.60	> 20 Years (Adults)	0.35-5.50	
	RECOM	MENDATIONS OF TSH LE	VELS DURING PRE	GNANCY (µIU/mL)		
	1st Trimester			0.10 - 2.50		
	2nd Trimester			0.20 - 3.00		
	3rd Trimester			0.30 - 4.10		

INCREASED TSH LEVELS:

1. Primary or untreated hypothyroidism may vary from 3 times to more than 100 times normal depending upon degree of hypofunction.

2. Hypothyroid patients receiving insufficient thyroid replacement therapy.

3.Hashimotos thyroiditis

4.DRUGS: Amphetamines, iodine containing agents & dopamine antagonist.

5.Neonatal period, increase in 1st 2-3 days of life due to post-natal surge

DECREASED TSH LEVELS:

1.Toxic multi-nodular goiter & Thyroiditis.

2. Over replacement of thyroid hormone in treatment of hypothyroidism.

3. Autonomously functioning Thyroid adenoma

4.Secondary pituitary or hypothalamic hypothyroidism

5. Acute psychiatric illness

6.Severe dehydration.

7.DRUGS: Glucocorticoids, Dopamine, Levodopa, T4 replacement therapy, Anti-thyroid drugs for thyrotoxicosis.

8. Pregnancy: 1st and 2nd Trimester



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)





NAME

: Mrs. MANILA VAID

PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

【 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

AGE/ GENDER: 45 YRS/FEMALECOLLECTED BY:REFERRED BY:BARCODE NO.: 12505766CLIENT CODE.: P.K.R JAIN HEALTHCARE INSTI		PATIENT ID REG. NO./LAB NO. REGISTRATION DATE COLLECTION DATE ITUTE REPORTING DATE		: 1677980 : 122411210013 : 21/Nov/2024 11:36 AM : 21/Nov/2024 11:38AM : 21/Nov/2024 01:30PM									
						CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AM	SIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA					
						Test Name		Value	Unit	Biological Reference interva			
								CLINICAL PATHO	LOGY				
							URINE RO	UTINE & MICROSCOP		ATION			
PHYSICAL EXAMIN	NATION												
QUANTITY RECIEVED by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY COLOUR		10	ml										
		PALE YELLOW		PALE YELLOW									
by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY TRANSPARANCY		TURBID		CLEAR									
by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY SPECIFIC GRAVITY		1.02 PKR		1.002 - 1.030									
by DIP STICK/REFLEC CHEMICAL EXAMI	TANCE SPECTROPHOTOMETRY												
REACTION		ACIDIC											
by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY PROTEIN		NEGATIVE (-ve)		NEGATIVE (-ve)									
by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY SUGAR		NEGATIVE (-ve)		NEGATIVE (-ve)									
by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY pH		5.5		5.0 - 7.5									
by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY													
BILIRUBIN by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY		NEGATIVE (-ve)		NEGATIVE (-ve)									
NITRITE by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY.		NEGATIVE (-ve)		NEGATIVE (-ve)									
UROBILINOGEN		NOT DETECTED	EU/dL	0.2 - 1.0									
by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY KETONE BODIES by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY BLOOD by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY ASCORBIC ACID by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY		NEGATIVE (-ve)		NEGATIVE (-ve)									
		NEGATIVE (-ve)		NEGATIVE (-ve)									
		NEGATIVE (-ve)		NEGATIVE (-ve)									
MICROSCOPIC EXA			/IIDE	0.2									
RED BLOOD CELLS	(RDUS)	NEGATIVE (-ve)	/HPF	0 - 3									



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS , MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST

NOT VALID FOR MEDICO LEGAL PURPOSE



A PIONEER DIAGNOSTIC CENTRE

【 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. MANILA VAID				
AGE/ GENDER	: 45 YRS/FEMALE	PATIENT ID	: 1677980		
COLLECTED BY	:	REG. NO./LAB NO.	: 122411210013		
REFERRED BY	:	REGISTRATION DATE	: 21/Nov/2024 11:36 AM		
BARCODE NO.	: 12505766	COLLECTION DATE	: 21/Nov/2024 11:38AM		
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	REPORTING DATE	: 21/Nov/2024 01:30PM		
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA				
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA				

Test Name	Value	Unit	Biological Reference interval
by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT			
PUS CELLS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	12-15	/HPF	0 - 5
EPITHELIAL CELLS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	6-8	/HPF	ABSENT
CRYSTALS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	NEGATIVE (-ve)		NEGATIVE (-ve)
CASTS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	NEGATIVE (-ve)		NEGATIVE (-ve)
BACTERIA by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	NEGATIVE (-ve)		NEGATIVE (-ve)
OTHERS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	NEGATIVE (-ve)		NEGATIVE (-ve)
TRICHOMONAS VAGINALIS (PROTOZOA) by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	ABSENT		ABSENT

* End Of Report



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS , MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST

