A PIONEER DIAGNOSTIC CENTRE

【 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME : N	Irs. KAMINI			
AGE/ GENDER : 6	0 YRS/FEMALE		PATIENT ID	: 1606656
COLLECTED BY :			REG. NO./LAB NO.	: 122411260010
REFERRED BY :			<b>REGISTRATION DATE</b>	: 26/Nov/2024 10:58 AM
BARCODE NO. : 1	2505850		COLLECTION DATE	: 26/Nov/2024 11:28AM
CLIENT CODE. : P	K.R JAIN HEALTHCARE INSTITU	ГЕ	REPORTING DATE	: 26/Nov/2024 01:00PM
CLIENT ADDRESS : N	ASIRPUR, HISSAR ROAD, AMBAL	A CITY - HAI	RYANA	
Test Name		Value	Unit	Biological Reference interval
	SWASTI	HYA WEI	LLNESS PANEL: 1.4	
	СОМР	LETE BLO	OOD COUNT (CBC)	
<u>RED BLOOD CELLS (R</u>	BCS) COUNT AND INDICES			
HAEMOGLOBIN (HB) by CALORIMETRIC		12.4	gm/dL	12.0 - 16.0
RED BLOOD CELL (RBC	) COUNT SING, ELECTRICAL IMPEDENCE	4.24	Millions/c	emm 3.50 - 5.00
PACKED CELL VOLUME	(PCV) MATED HEMATOLOGY ANALYZER	36.4 <sup>L</sup>	%	37.0 - 50.0
MEAN CORPUSCULAR V		86	KR fl	80.0 - 100.0
MEAN CORPUSCULAR I by CALCULATED BY AUTOR	HAEMOGLOBIN (MCH) MATED HEMATOLOGY ANALYZER	29.3	pg	27.0 - 34.0
	HEMOGLOBIN CONC. (MCHC) MATED HEMATOLOGY ANALYZER	34.1	g/dL	32.0 - 36.0
RED CELL DISTRIBUTION	ON WIDTH (RDW-CV) MATED HEMATOLOGY ANALYZER	12.7	%	11.00 - 16.00
RED CELL DISTRIBUTION	ON WIDTH (RDW-SD) mated hematology analyzer	41.6	fL	35.0 - 56.0
MENTZERS INDEX by CALCULATED		20.28	RATIO	BETA THALASSEMIA TRAIT: < 13.0 IRON DEFICIENCY ANEMIA: >13.0
GREEN & KING INDEX by CALCULATED		25.81	RATIO	BETA THALASSEMIA TRAIT:< 65.0 IRON DEFICIENCY ANEMIA: > 65.0
WHITE BLOOD CELLS				
TOTAL LEUCOCYTE CO by flow cytometry by : DIFFERENTIAL LEUCO	SF CUBE & MICROSCOPY	5110	/cmm	4000 - 11000
NEUTROPHILS by flow cytometry by s		55	%	50 - 70
LYMPHOCYTES		37	%	20 - 40

TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT

**DR.VINAY CHOPRA** CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS , MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST

**NOT VALID FOR MEDICO LEGAL PURPOSE** 



A PIONEER DIAGNOSTIC CENTRE

【 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. KAMINI			
AGE/ GENDER	: 60 YRS/FEMALE		PATIENT ID	: 1606656
<b>COLLECTED BY</b>	:		REG. NO./LAB NO.	: 122411260010
<b>REFERRED BY</b>	:		<b>REGISTRATION DATE</b>	: 26/Nov/2024 10:58 AM
BARCODE NO.	: 12505850		<b>COLLECTION DATE</b>	: 26/Nov/2024 11:28AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTIT	TUTE	<b>REPORTING DATE</b>	: 26/Nov/2024 01:00PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMB	ALA CITY - HA	ARYANA	
Test Name		Value	Unit	Biological Reference interval
by FLOW CYTOMETRY	Y BY SF CUBE & MICROSCOPY			
EOSINOPHILS	Y BY SF CUBE & MICROSCOPY	2	%	1 - 6
MONOCYTES		6	%	2 - 12
•	Y BY SF CUBE & MICROSCOPY	0	0/	0 1
BASOPHILS by FLOW CYTOMETRY	Y BY SF CUBE & MICROSCOPY	0	%	0 - 1
ABSOLUTE LEUKO	CYTES (WBC) COUNT			
ABSOLUTE NEUTR		2811	/cmm	2000 - 7500
ABSOLUTE LYMPH	Y BY SF CUBE & MICROSCOPY OCYTF COUNT	1891 <sup>L</sup>	/cmm	800 - 4900
by FLOW CYTOMETRY	Y BY SF CUBE & MICROSCOPY		KR	000 1000
ABSOLUTE EOSINC	PHIL COUNT Y BY SF CUBE & MICROSCOPY	102	/cmm	40 - 440
ABSOLUTE MONOC		307	/cmm	80 - 880
	Y BY SF CUBE & MICROSCOPY	0	,	0, 110
ABSOLUTE BASOPI	HIL COUN I Y BY SF CUBE & MICROSCOPY	0	/cmm	0 - 110
PLATELETS AND C	THER PLATELET PREDICTIVE	MARKERS.		
PLATELET COUNT by HYDRO DYNAMIC F	(PLT) OCUSING, ELECTRICAL IMPEDENCE	183000	/cmm	150000 - 450000
PLATELETCRIT (PC		0.17	%	0.10 - 0.36
by HYDRO DYNAMIC F MEAN PLATELET V	OCUSING, ELECTRICAL IMPEDENCE	9	fL	6.50 - 12.0
by HYDRO DYNAMIC F	OCUSING, ELECTRICAL IMPEDENCE			
	CELL COUNT (P-LCC)	39000	/cmm	30000 - 90000
PLATELET LARGE	CELL RATIO (P-LCR) FOCUSING, ELECTRICAL IMPEDENCE	21.3	%	11.0 - 45.0
PLATELET DISTRIE	BUTION WIDTH (PDW) FOCUSING, ELECTRICAL IMPEDENCE	15.9	%	15.0 - 17.0
	CTED ON EDTA WHOLE BLOOD			



**DR.VINAY CHOPRA** CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS , MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST

NOT VALID FOR MEDICO LEGAL PURPOSE



**A PIONEER DIAGNOSTIC CENTRE** 

💟 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. KAMINI				
AGE/ GENDER	: 60 YRS/FEMALE	PA	ATIENT ID	: 1606656	
COLLECTED BY	:	RI	EG. NO./LAB NO.	: 122411260010	)
REFERRED BY		RI	EGISTRATION DATE	: 26/Nov/2024 10	:58 AM
BARCODE NO.	: 12505850		DLLECTION DATE	: 26/Nov/2024 11	
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INST	TTUTE RI	EPORTING DATE	: 26/Nov/2024 05	:IIPM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AM	IBALA CITY - HARY	ANA		
Test Name		Value	Unit	Biologic	al Reference interval
WHOLE BLOOD	EMOGLOBIN (HbA1c):	7.3 <sup>H</sup>	MOGLOBIN (HBA1( %	4.0 - 6.4	
by HPLC (HIGH PERFO ESTIMATED AVERA	RMANCE LIQUID CHROMATOGRAPHY) IGE PLASMA GLUCOSE RMANCE LIQUID CHROMATOGRAPHY)	162.81 <sup>H</sup>	mg/dL	60.00 - 1	140.00
[	AS PER AMERICAN	DIABETES ASSOCIATI	ON (ADA):		Г
	REFERENCE GROUP		OSYLATED HEMOGLOGIB	(HBAIC) in %	-
Non di	abetic Adults >= 18 years	DI	<5.7		
	t Risk (Prediabetes)		5.7 – 6.4		
D	iagnosing Diabetes		>= 6.5		
			Age > 19 Years		
Thoropout	is goals for allocamic control		Therapy:	< 7.0	4
Therapeut	ic goals for glycemic control	Actions S		>8.0	-
		Coal of	Age < 19 Years therapy:	<7.5	
		0000		51.0	

1.Glycosylated hemoglobin (HbA1c) test is three monthly monitoring done to assess compliace with therapeutic regimen in diabetic patients. 2.Since Hb1c reflects long term fluctuations in blood glucose concentration, a diabetic patient who has recently under good control may still have high concentration of HbAlc. Converse is true for a diabetic previously under good control but now poorly controlled.

3. Target goals of < 7.0 % may be beneficial in patients with short duration of diabetes, long life expectancy and no significant cardiovascular disease. In patients with significant complications of diabetes, limited life expectancy or extensive co-morbid conditions, targetting a goal of < 7.0% may not be appropriate.

4.High HbA1c (>9.0 -9.5 %) is strongly associated with risk of development and rapid progression of microvascular and nerve complications 5.Any condition that shorten RBC life span like acute blood loss, hemolytic anemia falsely lower HbA1c results.

6.HbA1c results from patients with HbSS,HbSC and HbD must be interpreted with caution, given the pathological processes including anemia, increased red cell turnover, and transfusion requirement that adversely impact HbA1c as a marker of long-term gycemic control.

7.Specimens from patients with polycythemia or post-splenctomy may exhibit increse in HbA1c values due to a somewhat longer life span of the red cells.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)



A PIONEER DIAGNOSTIC CENTRE

🔽 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. KAMINI			
AGE/ GENDER	: 60 YRS/FEMALE	PA	ATIENT ID	: 1606656
COLLECTED BY	:	RI	EG. NO./LAB NO.	: 122411260010
REFERRED BY	:	RI	EGISTRATION DATE	: 26/Nov/2024 10:58 AM
BARCODE NO.	: 12505850	CO	DLLECTION DATE	: 26/Nov/2024 11:28AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTIT	UTE <b>RI</b>	EPORTING DATE	: 26/Nov/2024 04:08PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBA	LA CITY - HARY	ANA	
Test Name		Value	Unit	Biological Reference interval
	ERYTHROC	YTE SEDIME	ENTATION RATE (	ESR)
by RED CELL AGGRE	DIMENTATION RATE (ESR) gation by capillary photometry	7	mm/1st	
INTERPRETATION: 1. ESR is a non-specifimmune disease but	does not tell the health practitioner	exactly where the	ne inflammation is in the	ion associated with infection, cancer and auto e body or what is causing it. pically used in conjunction with other test suc

A low ESR can be seen with conditions that inhibit the normal sedimentation of red blood cells, such as a high red blood cell count

(polycythaemia), significantly high white blood cell count (leucocytosis), and some protein abnormalities. Some changes in red cell shape (such as sickle cells in sickle cell anaemia) also lower the ESR. NOTE:

LER and C - reactive protein (C-RP) are both markers of inflammation.
 Generally, ESR does not change as rapidly as does CRP, either at the start of inflammation or as it resolves.
 CRP is not affected by as many other factors as is ESR, making it a better marker of inflammation.
 If the ESR is elevated, it is typically a result of two types of proteins, globulins or fibrinogen.
 Women tend to have a higher ESR, and menstruation and pregnancy can cause temporary elevations.
 Drugs such as dovtram, motbuling, and vities and vit

6. Drugs such as dextran, methyldopa, oral contraceptives, penicillamine procainamide, theophylline, and vitamin A can increase ESR, while aspirin, cortisone, and quinine may decrease it



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)

NOT VALID FOR MEDICO LEGAL PURPOSE



A PIONEER DIAGNOSTIC CENTRE

🔽 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

: Mrs. KAMINI			
: 60 YRS/FEMALE	PA	FIENT ID	: 1606656
:	RE	G. NO./LAB NO.	: 122411260010
:	RE	GISTRATION DATE	: 26/Nov/2024 10:58 AM
: 12505850	CO	LLECTION DATE	: 26/Nov/2024 11:28AM
: P.K.R JAIN HEALTHCARE INS	TITUTE <b>RE</b>	PORTING DATE	: 26/Nov/2024 01:00PM
: NASIRPUR, HISSAR ROAD, AM	IBALA CITY - HARYA	NA	
	Value	Unit	Biological Reference interva
CLINIC	AL CHEMISTR	Y/BIOCHEMIST	RY
		IT DIO CIILINIS I	
	GLUCOSE FA		
-	: 60 YRS/FEMALE : : 12505850 : P.K.R JAIN HEALTHCARE INS : NASIRPUR, HISSAR ROAD, AN	: 60 YRS/FEMALE PA' : REA : REA : 12505850 CO : P.K.R JAIN HEALTHCARE INSTITUTE REA : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYA	<ul> <li>: 60 YRS/FEMALE</li> <li>: 60 YRS/FEMALE</li> <li>: REG. NO./LAB NO.</li> <li>: REGISTRATION DATE</li> <li>: 12505850</li> <li>: COLLECTION DATE</li> <li>: P.K.R JAIN HEALTHCARE INSTITUTE</li> <li>: REPORTING DATE</li> <li>: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA</li> </ul>

A fasting plasma glucose level below 100 mg/dl is considered normal.
 A fasting plasma glucose level between 100 - 125 mg/dl is considered as glucose intolerant or prediabetic. A fasting and post-prandial blood test (after consumption of 75 gms of glucose) is recommended for all such patients.
 A fasting plasma glucose level of above 125 mg/dl is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for all such patients.
 A fasting plasma glucose level in excess of 125 mg/dl on both occasions is confirmatory for diabetic state.





**DR.VINAY CHOPRA** CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



A PIONEER DIAGNOSTIC CENTRE

【 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. KAMINI			
AGE/ GENDER	: 60 YRS/FEMALE		PATIENT ID	: 1606656
COLLECTED BY	:		REG. NO./LAB NO.	: 122411260010
REFERRED BY	:		<b>REGISTRATION DATE</b>	: 26/Nov/2024 10:58 AM
BARCODE NO.	: 12505850		COLLECTION DATE	: 26/Nov/2024 11:28AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INS	STITUTE	<b>REPORTING DATE</b>	: 26/Nov/2024 01:00PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, A	MBALA CITY - HA	ARYANA	
Test Name		Value	Unit	<b>Biological Reference interval</b>
		LIPID PR	OFILE : BASIC	
CHOLESTEROL TO by CHOLESTEROL OX		196.15	mg/dL	OPTIMAL: < 200.0 BORDERLINE HIGH: 200.0 - 239.0 HIGH CHOLESTEROL: > OR = 240.0
TRIGLYCERIDES: S by GLYCEROL PHOSF	ERUM phate oxidase (enzymatic)	140.46	mg/dL	OPTIMAL: < 150.0 BORDERLINE HIGH: 150.0 - 199.0 HIGH: 200.0 - 499.0 VERY HIGH: > OR = 500.0
HDL CHOLESTERO	L (DIRECT): SERUM	64.74	mg/dL	LOW HDL: < 30.0 BORDERLINE HIGH HDL: 30.0 60.0 HIGH HDL: > OR = 60.0
LDL CHOLESTEROI by CALCULATED, SPE		103.32	mg/dL	OPTIMAL: < 100.0 ABOVE OPTIMAL: 100.0 - 129 BORDERLINE HIGH: 130.0 - 159.0 HIGH: 160.0 - 189.0 VERY HIGH: > OR = 190.0
NON HDL CHOLEST by calculated, spe		131.41 <sup>H</sup>	mg/dL	OPTIMAL: < 130.0 ABOVE OPTIMAL: 130.0 - 159 BORDERLINE HIGH: 160.0 - 189.0 HIGH: 190.0 - 219.0 VERY HIGH: > OR = 220.0
VLDL CHOLESTER( by CALCULATED, SPE		28.09	mg/dL	0.00 - 45.00
TOTAL LIPIDS: SER by CALCULATED, SPE	RUM	532.76	mg/dL	350.00 - 700.00
CHOLESTEROL/HD by CALCULATED, SPE		3.03	RATIO	LOW RISK: 3.30 - 4.40 AVERAGE RISK: 4.50 - 7.0 MODERATE RISK: 7.10 - 11.0 HIGH RISK: > 11.0



**DR.VINAY CHOPRA** CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS , MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST

**NOT VALID FOR MEDICO LEGAL PURPOSE** 

440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. **REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)** 



🔽 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. KAMINI		
AGE/ GENDER	: 60 YRS/FEMALE	PATIENT ID	: 1606656
<b>COLLECTED BY</b>	:	<b>REG. NO./LAB NO.</b>	: 122411260010
<b>REFERRED BY</b>	:	<b>REGISTRATION DATE</b>	: 26/Nov/2024 10:58 AM
BARCODE NO.	: 12505850	<b>COLLECTION DATE</b>	: 26/Nov/2024 11:28AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	<b>REPORTING DATE</b>	: 26/Nov/2024 01:00PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY	- HARYANA	

Test Name	Value	Unit	<b>Biological Reference interval</b>
LDL/HDL RATIO: SERUM by calculated, spectrophotometry	1.6	RATIO	LOW RISK: 0.50 - 3.0 MODERATE RISK: 3.10 - 6.0 HIGH RISK: > 6.0
TRIGLYCERIDES/HDL RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	2.17 <sup>L</sup>	RATIO	3.00 - 5.00

#### **INTERPRETATION:**

1. Measurements in the same patient can show physiological analytical variations. Three serial samples 1 week apart are recommended for Total Cholesterol, Triglycerides, HDL & LDL Cholesterol.

2. As per NLA-2014 guidelines, all adults above the age of 20 years should be screened for lipid status. Selective screening of children above the age of 2 years with a family history of premature cardiovascular disease or those with at least one parent with high total cholesterol is recommended.

3. Low HDL levels are associated with increased risk for Atherosclerotic Cardiovascular disease (ASCVD) due to insufficient HDL being available

to participate in reverse cholesterol transport, the process by which cholesterol is eliminated from peripheral tissues. 4. NLA-2014 identifies Non HDL Cholesterol (an indicator of all atherogeniclipoproteins such as LDL, VLDL, IDL, Lpa, Chylomicron remnants) along with LDL-cholesterol as co- primary target for cholesterol lowering therapy. Note that major risk factors can modify treatment goals for LDL & Non HDL.

5. Additional testing for Apolipoprotein B, hsCRP,Lp(a) & LP-PLA2 should be considered among patients with moderate risk for ASCVD for risk refinement





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



NAME	: Mrs. KAMINI			
AGE/ GENDER	: 60 YRS/FEMALE		PATIENT ID	: 1606656
COLLECTED BY	:		REG. NO./LAB NO.	: 122411260010
<b>REFERRED BY</b>	:		<b>REGISTRATION DATE</b>	: 26/Nov/2024 10:58 AM
BARCODE NO.	: 12505850		COLLECTION DATE	: 26/Nov/2024 11:28AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITU	JTE	<b>REPORTING DATE</b>	: 26/Nov/2024 01:00PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBA	LA CITY - HA	ARYANA	
Test Name		Value	Unit	<b>Biological Reference interval</b>
	LIVER	FUNCTIO	N TEST (COMPLETE)	
BILIRUBIN TOTAL: by DIAZOTIZATION, SF		0.59	mg/dL	INFANT: 0.20 - 8.00 ADULT: 0.00 - 1.20
	C (CONJUGATED): SERUM	0.12	mg/dL	0.00 - 0.40
,	CT (UNCONJUGATED): SERUM	0.47	mg/dL	0.10 - 1.00
SGOT/AST: SERUM by IFCC, WITHOUT PY	RIDOXAL PHOSPHATE	17.98	U/L	7.00 - 45.00
SGPT/ALT: SERUM		19.41	U/L	0.00 - 49.00
by IFCC, WITHOUT PY AST/ALT RATIO: SI by CALCULATED, SPE		0.93	RATIO	0.00 - 46.00
ALKALINE PHOSPH		106.06	U/L	40.0 - 130.0
GAMMA GLUTAMY	L TRANSFERASE (GGT): SERUM	30.49	U/L	0.00 - 55.0
TOTAL PROTEINS: by BIURET, SPECTRO	SERUM	6.81	gm/dL	6.20 - 8.00
ALBUMIN: SERUM by BROMOCRESOL G	REEN	4.71	gm/dL	3.50 - 5.50
GLOBULIN: SERUM		2.1 <sup>L</sup>	gm/dL	2.30 - 3.50
A : G RATIO: SERUN by CALCULATED, SPE	IN	2.24 <sup>H</sup>	RATIO	1.00 - 2.00

INTERPRETATION

**NOTE:** To be correlated in individuals having SGOT and SGPT values higher than Normal Referance Range. USE: Differential diagnosis of diseases of hepatobiliary system and pancreas.

INCREASED:

DRUG HEPATOTOXICITY	> 2
ALCOHOLIC HEPATITIS	> 2 (Highly Suggestive)
CIRRHOSIS	1.4 - 2.0
INTRAHEPATIC CHOLESTATIS	> 1.5
HEPATOCELLULAR CARCINOMA & CHRONIC HEPATITIS	> 1.3 (Slightly Increased)





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)





A PIONEER DIAGNOSTIC CENTRE

【 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. KAMINI		
AGE/ GENDER	: 60 YRS/FEMALE	PATIENT ID	: 1606656
COLLECTED BY	:	REG. NO./LAB NO.	: 122411260010
<b>REFERRED BY</b>	:	<b>REGISTRATION DATE</b>	: 26/Nov/2024 10:58 AM
BARCODE NO.	: 12505850	<b>COLLECTION DATE</b>	: 26/Nov/2024 11:28AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	<b>REPORTING DATE</b>	: 26/Nov/2024 01:00PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - I	HARYANA	

|--|

### **DECREASED:**

1. Acute Hepatitis due to virus, drugs, toxins (with AST increased 3 to 10 times upper limit of normal)

2. Extra Hepatic cholestatis: 0.8 (normal or slightly decreased).

NORMAL	< 0.65
GOOD PROGNOSTIC SIGN	0.3 - 0.6
POOR PROGNOSTIC SIGN	1.2 - 1.6



**DR.VINAY CHOPRA** CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)





A PIONEER DIAGNOSTIC CENTRE

【 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. KAMINI			
AGE/ GENDER	: 60 YRS/FEMALE		PATIENT ID	: 1606656
COLLECTED BY	:		REG. NO./LAB NO.	: 122411260010
REFERRED BY	:		<b>REGISTRATION DATE</b>	: 26/Nov/2024 10:58 AM
BARCODE NO.	: 12505850		COLLECTION DATE	: 26/Nov/2024 11:28AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTI	TUTE	<b>REPORTING DATE</b>	: 26/Nov/2024 01:00PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMI	BALA CITY - HA	ARYANA	
Test Name		Value	Unit	Biological Reference interva
	KIDNI	EY FUNCTIO	ON TEST (COMPLETE)	1
UREA: SERUM by UREASE - GLUTAM	IATE DEHYDROGENASE (GLDH)	27.87	mg/dL	10.00 - 50.00
CREATININE: SERU		0.87	mg/dL	0.40 - 1.20
BLOOD UREA NITR by CALCULATED, SPE	COGEN (BUN): SERUM	13.02	mg/dL	7.0 - 25.0
BLOOD UREA NITE RATIO: SERUM by CALCULATED, SPE	COGEN (BUN)/CREATININE	14.97	RATIO	10.0 - 20.0
UREA/CREATININ		32.03	RATIO	
URIC ACID: SERUM by URICASE - OXIDAS		5.24	mg/dL	2.50 - 6.80
CALCIUM: SERUM by ARSENAZO III, SPE		10.45	mg/dL	8.50 - 10.60
PHOSPHOROUS: SE by phosphomolybe ELECTROLYTES	ERUM DATE, SPECTROPHOTOMETRY	2.94	mg/dL	2.30 - 4.70
SODIUM: SERUM	'E ELECTRODE)	141	mmol/L	135.0 - 150.0
POTASSIUM: SERUI	M	5	mmol/L	3.50 - 5.00
CHLORIDE: SERUM		105.75	mmol/L	90.0 - 110.0
ESTIMATED GLOM	IERULAR FILTERATION RATE			
ESTIMATED GLOM (eGFR): SERUM by CALCULATED INTERPRETATION:	ERULAR FILTERATION RATE	76.2		
To differentiate betw	een pre- and post renal azotemia.			

INCREASED RATIO (>20:1) WITH NORMAL CREATININE:

1. Prerenal azotemia (BUN rises without increase in creatinine) e.g. heart failure, salt depletion, dehydration, blood loss) due to decreased glomerular filtration rate.

2. Catabolic states with increased tissue breakdown.

3. GI haemorrhage.



**DR.VINAY CHOPRA** CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS , MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST



A PIONEER DIAGNOSTIC CENTRE

🔽 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. KAMINI		
AGE/ GENDER	: 60 YRS/FEMALE	PATIENT ID	: 1606656
COLLECTED BY	:	<b>REG. NO./LAB NO.</b>	: 122411260010
REFERRED BY	:	<b>REGISTRATION DATI</b>	E : 26/Nov/2024 10:58 AM
BARCODE NO.	: 12505850	COLLECTION DATE	: 26/Nov/2024 11:28AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITU	TE <b>REPORTING DATE</b>	: 26/Nov/2024 01:00PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBAL	A CITY - HARYANA	
Test Name		Value Unit	Biological Reference interval
8. Reduced muscle m 9. Certain drugs (e.g. INCREASED RATIO (>2 1. Postrenal azotemia 2. Prerenal azotemia	n (e.g. ureter colostomy) hass (subnormal creatinine production tetracycline, glucocorticoids) 20:1) WITH ELEVATED CREATININE LEVE a (BUN rises disproportionately more t superimposed on renal disease. 10:1) WITH DECREASED BUN :	ELS:	opathy).
1. Acute tubular necr			
2. Low protein diet a			
<ol> <li>Severe liver diseas</li> <li>Other causes of de</li> </ol>	e. ecreased urea synthesis.		
5. Repeated dialysis	(urea rather than creatinine diffuses c		
	monemias (urea is virtually absent in		
	of inappropiate antidiuretic harmone)	due to tubular secretion of urea.	
8. Pregnancy. DECREASED RATIO (~	10:1) WITH INCREASED CREATININE:		
	apy (accelerates conversion of creating	e to creatinine).	
	releases muscle creatinine).	······································	
	who develop renal failure.		
INAPPROPIATE RATIO		- to consistent of the constant of the second state of the second	
Diabatic katoocide	eis (acotoacotato causos falso incroas	o in croatining with cortain mothod	lologios resulting in normal ratio when dehydrat

1. Diabetic ketoacidosis (acetoacetate causes false increase in creatinine with certain methodologies, resulting in normal ratio when dehydration should produce an increased BUN/creatinine ratio).

Cephalosporin therapy (interferes with creatinine measurement).
 ESTIMATED GLOMERULAR FILTERATION RATE:

CKD STAGE	DESCRIPTION	GFR ( mL/min/1.73m2 )	ASSOCIATED FINDINGS
G1	Normal kidney function	>90	No proteinuria
G2	Kidney damage with normal or high GFR	>90	Presence of Protein , Albumin or cast in urine
G3a	Mild decrease in GFR	60 -89	
G3b	Moderate decrease in GFR	30-59	
G4	Severe decrease in GFR	15-29	
G5	Kidney failure	<15	



**DR.VINAY CHOPRA** CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS , MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST





A PIONEER DIAGNOSTIC CENTRE

0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. KAMINI		
AGE/ GENDER	: 60 YRS/FEMALE	PATIENT ID	: 1606656
COLLECTED BY	:	REG. NO./LAB NO.	: 122411260010
<b>REFERRED BY</b>	:	<b>REGISTRATION DATE</b>	: 26/Nov/2024 10:58 AM
BARCODE NO.	: 12505850	<b>COLLECTION DATE</b>	: 26/Nov/2024 11:28AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	<b>REPORTING DATE</b>	: 26/Nov/2024 01:00PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - H	IARYANA	

Test Name	Value	Unit	<b>Biological Reference interval</b>

COMMENTS:

1. Estimated Glomerular filtration rate (eGFR) is the sum of filtration rates in all functioning nephrons and so an estimation of the GFR provides a measure of functioning nephrons of the kidney. 2. eGFR calculated using the 2009 CKD-EPI creatinine equation and GFR category reported as per KDIGO guideline 2012

3. In patients, with eGFR creatinine between 45-59 ml/min/1.73 m2 (G3) and without any marker of Kidney damage, It is recommended to measure eGFR with Cystatin C for confirmation of CKD

4. eGFR category G1 OR G2 does not fullfill the criteria for CKD, in the absence of evidence of Kidney Damage 5. In a suspected case of Acute Kidney Injury (AKI), measurement of eGFR should be done after 48-96 hours of any Intervention or procedure 6. eGFR calculated by Serum Creatinine may be less accurate due to certain factors like Race, Muscle Mass, Diet, Certain Drugs. In such cases, eGFR should be calculated using Serum Cystatin C 7. A decrease in eGFR implies either progressive renal disease, or a reversible process causing decreased nephron function (eg, severe dehydration).

ADVICE:

KDIGO guideline, 2012 recommends Chronic Kidney Disease (CKD) should be classified based on cause, eGFR category and Albuminuria (ACR) category. GFR & ACR category combined together reflect risk of progression and helps Clinician to identify the individual who are progressing at more rapid rate than anticipated



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)





A PIONEER DIAGNOSTIC CENTRE

🔽 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. KAMINI			
AGE/ GENDER	: 60 YRS/FEMALE		PATIENT ID	: 1606656
COLLECTED BY	:		REG. NO./LAB NO.	: 122411260010
REFERRED BY	:		<b>REGISTRATION DATE</b>	: 26/Nov/2024 10:58 AM
BARCODE NO.	: 12505850		<b>COLLECTION DATE</b>	: 26/Nov/2024 11:28AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTIT	ΓUTE	<b>REPORTING DATE</b>	: 26/Nov/2024 04:44PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMB	ALA CITY - H	ARYANA	
Test Name		Value	Unit	Biological Reference interval
		IRON	N PROFILE	
IRON: SERUM by FERROZINE, SPEC	CTROPHOTOMETRY	89.3	µg/dL	37.0 - 145.0
THE ADDRESS AND ADDRESS ADDRES	ALL DIVERSION OF ALL OF			

UNSATURATED IRON BINDING CAPACITY (UIBC)	236.47	µg/dL	150.0 - 336.0
:SERUM			
by FERROZINE, SPECTROPHOTOMETERY			
TOTAL IRON BINDING CAPACITY (TIBC)	325.77	µg/dL	230 - 430
:SERUM			
by SPECTROPHOTOMETERY			
%TRANSFERRIN SATURATION: SERUM	27.41	%	15.0 - 50.0
by CALCULATED, SPECTROPHOTOMETERY (FERENE)			
TRANSFERRIN: SERUM	231.3	mg/dL	200.0 - 350.0
by SPECTROPHOTOMETERY (FERENE)		0	
INTERPRETATION:-			

VARIABLES	ANEMIA OF CHRONIC DISEASE	IRON DEFICIENCY ANEMIA	THALASSEMIA α/β TRAIT
SERUM IRON:	Normal to Reduced	Reduced	Normal
TOTAL IRON BINDING CAPACITY:	Decreased	Increased	Normal
% TRANSFERRIN SATURATION:	Decreased	Decreased < 12-15 %	Normal
SERUM FERRITIN:	Normal to Increased	Decreased	Normal or Increased
IDON:			

#### IRON

1.Serum iron studies is recommended for differential diagnosis of microcytic hypochromic anemia.i.e iron deficiency anemia, zinc deficiency anemia, anemia of chronic disease and thalassemia syndromes.

It is essential to isolate iron deficiency anemia from Beta thalassemia syndromes because during iron replacement which is therapeutic for iron deficiency anemia, is severely contra-indicated in Thalassemia.
 TOTAL IRON BINDING CAPACITY (TIBC):

1.It is a direct measure of protein transferrin which transports iron from the gut to storage sites in the bone marrow.

## % TRANSFERRIN SATURATION:

1. Occurs in idiopathic hemochromatosis and transfusional hemosiderosis where no unsaturated iron binding capacity is available for iron mobilization. Similar condition is seen in congenital deficiency of transferrin.



**DR.VINAY CHOPRA** CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



🔽 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. KAMINI			
AGE/ GENDER	: 60 YRS/FEMALE	PAT	IENT ID	: 1606656
COLLECTED BY	:	REG.	. NO./LAB NO.	: 122411260010
<b>REFERRED BY</b>	:	REG	ISTRATION DATE	: 26/Nov/2024 10:58 AM
BARCODE NO.	: 12505850	COL	LECTION DATE	: 26/Nov/2024 11:28AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITU	TE <b>REP</b> (	ORTING DATE	: 26/Nov/2024 01:28PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBAL	A CITY - HARYAN	JA	
Test Name		Value	Unit	Biological Reference interval
		ENDOCDIN	OLOCY	
		ENDOCRIN		
	THYRO		OLOGY N TEST: TOTAL	
				0.35 - 1.93
THYROXINE (T4): S	NE (T3): SERUM ESCENT MICROPARTICLE IMMUNOASSAY)	DID FUNCTIO	N TEST: TOTAL	0.35 - 1.93 4.87 - 12.60
by CMIA (CHEMILUMINI THYROXINE (T4): S by CMIA (CHEMILUMINI THYROID STIMULA	NE (T3): SERUM ESCENT MICROPARTICLE IMMUNOASSAY) ERUM ESCENT MICROPARTICLE IMMUNOASSAY) TING HORMONE (TSH): SERUM ESCENT MICROPARTICLE IMMUNOASSAY)	DID FUNCTIO 1.29	<b>N TEST: TOTAL</b> ng/mL	

TSH levels are subject to circadian variation, reaching peak levels between 2-4 a.m and at a minimum between 6-10 pm. The variation is of the order of 50%. Hence time of the day has influence on the measured serum TSH concentrations. TSH stimulates the production and secretion of the metabolically active hormones, thyroxine (T4) and triiodothyronine (T3). Failure at any level of regulation of the hypothalamic-pituitary-thyroid axis will result in either underproduction (hypothyroidism) or overproduction(hyperthyroidism) of T4 and/or T3.

CLINICAL CONDITION	T3	T4	TSH
Primary Hypothyroidism:	Reduced	Reduced	Increased (Significantly)
Subclinical Hypothyroidism:	Normal or Low Normal	Normal or Low Normal	High
Primary Hyperthyroidism:	Increased	Increased	Reduced (at times undetectable)
Subclinical Hyperthyroidism:	Normal or High Normal	Normal or High Normal	Reduced

#### LIMITATIONS:-

1. T3 and T4 circulates in reversibly bound form with Thyroid binding globulins (TBG), and to a lesser extent albumin and Thyroid binding Pre Albumin so conditions in which TBG and protein levels alter such as pregnancy, excess estrogens, androgens, anabolic steroids and glucocorticoids may falsely affect the T3 and T4 levels and may cause false thyroid values for thyroid function tests.

2. Normal levels of T4 can also be seen in Hyperthyroid patients with :T3 Thyrotoxicosis, Decreased binding capacity due to hypoproteinemia or ingestion of certain drugs (e.g.: phenytoin , salicylates).

3. Serum T4 levels in neonates and infants are higher than values in the normal adult , due to the increased concentration of TBG in neonate serum.

4. TSH may be normal in central hypothyroidism , recent rapid correction of hyperthyroidism or hypothyroidism , pregnancy , phenytoin therapy.

TRIIODOTHYRONINE (T3)		THYROX	INE (T4)	THYROID STIMULATING HORMONE (TSH)	
Age	Refferance Range (ng/mL)	Age	Refferance Range (µg/dL)	Age	Reference Range ( μIU/mL)
0-7 Days	0.20 - 2.65	0 - 7 Days	5.90 - 18.58	0 - 7 Days	2.43 - 24.3
7 Days - 3 Months	0.36 - 2.59	7 Days - 3 Months	6.39 - 17.66	7 Days - 3 Months	0.58 - 11.00
3 - 6 Months	0.51 - 2.52	3 - 6 Months	6.75 - 17.04	3 Days – 6 Months	0.70 - 8.40
6 - 12 Months	0.74 - 2.40	6 - 12 Months	7.10 - 16.16	6 – 12 Months	0.70 - 7.00





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)





NAME	: Mrs. KAMINI		
AGE/ GENDER	: 60 YRS/FEMALE	PATIENT ID	: 1606656
COLLECTED BY	:	<b>REG. NO./LAB NO.</b>	: 122411260010
<b>REFERRED BY</b>	:	<b>REGISTRATION DATE</b>	: 26/Nov/2024 10:58 AM
BARCODE NO.	: 12505850	<b>COLLECTION DATE</b>	: 26/Nov/2024 11:28AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	<b>REPORTING DATE</b>	: 26/Nov/2024 01:28PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - H	IARYANA	

Test Name			Value	Unit		Biological Reference interval
1 - 10 Years	0.92 - 2.28	1 - 10 Years	6.00 - 13.80	1 – 10 Years	0.60 - 5.50	
11- 19 Years	0.35 - 1.93	11 - 19 Years	4.87-13.20	11 – 19 Years	0.50 - 5.50	
> 20 years (Adults)	0.35 - 1.93	> 20 Years (Adults)	4.87 - 12.60	> 20 Years (Adults)	0.35-5.50	
	RECOM	MENDATIONS OF TSH LE	VELS DURING PREC	GNANCY ( µIU/mL)		
	1st Trimester			0.10 - 2.50		
	2nd Trimester			0.20 - 3.00		
	3rd Trimester			0.30 - 4.10		

### **INCREASED TSH LEVELS:**

1. Primary or untreated hypothyroidism may vary from 3 times to more than 100 times normal depending upon degree of hypofunction.

2. Hypothyroid patients receiving insufficient thyroid replacement therapy.

3.Hashimotos thyroiditis

4.DRUGS: Amphetamines, iodine containing agents & dopamine antagonist.

5.Neonatal period, increase in 1st 2-3 days of life due to post-natal surge

## DECREASED TSH LEVELS:

1.Toxic multi-nodular goiter & Thyroiditis.

2. Over replacement of thyroid hormone in treatment of hypothyroidism.

3. Autonomously functioning Thyroid adenoma

4.Secondary pituitary or hypothalamic hypothyroidism

5. Acute psychiatric illness

6.Severe dehydration.

7.DRUGS: Glucocorticoids, Dopamine, Levodopa, T4 replacement therapy, Anti-thyroid drugs for thyrotoxicosis.

8. Pregnancy: 1st and 2nd Trimester





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)



A PIONEER DIAGNOSTIC CENTRE

🕻 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. KAMINI			
AGE/ GENDER	: 60 YRS/FEMALE	РАТ	TENT ID	: 1606656
COLLECTED BY	:	REG	. NO./LAB NO.	: 122411260010
REFERRED BY	:	REG	ISTRATION DATE	: 26/Nov/2024 10:58 AM
BARCODE NO.	: 12505850	COL	LECTION DATE	: 26/Nov/2024 11:28AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INST	ITUTE <b>REP</b>	ORTING DATE	: 26/Nov/2024 04:44PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AM	BALA CITY - HARYAI	NA	
Test Name		Value	Unit	Biological Reference interval
	IMM	UNOPATHOLO	GY/SEROLOGY	Y
	(	C-REACTIVE PRO	OTEIN (CRP)	
C-REACTIVE PROT SERUM by NEPHLOMETRY	EIN (CRP) QUANTITATIVE:	0.96	mg/L	0.0 - 6.0
INTERPRETATION:			ts for inflammation.	

4. As compared to ESR, CRP shows an earlier rise in inflammatory disorders which begins in 4-6 hrs, the intensity of the rise being higher than ESR and the recovery being earlier than ESR. Unlike ESR, CRP levels are not influenced by hematologic conditions like Anemia, Polycythemia etc., 5. Elevated values are consistent with an acute inflammatory process. NOTE:

1. Elevated C-reactive protein (CRP) values are nonspecific and should not be interpreted without a complete clinical history.

2. Oral contraceptives may increase CRP levels.





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)



A PIONEER DIAGNOSTIC CENTRE

0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. KAMINI					
AGE/ GENDER	: 60 YRS/FEMALE	PA	FIENT ID	: 1606656		
COLLECTED BY	:	REG	G. NO./LAB NO.	: 1224112	60010	
REFERRED BY	:	REG	GISTRATION DATE	:26/Nov/2	024 10:58 AM	
BARCODE NO.	: 12505850	COL	LECTION DATE	:26/Nov/2	024 11:28AM	
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INST	ITUTE <b>REI</b>	PORTING DATE	:26/Nov/2	: 26/Nov/2024 01:29PM	
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMI	BALA CITY - HARYA	NA			
Fest Name		Value	Unit	Bi	ological Reference interva	
Test Name	VITAM	VITAM			ological Reference interva	
VITAMIN D (25-HY	<b>VITAM</b> DROXY VITAMIN D3): SERUM ESCENCE IMMUNOASSAY)	VITAM	IINS	B DI IN SU	ological Reference interva EFICIENCY: < 20.0 ISUFFICIENCY: 20.0 - 30.0 JFFICIENCY: 30.0 - 100.0 DXICITY: > 100.0	
VITAMIN D (25-HY by CLIA (CHEMILUMIN INTERPRETATION:	DROXY VITAMIN D3): SERUM escence immunoassay)	VITAM IIN D/25 HYDE 12.8 <sup>L</sup>	IINS COXY VITAMIN D3 ng/mL	B IN SU TO	EFICIENCY: < 20.0 ISUFFICIENCY: 20.0 - 30.0 JFFICIENCY: 30.0 - 100.0	
VITAMIN D (25-HY by CLIA (CHEMILUMIN INTERPRETATION: DEFI	DROXY VITAMIN D3): SERUM ESCENCE IMMUNOASSAY)	<b>VITAM</b> IIN D/25 HYDF 12.8 <sup>L</sup>	IINS COXY VITAMIN D3 ng/mL	B DI IN SU TO	EFICIENCY: < 20.0 ISUFFICIENCY: 20.0 - 30.0 JFFICIENCY: 30.0 - 100.0	
by CLIA (CHEMILUMIN <u>INTERPRETATION:</u> DEFI INSUFI	DROXY VITAMIN D3): SERUM escence immunoassay)	VITAM IIN D/25 HYDE 12.8 <sup>L</sup>	UNS COXY VITAMIN D3 ng/mL	B IN SU TO	EFICIENCY: < 20.0 ISUFFICIENCY: 20.0 - 30.0 JFFICIENCY: 30.0 - 100.0	

conversion of 7- dihydrocholecalciferol to Vitamin D3 in the skin upon Ultraviolet exposure. 2.25-OH--Vitamin D represents the main body resevoir and transport form of Vitamin D and transport form of Vitamin D, being stored in adipose tissue and tightly bound by a transport protein while in circulation.

3. Vitamin D plays a primary role in the maintenance of calcium homeostatis. It promotes calcium absorption, renal calcium absorption and phosphate reabsorption, skeletal calcium deposition, calcium mobilization, mainly regulated by parathyroid harmone (PTH).

4. Severe deficiency may lead to failure to mineralize newly formed osteoid in bone, resulting in rickets in children and osteomalacia in adults. DECREASED:

1.Lack of sunshine exposure.

2.Inadequate intake, malabsorption (celiac disease)

3. Depressed Hepatic Vitamin D 25- hydroxylase activity

4. Secondary to advanced Liver disease

5. Osteoporosis and Secondary Hyperparathroidism (Mild to Moderate deficiency)

6.Enzyme Inducing drugs: anti-epileptic drugs like phenytoin, phenobarbital and carbamazepine, that increases Vitamin D metabolism.

INCREASED:

1. Hypervitaminosis D is Rare, and is seen only after prolonged exposure to extremely high doses of Vitamin D. When it occurs, it can result in severe hypercalcemia and hyperphophatemia.

CAUTION: Replacement therapy in deficient individuals must be monitored by periodic assessment of Vitamin D levels in order to prevent hypervitaminosis D

NOTE:-Dark coloured individuals as compare to whites, is at higher risk of developing Vitamin D deficiency due to excess of melanin pigment which interefere with Vitamin D absorption.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



【 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. KAMINI			
AGE/ GENDER	: 60 YRS/FEMALE	PA	ATIENT ID	: 1606656
COLLECTED BY	:	R	EG. NO./LAB NO.	: 122411260010
REFERRED BY	:	R	EGISTRATION DATE	: 26/Nov/2024 10:58 AM
BARCODE NO.	: 12505850	CO	DLLECTION DATE	: 26/Nov/2024 11:28AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE IN	STITUTE <b>R</b> I	EPORTING DATE	: 26/Nov/2024 01:00PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, A	MBALA CITY - HARY	ANA	
Test Name		Value	Unit	Biological Reference interval
		CLINICAL P	ATHOLOGY	
	URINE RO	DUTINE & MICR	OSCOPIC EXAMINA	ATION
PHYSICAL EXAMI	NATION			
QUANTITY RECIEV		20	ml	
COLOUR	TANCE SPECTROPHOTOMETRY	PALE YELL	)W	PALE YELLOW
by DIP STICK/REFLEC	TANCE SPECTROPHOTOMETRY			
TRANSPARANCY	TANCE SPECTROPHOTOMETRY	HAZY		CLEAR
SPECIFIC GRAVITY		1.02 PK		1.002 - 1.030
,	TANCE SPECTROPHOTOMETRY			
CHEMICAL EXAMI	<u>NATION</u>			
REACTION	TANCE SPECTROPHOTOMETRY	ACIDIC		
PROTEIN		NEGATIVE (	(-ve)	NEGATIVE (-ve)
•	TANCE SPECTROPHOTOMETRY			
SUGAR by DIP STICK/REFLEC	TANCE SPECTROPHOTOMETRY	NEGATIVE (	-ve)	NEGATIVE (-ve)
pH		5.5		5.0 - 7.5
by DIP STICK/REFLEC BILIRUBIN	TANCE SPECTROPHOTOMETRY	NECATIVE		NEGATIVE (-ve)
	TANCE SPECTROPHOTOMETRY	NEGATIVE (	-ve)	NEGATIVE (-ve)
NITRITE		NEGATIVE (	(-ve)	NEGATIVE (-ve)
UROBILINOGEN	TANCE SPECTROPHOTOMETRY.	NOT DETEC	TED EU/dL	0.2 - 1.0
	TANCE SPECTROPHOTOMETRY			
KETONE BODIES	TANCE SPECTROPHOTOMETRY	NEGATIVE (	(-ve)	NEGATIVE (-ve)
BLOOD		NEGATIVE (	-ve)	NEGATIVE (-ve)
•	TANCE SPECTROPHOTOMETRY			
ASCORBIC ACID by DIP STICK/REFLEC	TANCE SPECTROPHOTOMETRY	NEGATIVE (	-ve)	NEGATIVE (-ve)
MICROSCOPIC EX				
	(RBCs)	NEGATIVE (	-ve) /HPF	0 - 3



**DR.VINAY CHOPRA** CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS , MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST

**NOT VALID FOR MEDICO LEGAL PURPOSE** 



【 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mrs. KAMINI				
AGE/ GENDER	: 60 YRS/FEMALE		PATIENT ID	: 1606656	
<b>COLLECTED BY</b>	COLLECTED BY :		<b>REG. NO./LAB NO.</b>	: 122411260010	
<b>REFERRED BY</b>	:	<b>REGISTRATION DATE</b>		: 26/Nov/2024 10:58 AM : 26/Nov/2024 11:28AM : 26/Nov/2024 01:00PM	
<b>BARCODE NO.</b> : 12505850			<b>COLLECTION DATE</b>		
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTI	<b>X.R JAIN HEALTHCARE INSTITUTE</b>			
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY		IARYANA		
Test Name		Value	Unit	<b>Biological Reference interval</b>	
by MICROSCOPY ON	CENTRIFUGED URINARY SEDIMENT				
PUS CELLS by MICROSCOPY ON	CENTRIFUGED URINARY SEDIMENT	5-7	/HPF	0 - 5	

by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT				
EPITHELIAL CELLS	6-8	/HPF	ABSENT	
by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT				
CRYSTALS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	NEGATIVE (-ve)		NEGATIVE (-ve)	
CASTS	NEGATIVE (-ve)		NEGATIVE (-ve)	
by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT				
BACTERIA	NEGATIVE (-ve)		NEGATIVE (-ve)	
by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT				
OTHERS	NEGATIVE (-ve)		NEGATIVE (-ve)	
by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT				
TRICHOMONAS VAGINALIS (PROTOZOA)	ABSENT		ABSENT	
by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT				

\*\*\* End Of Report



**DR.VINAY CHOPRA** CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS , MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST

