

A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

NAME : Mrs. KAMALJEET KAUR

AGE/ GENDER : 53 YRS/FEMALE **PATIENT ID** : 1682518

COLLECTED BY REG. NO./LAB NO. : 122411260012

REFERRED BY **REGISTRATION DATE** : 26/Nov/2024 11:46 AM BARCODE NO. : 12505852 **COLLECTION DATE** : 26/Nov/2024 11:49AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 26/Nov/2024 01:01PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Value Unit **Biological Reference interval Test Name**

SWASTHYA WELLNESS PANEL: GT **COMPLETE BLOOD COUNT (CBC)**

RED BLOOD CELLS (RBCS) COUNT AND INDICES

13.6	gm/dL	12.0 - 16.0
4.47	Millions/cmm	3.50 - 5.00
40.5	%	37.0 - 50.0
90.5	fL	80.0 - 100.0
30.5	pg	27.0 - 34.0
33.6	g/dL	32.0 - 36.0
13.9	%	11.00 - 16.00
46.5	fL	35.0 - 56.0
20.25	RATIO	BETA THALASSEMIA TRAIT: < 13.0 IRON DEFICIENCY ANEMIA: >13.0
28.21	RATIO	BETA THALASSEMIA TRAIT:<= 65.0 IRON DEFICIENCY ANEMIA: > 65.0
8120	/cmm	4000 - 11000
64	%	50 - 70
28	%	20 - 40
	4.47 40.5 90.5 30.5 33.6 13.9 46.5 20.25 28.21	4.47 Millions/cmm 40.5 % 90.5 fL 30.5 pg 33.6 g/dL 13.9 % 46.5 fL 20.25 RATIO 8120 /cmm 64 %



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)







A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

NAME : Mrs. KAMALJEET KAUR

AGE/ GENDER : 53 YRS/FEMALE **PATIENT ID** : 1682518

COLLECTED BY REG. NO./LAB NO. : 122411260012

REFERRED BY **REGISTRATION DATE** : 26/Nov/2024 11:46 AM BARCODE NO. : 12505852 **COLLECTION DATE** : 26/Nov/2024 11:49AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 26/Nov/2024 01:01PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name	Value	Unit	Biological Reference interval
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY			
EOSINOPHILS	3	%	1 - 6
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY			
MONOCYTES by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	5	%	2 - 12
BASOPHILS by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	0	%	0 - 1
ABSOLUTE LEUKOCYTES (WBC) COUNT			
ABSOLUTE NEUTROPHIL COUNT by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	5197	/cmm	2000 - 7500
ABSOLUTE LYMPHOCYTE COUNT by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	2274 ^L	/cmm	800 - 4900
ABSOLUTE EOSINOPHIL COUNT by flow cytometry by sf cube & microscopy	244	/cmm	40 - 440
ABSOLUTE MONOCYTE COUNT by flow cytometry by sf cube & microscopy	406	/cmm	80 - 880
ABSOLUTE BASOPHIL COUNT by flow cytometry by sf cube & microscopy	0	/cmm	0 - 110
PLATELETS AND OTHER PLATELET PREDICTIVE	MARKERS.		
PLATELET COUNT (PLT) by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	443000	/cmm	150000 - 450000
PLATELETCRIT (PCT) by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	0.4 ^H	%	0.10 - 0.36
MEAN PLATELET VOLUME (MPV) by hydro dynamic focusing, electrical impedence	9	fL	6.50 - 12.0
PLATELET LARGE CELL COUNT (P-LCC) by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	83000	/cmm	30000 - 90000
PLATELET LARGE CELL RATIO (P-LCR) by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	18.8	%	11.0 - 45.0
PLATELET DISTRIBUTION WIDTH (PDW) by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE NOTE: TEST CONDUCTED ON EDTA WHOLE BLOOD	15.6	%	15.0 - 17.0



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS, MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600.



A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 pkrjainhealthcare@gmail.com

NAME : Mrs. KAMALJEET KAUR

AGE/ GENDER : 53 YRS/FEMALE **PATIENT ID** : 1682518

COLLECTED BY REG. NO./LAB NO. : 122411260012

REFERRED BY **REGISTRATION DATE** : 26/Nov/2024 11:46 AM BARCODE NO. : 12505852 **COLLECTION DATE** : 26/Nov/2024 11:49AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 26/Nov/2024 04:49PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Value Unit **Biological Reference interval Test Name**

GLYCOSYLATED HAEMOGLOBIN (HBA1C)

5.5 % GLYCOSYLATED HAEMOGLOBIN (HbA1c): 4.0 - 6.4

WHOLE BLOOD

by HPLC (HIGH PERFORMANCE LIQUID CHROMATOGRAPHY)

ESTIMATED AVERAGE PLASMA GLUCOSE mg/dL 60.00 - 140.00 111.15

by HPLC (HIGH PERFORMANCE LIQUID CHROMATOGRAPHY)

INTERPRETATION:

AS PER AMERICAN DIABETES ASSOCIATION (ADA):			
REFERENCE GROUP GLYCOSYLATED HEMOGLOGIB (HBAIC) in %		OGIB (HBAIC) in %	
Non diabetic Adults >= 18 years	<5.7		
At Risk (Prediabetes)	5.7 – 6.4		
Diagnosing Diabetes	>= 6.5		
9 9	Age > 19 Ye	ears	
	Goals of Therapy:	< 7.0	
Therapeutic goals for glycemic control	Actions Suggested:	>8.0	
	Age < 19 Ye	ears	
	Goal of therapy:	<7.5	

COMMENTS:

- 1. Glycosylated hemoglobin (HbA1c) test is three monthly monitoring done to assess compliace with therapeutic regimen in diabetic patients. 2. Since Hb1c reflects long term fluctuations in blood glucose concentration, a diabetic patient who has recently under good control may still have high concentration of HbAlc. Converse is true for a diabetic previously under good control but now poorly controlled.
- 3. Target goals of < 7.0 % may be beneficial in patients with short duration of diabetes, long life expectancy and no significant cardiovascular disease. In patients with significant complications of diabetes, limited life expectancy or extensive co-morbid conditions, targetting a goal of < 7.0% may not be
- 4.High HbA1c (>9.0 -9.5 %) is strongly associated with risk of development and rapid progression of microvascular and nerve complications 5. Any condition that shorten RBC life span like acute blood loss, hemolytic anemia falsely lower HbA1c results.
- 6.HbA1c results from patients with HbSS,HbSC and HbD must be interpreted with caution, given the pathological processes including anemia, increased red cell turnover, and transfusion requirement that adversely impact HbA1c as a marker of long-term gycemic control.
- 7. Specimens from patients with polycythemia or post-splenctomy may exhibit increse in HbA1c values due to a somewhat longer life span of the red cells.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)





A PIONEER DIAGNOSTIC CENTRE

NAME : Mrs. KAMALJEET KAUR

: 1682518 AGE/ GENDER : 53 YRS/FEMALE **PATIENT ID**

COLLECTED BY REG. NO./LAB NO. : 122411260012

REFERRED BY **REGISTRATION DATE** : 26/Nov/2024 11:46 AM BARCODE NO. **COLLECTION DATE** : 26/Nov/2024 11:49AM : 12505852 CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 26/Nov/2024 04:08PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Value Unit **Test Name Biological Reference interval**

ERYTHROCYTE SEDIMENTATION RATE (ESR)

ERYTHROCYTE SEDIMENTATION RATE (ESR)

mm/1st hr 0 - 20 39H

by RED CELL AGGREGATION BY CAPILLARY PHOTOMETRY

INTERPRETATION:

- 1. ESR is a non-specific test because an elevated result often indicates the presence of inflammation associated with infection, cancer and autoimmune disease, but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it.
- 2. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other test such as C-reactive protein
- 3. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as some others, such as systemic lupus erythematosus

CONDITION WITH LOW ESR

A low ESR can be seen with conditions that inhibit the normal sedimentation of red blood cells, such as a high red blood cell count (polycythaemia), significantly high white blood cell count (leucocytosis), and some protein abnormalities. Some changes in red cell shape (such as sickle cells in sickle cell anaemia) also lower the ESR.

NOTE:

- 1. ESR and C reactive protein (C-RP) are both markers of inflammation.
- 2. Generally, ESR does not change as rapidly as does CRP, either at the start of inflammation or as it resolves.
 3. CRP is not affected by as many other factors as is ESR, making it a better marker of inflammation.
 4. If the ESR is elevated, it is typically a result of two types of proteins, globulins or fibringen.
 5. Women tend to average mathyldone and entraceptives professional processing mathyldone and with the opposition of the oppositio

- 6. Drugs such as dextran, methyldopa, oral contraceptives, penicillamine procainamide, theophylline, and vitamin A can increase ESR, while aspirin, cortisone, and quinine may decrease it



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)







A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 pkrjainhealthcare@gmail.com

NAME : Mrs. KAMALJEET KAUR

AGE/ GENDER : 53 YRS/FEMALE **PATIENT ID** : 1682518

COLLECTED BY REG. NO./LAB NO. : 122411260012

REFERRED BY **REGISTRATION DATE** : 26/Nov/2024 11:46 AM BARCODE NO. : 12505852 **COLLECTION DATE** : 26/Nov/2024 11:49AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 26/Nov/2024 01:01PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Value Unit **Biological Reference interval Test Name**

CLINICAL CHEMISTRY/BIOCHEMISTRY GLUCOSE FASTING (F)

GLUCOSE FASTING (F): PLASMA 86.87 NORMAL: < 100.0 mg/dL

by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD) PREDIABETIC: 100.0 - 125.0

DIABETIC: > 0R = 126.0

INTERPRETATION
IN ACCORDANCE WITH AMERICAN DIABETES ASSOCIATION GUIDELINES:

1. A fasting plasma glucose level below 100 mg/dl is considered normal.

2. A fasting plasma glucose level between 100 - 125 mg/dl is considered as glucose intolerant or prediabetic. A fasting and post-prandial blood

test (after consumption of 75 gms of glucose) is recommended for all such patients.

3. A fasting plasma glucose level of above 125 mg/dl is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for all such patients. A fasting plasma glucose level in excess of 125 mg/dl on both occasions is confirmatory for diabetic state.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)



CLIENT CODE.



PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

REPORTING DATE

■ 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

: 26/Nov/2024 01:01PM

NAME : Mrs. KAMALJEET KAUR

AGE/ GENDER : 53 YRS/FEMALE **PATIENT ID** : 1682518

COLLECTED BY REG. NO./LAB NO. : 122411260012

REFERRED BY **REGISTRATION DATE** : 26/Nov/2024 11:46 AM BARCODE NO. : 12505852 **COLLECTION DATE** : 26/Nov/2024 11:49AM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

Test Name	Value	Unit	Biological Reference interval
	LIPID PROFILE	: BASIC	
CHOLESTEROL TOTAL: SERUM by CHOLESTEROL OXIDASE PAP	192.17	mg/dL	OPTIMAL: < 200.0 BORDERLINE HIGH: 200.0 - 239.0 HIGH CHOLESTEROL: > OR = 240.0
TRIGLYCERIDES: SERUM by GLYCEROL PHOSPHATE OXIDASE (ENZYMATIC)	171.51 ^H	mg/dL	OPTIMAL: < 150.0 BORDERLINE HIGH: 150.0 - 199.0 HIGH: 200.0 - 499.0 VERY HIGH: > OR = 500.0
HDL CHOLESTEROL (DIRECT): SERUM by SELECTIVE INHIBITION	60.04	mg/dL	LOW HDL: < 30.0 BORDERLINE HIGH HDL: 30.0 - 60.0 HIGH HDL: > OR = 60.0
LDL CHOLESTEROL: SERUM by CALCULATED, SPECTROPHOTOMETRY	97.83	mg/dL	OPTIMAL: < 100.0 ABOVE OPTIMAL: 100.0 - 129.0 BORDERLINE HIGH: 130.0 - 159.0 HIGH: 160.0 - 189.0 VERY HIGH: > OR = 190.0
NON HDL CHOLESTEROL: SERUM by CALCULATED, SPECTROPHOTOMETRY	132.13 ^H	mg/dL	OPTIMAL: < 130.0 ABOVE OPTIMAL: 130.0 - 159.0 BORDERLINE HIGH: 160.0 - 189.0 HIGH: 190.0 - 219.0 VERY HIGH: > OR = 220.0
VLDL CHOLESTEROL: SERUM by CALCULATED, SPECTROPHOTOMETRY	34.3	mg/dL	0.00 - 45.00
TOTAL LIPIDS: SERUM by CALCULATED, SPECTROPHOTOMETRY	555.85	mg/dL	350.00 - 700.00
CHOLESTEROL/HDL RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	3.2	RATIO	LOW RISK: 3.30 - 4.40 AVERAGE RISK: 4.50 - 7.0 MODERATE RISK: 7.10 - 11.0 HIGH RISK: > 11.0



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)







A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

NAME : Mrs. KAMALJEET KAUR

AGE/ GENDER : 53 YRS/FEMALE **PATIENT ID** : 1682518

COLLECTED BY REG. NO./LAB NO. : 122411260012

REFERRED BY **REGISTRATION DATE** : 26/Nov/2024 11:46 AM BARCODE NO. **COLLECTION DATE** : 26/Nov/2024 11:49AM : 12505852 CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 26/Nov/2024 01:01PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name	Value	Unit	Biological Reference interval
LDL/HDL RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	1.63	RATIO	LOW RISK: 0.50 - 3.0 MODERATE RISK: 3.10 - 6.0 HIGH RISK: > 6.0
TRIGLYCERIDES/HDL RATIO: SERUM	2.86 ^L	RATIO	3.00 - 5.00

INTERPRETATION:

1.Measurements in the same patient can show physiological& analytical variations. Three serial samples 1 week apart are recommended for Total Cholesterol, Triglycerides, HDL & LDL Cholesterol.

2. As per NLA-2014 guidelines, all adults above the age of 20 years should be screened for lipid status. Selective screening of children above the age of 2 years with a family history of premature cardiovascular disease or those with at least one parent with high total cholesterol is recommended.

3. Low HDL levels are associated with increased risk for Atherosclerotic Cardiovascular disease (ASCVD) due to insufficient HDL being available

to participate in reverse cholesterol transport, the process by which cholesterol is eliminated from peripheral tissues.
4. NLA-2014 identifies Non HDL Cholesterol (an indicator of all atherogeniclipoproteins such as LDL, VLDL, IDL, Lpa, Chylomicron remnants) along with LDL-cholesterol as co-primary target for cholesterol lowering therapy. Note that major risk factors can modify treatment goals for LDL &Non

5. Additional testing for Apolipoprotein B, hsCRP,Lp(a) & LP-PLA2 should be considered among patients with moderate risk for ASCVD for risk refinement



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)





A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

NAME : Mrs. KAMALJEET KAUR

AGE/ GENDER : 53 YRS/FEMALE **PATIENT ID** : 1682518

COLLECTED BY REG. NO./LAB NO. : 122411260012

REFERRED BY **REGISTRATION DATE** : 26/Nov/2024 11:46 AM BARCODE NO. : 12505852 **COLLECTION DATE** : 26/Nov/2024 11:49AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 26/Nov/2024 01:01PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name Value Unit **Biological Reference interval**

LIVER FUNCTION TEST (COMPLETE)

BILIRUBIN TOTAL: SERUM by DIAZOTIZATION, SPECTROPHOTOMETRY	0.83	mg/dL	INFANT: 0.20 - 8.00 ADULT: 0.00 - 1.20
BILIRUBIN DIRECT (CONJUGATED): SERUM by DIAZO MODIFIED, SPECTROPHOTOMETRY	0.16	mg/dL	0.00 - 0.40
BILIRUBIN INDIRECT (UNCONJUGATED): SERUM by CALCULATED, SPECTROPHOTOMETRY	0.67	mg/dL	0.10 - 1.00
SGOT/AST: SERUM by IFCC, WITHOUT PYRIDOXAL PHOSPHATE	21.75	U/L	7.00 - 45.00
SGPT/ALT: SERUM by IFCC, WITHOUT PYRIDOXAL PHOSPHATE	18.83	U/L	0.00 - 49.00
AST/ALT RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	1.16	RATIO	0.00 - 46.00
ALKALINE PHOSPHATASE: SERUM by Para nitrophenyl phosphatase by amino methyl propanol	81.16	U/L	40.0 - 130.0
GAMMA GLUTAMYL TRANSFERASE (GGT): SERUM by SZASZ, SPECTROPHTOMETRY	36.75	U/L	0.00 - 55.0
TOTAL PROTEINS: SERUM by BIURET, SPECTROPHOTOMETRY	7.05	gm/dL	6.20 - 8.00
ALBUMIN: SERUM by BROMOCRESOL GREEN	4.66	gm/dL	3.50 - 5.50
GLOBULIN: SERUM by CALCULATED, SPECTROPHOTOMETRY	2.39	gm/dL	2.30 - 3.50
A : G RATIO: SERUM	1.95	RATIO	1.00 - 2.00

INTERPRETATION

by CALCULATED, SPECTROPHOTOMETRY

NOTE:- To be correlated in individuals having SGOT and SGPT values higher than Normal Referance Range.

USE:- Differential diagnosis of diseases of hepatobiliary system and pancreas.

INCREASED:

DRUG HEPATOTOXICITY	> 2
ALCOHOLIC HEPATITIS	> 2 (Highly Suggestive)
CIRRHOSIS	1.4 - 2.0
INTRAHEPATIC CHOLESTATIS	> 1.5
HEPATOCELLULAR CARCINOMA & CHRONIC HEPATITIS	> 1.3 (Slightly Increased)



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST







A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

NAME : Mrs. KAMALJEET KAUR

AGE/ GENDER : 53 YRS/FEMALE **PATIENT ID** : 1682518

COLLECTED BY REG. NO./LAB NO. : 122411260012

REFERRED BY **REGISTRATION DATE** : 26/Nov/2024 11:46 AM BARCODE NO. : 12505852 **COLLECTION DATE** : 26/Nov/2024 11:49AM

CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 26/Nov/2024 01:01PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name Value Unit **Biological Reference interval**

DECREASED:

1. Acute Hepatitis due to virus, drugs, toxins (with AST increased 3 to 10 times upper limit of normal)

2. Extra Hepatic cholestatis: 0.8 (normal or slightly decreased).

PROGNOSTIC SIGNIFICANCE:

NORMAL	< 0.65
GOOD PROGNOSTIC SIGN	0.3 - 0.6
POOR PROGNOSTIC SIGN	1.2 - 1.6



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)

NOT VALID FOR MEDICO LEGAL PURPOSE



PKR JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

2.50 - 6.80

8.50 - 10.60

NAME : Mrs. KAMALJEET KAUR

AGE/ GENDER : 53 YRS/FEMALE **PATIENT ID** : 1682518

COLLECTED BY REG. NO./LAB NO. : 122411260012

REFERRED BY **REGISTRATION DATE** : 26/Nov/2024 11:46 AM : 12505852 BARCODE NO. **COLLECTION DATE** : 26/Nov/2024 11:49AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 26/Nov/2024 01:01PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name	Value	Unit	Biological Reference interval
KIDN	EY FUNCTION T	TEST (COMPLETE)	
UREA: SERUM by UREASE - GLUTAMATE DEHYDROGENASE (GLDH)	20.08	mg/dL	10.00 - 50.00
CREATININE: SERUM by ENZYMATIC, SPECTROPHOTOMETERY	0.75	mg/dL	0.40 - 1.20
BLOOD UREA NITROGEN (BUN): SERUM by CALCULATED, SPECTROPHOTOMETRY	9.38	mg/dL	7.0 - 25.0
BLOOD UREA NITROGEN (BUN)/CREATININE RATIO: SERUM by CALCULATED. SPECTROPHOTOMETRY	12.51	RATIO	10.0 - 20.0
UREA/CREATININE RATIO: SERUM	26.77	RATIO	

by CALCULATED, SPECTROPHOTOMETRY		
URIC ACID: SERUM	4.59	mg/dL
by URICASE - OXIDASE PEROXIDASE		
CALCIUM: SERUM	10.3	mg/dL
by ARSENAZO III, SPECTROPHOTOMETRY		

PHOSPHOROUS: SERUM 2.88 2.30 - 4.70 mg/dL

by PHOSPHOMOLYBDATE, SPECTROPHOTOMETRY

ELECTROLYTES SODIUM: SERUM 138.7 mmol/L 135.0 - 150.0 by ISE (ION SELECTIVE ELECTRODE) POTASSIUM: SERUM mmol/L 3.50 - 5.00by ISE (ION SELECTIVE ELECTRODE) 90.0 - 110.0 CHLORIDE: SERUM 104.03 mmol/L

by ISE (ION SELECTIVE ELECTRODE)

ESTIMATED GLOMERULAR FILTERATION RATE

ESTIMATED GLOMERULAR FILTERATION RATE 95.1

(eGFR): SERUM by CALCULATED

INTERPRETATION: To differentiate between pre- and post renal azotemia.

INCREASED RATIO (>20:1) WITH NORMAL CREATININE:

1. Prerenal azotemia (BUN rises without increase in creatinine) e.g. heart failure, salt depletion, dehydration, blood loss) due to decreased glomerular filtration rate.

Catabolic states with increased tissue breakdown.

3. GI haemorrhage.



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)





A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

NAME : Mrs. KAMALJEET KAUR

: 1682518 AGE/ GENDER : 53 YRS/FEMALE **PATIENT ID**

COLLECTED BY REG. NO./LAB NO. : 122411260012

REFERRED BY **REGISTRATION DATE** : 26/Nov/2024 11:46 AM BARCODE NO. **COLLECTION DATE** : 26/Nov/2024 11:49AM : 12505852 CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 26/Nov/2024 01:01PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name Value Unit **Biological Reference interval**

4. High protein intake.

5. Impaired renal function plus

6. Excess protein intake or production or tissue breakdown (e.g. infection, GI bleeding, thyrotoxicosis, Cushing's syndrome, high protein diet, burns, surgery, cachexia, high fever).

7. Urine reabsorption (e.g. ureter colostomy)

8. Reduced muscle mass (subnormal creatinine production)

9. Certain drugs (e.g. tetracycline, glucocorticoids)

INCREASED RATIO (>20:1) WITH ELEVATED CREATININE LEVELS:

- 1. Postrenal azotemia (BUN rises disproportionately more than creatinine) (e.g. obstructive uropathy).
- 2. Prerenal azotemia superimposed on renal disease.

DECREASED RATIO (<10:1) WITH DECREASED BUN:

- 1. Acute tubular necrosis.
- 2. Low protein diet and starvation.
- 3. Severe liver disease.
- 4. Other causes of decreased urea synthesis.
- 5. Repeated dialysis (urea rather than creatinine diffuses out of extracellular fluid).
- 6. Inherited hyperammonemias (urea is virtually absent in blood).
- 7. SIADH (syndrome of inappropiate antidiuretic harmone) due to tubular secretion of urea.

8. Pregnancy.

DECREASED RATIO (<10:1) WITH INCREASED CREATININE:

- 1. Phenacimide therapy (accelerates conversion of creatine to creatinine).
- 2. Rhabdomyolysis (releases muscle creatinine).
- 3. Muscular patients who develop renal failure.

INAPPROPIATE RATIO:

1. Diabetic ketoacidosis (acetoacetate causes false increase in creatinine with certain methodologies, resulting in normal ratio when dehydration should produce an increased BUN/creatinine ratio).

2. Cephalosporin therapy (interferes with creatinine measurement). **ESTIMATED GLOMERULAR FILTERATION RATE**:

SHIVIATED GEOMEROPIK FIETERATION RATE:			
CKD STAGE	DESCRIPTION	GFR (mL/min/1.73m2)	ASSOCIATED FINDINGS
G1	Normal kidney function	>90	No proteinuria
G2	Kidney damage with	>90	Presence of Protein,
	normal or high GFR		Albumin or cast in urine
G3a	Mild decrease in GFR	60 -89	
G3b	Moderate decrease in GFR	30-59	
G4	Severe decrease in GFR	15-29	
G5	Kidney failure	<15	



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)





A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

NAME : Mrs. KAMALJEET KAUR

: 1682518 AGE/ GENDER : 53 YRS/FEMALE **PATIENT ID**

COLLECTED BY REG. NO./LAB NO. : 122411260012

REFERRED BY **REGISTRATION DATE** : 26/Nov/2024 11:46 AM BARCODE NO. **COLLECTION DATE** : 26/Nov/2024 11:49AM : 12505852 CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 26/Nov/2024 01:01PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name Value Unit **Biological Reference interval**

COMMENTS:

1. Estimated Glomerular filtration rate (eGFR) is the sum of filtration rates in all functioning nephrons and so an estimation of the GFR provides a measure of functioning nephrons of the kidney.

2. eGFR calculated using the 2009 CKD-EPI creatinine equation and GFR category reported as per KDIGO guideline 2012

3. In patients, with eGFR creatinine between 45-59 ml/min/1.73 m2 (G3) and without any marker of Kidney damage, It is recommended to measure eGFR with Cystatin C for confirmation of CKD

4. eGFR category G1 OR G2 does not fullfill the criteria for CKD, in the absence of evidence of Kidney Damage
5. In a suspected case of Acute Kidney Injury (AKI), measurement of eGFR should be done after 48-96 hours of any Intervention or procedure
6. eGFR calculated by Serum Creatinine may be less accurate due to certain factors like Race, Muscle Mass, Diet, Certain Drugs. In such cases, eGFR should be calculated using Serum Cystatin C
7. A decrease in eGFR implies either progressive renal disease, or a reversible process causing decreased nephron function (eg, severe dehydration).

KDIGO guideline, 2012 recommends Chronic Kidney Disease (CKD) should be classified based on cause, eGFR category and Albuminuria (ACR) category. GFR & ACR category combined together reflect risk of progression and helps Clinician to identify the individual who are progressing at more rapid rate than anticipated



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)



CLIENT CODE.





PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

: 26/Nov/2024 01:28PM

NAME : Mrs. KAMALJEET KAUR

AGE/ GENDER : 53 YRS/FEMALE **PATIENT ID** : 1682518

COLLECTED BY REG. NO./LAB NO. : 122411260012

REFERRED BY **REGISTRATION DATE** : 26/Nov/2024 11:46 AM BARCODE NO. : 12505852 **COLLECTION DATE** : 26/Nov/2024 11:49AM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

Value Unit **Biological Reference interval Test Name**

REPORTING DATE

ENDOCRINOLOGY

THYROID FUNCTION TEST: TOTAL

TRIIODOTHYRONINE (T3): SERUM 0.35 - 1.93ng/mL

by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

THYROXINE (T4): SERUM 7.26 μgm/dL 4.87 - 12.60by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

THYROID STIMULATING HORMONE (TSH): SERUM 36.54^H 0.35 - 5.50μIU/mL

by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

3rd GENERATION, ULTRASENSITIVE INTERPRETATION:

TSH levels are subject to circadian variation, reaching peak levels between 2-4 a.m and at a minimum between 6-10 pm. The variation is of the order of 50%. Hence time of the day has influence on the measured serum TSH concentrations. TSH stimulates the production and secretion of the metabolically active hormones, thyroxine (T4) and triliodothyronine (T3). Failure at any level of regulation of the hypothalamic-pituitary-thyroid axis will result in either underproduction (hypothyroidism) or overproduction(hyperthyroidism) of T4 and/or T3.

CLINICAL CONDITION	Т3	T4	TSH
Primary Hypothyroidism:	Reduced	Reduced	Increased (Significantly)
Subclinical Hypothyroidism:	Normal or Low Normal	Normal or Low Normal	High
Primary Hyperthyroidism:	Increased	Increased	Reduced (at times undetectable)
Subclinical Hyperthyroidism:	Normal or High Normal	Normal or High Normal	Reduced

- 1. T3 and T4 circulates in reversibly bound form with Thyroid binding globulins (TBG), and to a lesser extent albumin and Thyroid binding Pre Albumin so conditions in which TBG and protein levels alter such as pregnancy, excess estrogens, androgens, anabolic steroids and glucocorticoids may falsely affect the T3 and T4 levels and may cause false thyroid values for thyroid function tests
- 2. Normal levels of T4 can also be seen in Hyperthyroid patients with :T3 Thyrotoxicosis, Decreased binding capacity due to hypoproteinemia or ingestion of certain drugs
- 3. Serum T4 levels in neonates and infants are higher than values in the normal adult, due to the increased concentration of TBG in neonate serum.
- 4. TSH may be normal in central hypothyroidism, recent rapid correction of hyperthyroidism or hypothyroidism, pregnancy, phenytoin therapy.

TRIIODOTHYRONINE (T3)		THYROXINE (T4)		THYROID STIMULATING HORMONE (TSH)	
Age	Refferance Range (ng/mL)	Age	Refferance Range (µg/dL)	Age	Reference Range (μIU/mL)
0-7 Days	0.20 - 2.65	0 - 7 Days	5.90 - 18.58	0 - 7 Days	2.43 - 24.3
7 Days - 3 Months	0.36 - 2.59	7 Days - 3 Months	6.39 - 17.66	7 Days - 3 Months	0.58 - 11.00
3 - 6 Months	0.51 - 2.52	3 - 6 Months	6.75 – 17.04	3 Days – 6 Months	0.70 - 8.40
6 - 12 Months	0.74 - 2.40	6 - 12 Months	7.10 – 16.16	6 – 12 Months	0.70 - 7.00



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)





A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 pkrjainhealthcare@gmail.com

NAME : Mrs. KAMALJEET KAUR

AGE/ GENDER : 53 YRS/FEMALE **PATIENT ID** : 1682518

COLLECTED BY : 122411260012 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 26/Nov/2024 11:46 AM BARCODE NO. : 12505852 **COLLECTION DATE** : 26/Nov/2024 11:49AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 26/Nov/2024 01:28PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name			Value	Unit		Biological Reference interval
1 - 10 Years	0.92 - 2.28	1 - 10 Years	6.00 - 13.80	1 – 10 Years	0.60 - 5.50	
11- 19 Years	0.35 - 1.93	11 - 19 Years	4.87- 13.20	11 – 19 Years	0.50 - 5.50	
> 20 years (Adults)	0.35 - 1.93	> 20 Years (Adults)	4.87 - 12.60	> 20 Years (Adults)	0.35-5.50	
1st Trimester			0.10 - 2.50			
2nd Trimester			0.20 - 3.00			
3rd Trimester			0.30 - 4.10			

INCREASED TSH LEVELS:

- 1. Primary or untreated hypothyroidism may vary from 3 times to more than 100 times normal depending upon degree of hypofunction.
- 2. Hypothyroid patients receiving insufficient thyroid replacement therapy.
- 3. Hashimotos thyroiditis
- 4.DRUGS: Amphetamines, iodine containing agents & dopamine antagonist.
- 5. Neonatal period, increase in 1st 2-3 days of life due to post-natal surge

DECREASED TSH LEVELS:

- 1. Toxic multi-nodular goiter & Thyroiditis.
- 2. Over replacement of thyroid hormone in treatment of hypothyroidism.
- 3. Autonomously functioning Thyroid adenoma
- 4. Secondary pituitary or hypothalamic hypothyroidism
- 5. Acute psychiatric illness
- 6. Severe dehydration.
- 7.DRUGS: Glucocorticoids, Dopamine, Levodopa, T4 replacement therapy, Anti-thyroid drugs for thyrotoxicosis.
- 8. Pregnancy: 1st and 2nd Trimester

*** End Of Report ***



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)