

# A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

**NAME** : Mr. RAM SINGH

**AGE/ GENDER** : 70 YRS/MALE **PATIENT ID** : 1691245

**COLLECTED BY** REG. NO./LAB NO. : 122412050003

REFERRED BY **REGISTRATION DATE** : 05/Dec/2024 08:31 AM BARCODE NO. : 12506000 **COLLECTION DATE** : 05/Dec/2024 03:10PM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 05/Dec/2024 01:20PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**Value** Unit **Biological Reference interval Test Name** 

# **SWASTHYA WELLNESS PANEL: 1.0 COMPLETE BLOOD COUNT (CBC)**

#### **RED BLOOD CELLS (RBCS) COUNT AND INDICES**

| HAEMOGLOBIN (HB) by CALORIMETRIC  | 15.6              | gm/dL        | 12.0 - 17.0  |
|---|-------------------|--------------|--|
| RED BLOOD CELL (RBC) COUNT by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE              | 5.41 <sup>H</sup> | Millions/cmm | 3.50 - 5.00  |
| PACKED CELL VOLUME (PCV) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER                 | 46.4              | %            | 40.0 - 54.0  |
| MEAN CORPUSCULAR VOLUME (MCV) by calculated by automated hematology analyzer            | 85.8              | fL           | 80.0 - 100.0   |
| MEAN CORPUSCULAR HAEMOGLOBIN (MCH) by calculated by automated hematology analyzer       | 28.9              | pg           | 27.0 - 34.0  |
| MEAN CORPUSCULAR HEMOGLOBIN CONC. (MCHC) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER | 33.7              | g/dL         | 32.0 - 36.0  |
| RED CELL DISTRIBUTION WIDTH (RDW-CV) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER     | 12.8              | %            | 11.00 - 16.00  |
| RED CELL DISTRIBUTION WIDTH (RDW-SD) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER     | 41.8              | fL           | 35.0 - 56.0  |
| MENTZERS INDEX by CALCULATED  | 15.86             | RATIO        | BETA THALASSEMIA TRAIT: < 13.0<br>IRON DEFICIENCY ANEMIA: >13.0  |
| GREEN & KING INDEX by CALCULATED  | 20.35             | RATIO        | BETA THALASSEMIA TRAIT:<= 65.0<br>IRON DEFICIENCY ANEMIA: > 65.0 |
| WHITE BLOOD CELLS (WBCS)  |                   |              |  |
| TOTAL LEUCOCYTE COUNT (TLC) by flow cytometry by sf cube & microscopy                   | 10800             | /cmm         | 4000 - 11000   |
| <u>DIFFERENTIAL LEUCOCYTE COUNT (DLC)</u>   |                   |              |  |
| NEUTROPHILS by flow cytometry by sf cube & microscopy                                   | 67                | %            | 50 - 70  |
| LYMPHOCYTES   | 21                | %            | 20 - 40  |



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST







# A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

NAME : Mr. RAM SINGH

**AGE/ GENDER** : 70 YRS/MALE **PATIENT ID** : 1691245

**COLLECTED BY** REG. NO./LAB NO. : 122412050003

REFERRED BY **REGISTRATION DATE** : 05/Dec/2024 08:31 AM BARCODE NO. : 12506000 **COLLECTION DATE** : 05/Dec/2024 03:10PM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 05/Dec/2024 01:20PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

| Test Name  | Value             | Unit     | Biological Reference interval |
|--|-------------------|----------|-------------------------------|
| by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY                                  |                   |          |                               |
| EOSINOPHILS  | 5                 | %        | 1 - 6                         |
| by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY                                  |                   |          |                               |
| MONOCYTES  | 7                 | %        | 2 - 12                        |
| by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY BASOPHILS                        | 0                 | %        | 0 - 1                         |
| by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY                                  | 0                 | %        | 0 - 1                         |
| ABSOLUTE LEUKOCYTES (WBC) COUNT  |                   |          |                               |
| ABSOLUTE NEUTROPHIL COUNT  | 7236              | /cmm     | 2000 - 7500                   |
| by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY                                  |                   | , 611111 | 2000 1000                     |
| ABSOLUTE LYMPHOCYTE COUNT  | 2268 <sup>L</sup> | /cmm     | 800 - 4900                    |
| by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY                                  |                   |          |                               |
| ABSOLUTE EOSINOPHIL COUNT  | 540 <sup>H</sup>  | /cmm     | 40 - 440                      |
| by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY ABSOLUTE MONOCYTE COUNT          | 750               | /        | 80 - 880                      |
| by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY                                  | 756               | /cmm     | 00 - 00U                      |
| ABSOLUTE BASOPHIL COUNT  | 0                 | /cmm     | 0 - 110                       |
| by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY                                  |                   |          |                               |
| PLATELETS AND OTHER PLATELET PREDICTIVE                                    | MARKERS.          |          |                               |
| PLATELET COUNT (PLT)   | 224000            | /cmm     | 150000 - 450000               |
| by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE                            |                   |          |                               |
| PLATELETCRIT (PCT)   | 0.21              | %        | 0.10 - 0.36                   |
| by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE                            |                   | CT.      | 0.50 10.0                     |
| MEAN PLATELET VOLUME (MPV) by HYDRO DYNAMIC FOCUSING. ELECTRICAL IMPEDENCE | 9                 | fL       | 6.50 - 12.0                   |
| PLATELET LARGE CELL COUNT (P-LCC)  | 52000             | /cmm     | 30000 - 90000                 |
| by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE                            | 32000             | / ciiiii | 30000 30000                   |
| PLATELET LARGE CELL RATIO (P-LCR)  | 23.2              | %        | 11.0 - 45.0                   |
| by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE                            |                   |          |                               |
| PLATELET DISTRIBUTION WIDTH (PDW)  | 16.2              | %        | 15.0 - 17.0                   |
| by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE                            |                   |          |                               |
| NOTE: TEST CONDUCTED ON EDTA WHOLE BLOOD                                   |                   |          |                               |



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS, MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST





# A PIONEER DIAGNOSTIC CENTRE

: 05/Dec/2024 01:20PM

**NAME** : Mr. RAM SINGH

AGE/ GENDER : 70 YRS/MALE **PATIENT ID** : 1691245

**COLLECTED BY** REG. NO./LAB NO. : 122412050003

REFERRED BY **REGISTRATION DATE** : 05/Dec/2024 08:31 AM BARCODE NO. **COLLECTION DATE** : 05/Dec/2024 03:10PM : 12506000

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

**Value** Unit **Test Name Biological Reference interval** 

#### **ERYTHROCYTE SEDIMENTATION RATE (ESR)**

ERYTHROCYTE SEDIMENTATION RATE (ESR)

mm/1st hr 0 - 20

REPORTING DATE

by RED CELL AGGREGATION BY CAPILLARY PHOTOMETRY

#### INTERPRETATION:

CLIENT CODE.

1. ESR is a non-specific test because an elevated result often indicates the presence of inflammation associated with infection, cancer and auto-immune disease, but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it.

2. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other test such as C-reactive protein

3. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as some others, such as systemic lupus erythematosus

#### CONDITION WITH LOW ESR

A low ESR can be seen with conditions that inhibit the normal sedimentation of red blood cells, such as a high red blood cell count (polycythaemia), significantly high white blood cell count (leucocytosis), and some protein abnormalities. Some changes in red cell shape (such as sickle cells in sickle cell anaemia) also lower the ESR.

NOTE:

- 1. ESR and C reactive protein (C-RP) are both markers of inflammation.
- 2. Generally, ESR does not change as rapidly as does CRP, either at the start of inflammation or as it resolves.
   3. CRP is not affected by as many other factors as is ESR, making it a better marker of inflammation.
   4. If the ESR is elevated, it is typically a result of two types of proteins, globulins or fibringen.
   5. Women tend to average mathyldone and entraceptives professional processing mathyldone and with the opposition of the oppositio

- 6. Drugs such as dextran, methyldopa, oral contraceptives, penicillamine procainamide, theophylline, and vitamin A can increase ESR, while aspirin, cortisone, and quinine may decrease it



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)





# A PIONEER DIAGNOSTIC CENTRE

**NAME** : Mr. RAM SINGH

AGE/ GENDER : 70 YRS/MALE **PATIENT ID** : 1691245

**COLLECTED BY** REG. NO./LAB NO. : 122412050003

REFERRED BY **REGISTRATION DATE** : 05/Dec/2024 08:31 AM BARCODE NO. : 12506000 **COLLECTION DATE** : 05/Dec/2024 03:10PM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 05/Dec/2024 01:20PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**Value** Unit **Biological Reference interval Test Name** 

# **CLINICAL CHEMISTRY/BIOCHEMISTRY GLUCOSE FASTING (F)**

GLUCOSE FASTING (F): PLASMA 81.75 NORMAL: < 100.0 mg/dL

by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD) PREDIABETIC: 100.0 - 125.0 DIABETIC: > 0R = 126.0

INTERPRETATION
IN ACCORDANCE WITH AMERICAN DIABETES ASSOCIATION GUIDELINES:

1. A fasting plasma glucose level below 100 mg/dl is considered normal.

2. A fasting plasma glucose level between 100 - 125 mg/dl is considered as glucose intolerant or prediabetic. A fasting and post-prandial blood

test (after consumption of 75 gms of glucose) is recommended for all such patients.

3. A fasting plasma glucose level of above 125 mg/dl is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for all such patients. A fasting plasma glucose level in excess of 125 mg/dl on both occasions is confirmatory for diabetic state.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)







# PKR JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

# A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

**NAME** : Mr. RAM SINGH

**AGE/ GENDER** : 70 YRS/MALE **PATIENT ID** : 1691245

**COLLECTED BY** REG. NO./LAB NO. : 122412050003

REFERRED BY **REGISTRATION DATE** : 05/Dec/2024 08:31 AM BARCODE NO. : 12506000 **COLLECTION DATE** : 05/Dec/2024 03:10PM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 05/Dec/2024 01:20PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

| Test Name  | Value         | Unit      | Biological Reference interval   |
|--|---------------|-----------|---|
|  | LIPID PROFILI | E : BASIC |   |
| CHOLESTEROL TOTAL: SERUM by CHOLESTEROL OXIDASE PAP            | 157.89        | mg/dL     | OPTIMAL: < 200.0<br>BORDERLINE HIGH: 200.0 -<br>239.0<br>HIGH CHOLESTEROL: > OR =<br>240.0  |
| TRIGLYCERIDES: SERUM by GLYCEROL PHOSPHATE OXIDASE (ENZYMATIC) | 140.47        | mg/dL     | OPTIMAL: < 150.0<br>BORDERLINE HIGH: 150.0 -<br>199.0<br>HIGH: 200.0 - 499.0<br>VERY HIGH: > OR = 500.0                                 |
| HDL CHOLESTEROL (DIRECT): SERUM by SELECTIVE INHIBITION        | 36.8          | mg/dL     | LOW HDL: < 30.0<br>BORDERLINE HIGH HDL: 30.0 -<br>60.0<br>HIGH HDL: > OR = 60.0   |
| LDL CHOLESTEROL: SERUM by CALCULATED, SPECTROPHOTOMETRY        | 93            | mg/dL     | OPTIMAL: < 100.0<br>ABOVE OPTIMAL: 100.0 - 129.0<br>BORDERLINE HIGH: 130.0 -<br>159.0<br>HIGH: 160.0 - 189.0<br>VERY HIGH: > OR = 190.0 |
| NON HDL CHOLESTEROL: SERUM by CALCULATED, SPECTROPHOTOMETRY    | 121.09        | mg/dL     | OPTIMAL: < 130.0<br>ABOVE OPTIMAL: 130.0 - 159.0<br>BORDERLINE HIGH: 160.0 -<br>189.0<br>HIGH: 190.0 - 219.0<br>VERY HIGH: > OR = 220.0 |
| VLDL CHOLESTEROL: SERUM by CALCULATED, SPECTROPHOTOMETRY       | 28.09         | mg/dL     | 0.00 - 45.00  |
| TOTAL LIPIDS: SERUM by CALCULATED, SPECTROPHOTOMETRY           | 456.25        | mg/dL     | 350.00 - 700.00   |
| CHOLESTEROL/HDL RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY  | 4.29          | RATIO     | LOW RISK: 3.30 - 4.40<br>AVERAGE RISK: 4.50 - 7.0<br>MODERATE RISK: 7.10 - 11.0<br>HIGH RISK: > 11.0                                    |



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)







# A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

NAME : Mr. RAM SINGH

AGE/ GENDER : 70 YRS/MALE **PATIENT ID** : 1691245

**COLLECTED BY** REG. NO./LAB NO. : 122412050003

REFERRED BY **REGISTRATION DATE** : 05/Dec/2024 08:31 AM BARCODE NO. **COLLECTION DATE** : 05/Dec/2024 03:10PM : 12506000 CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 05/Dec/2024 01:20PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

| Test Name   | Value | Unit  | Biological Reference interval   |
|---|-------|-------|---|
| LDL/HDL RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY           | 2.53  | RATIO | LOW RISK: 0.50 - 3.0<br>MODERATE RISK: 3.10 - 6.0<br>HIGH RISK: > 6.0 |
| TRIGLYCERIDES/HDL RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY | 3.82  | RATIO | 3.00 - 5.00   |

#### **INTERPRETATION:**

1. Measurements in the same patient can show physiological analytical variations. Three serial samples 1 week apart are recommended for Total Cholesterol, Triglycerides, HDL & LDL Cholesterol.

2. As per NLA-2014 guidelines, all adults above the age of 20 years should be screened for lipid status. Selective screening of children above the age of 2 years with a family history of premature cardiovascular disease or those with at least one parent with high total cholesterol is recommended.

3. Low HDL levels are associated with increased risk for Atherosclerotic Cardiovascular disease (ASCVD) due to insufficient HDL being available

to participate in reverse cholesterol transport, the process by which cholesterol is eliminated from peripheral tissues.

4. NLA-2014 identifies Non HDL Cholesterol (an indicator of all atherogeniclipoproteins such as LDL, VLDL, IDL, Lpa, Chylomicron remnants) along with LDL-cholesterol as co-primary target for cholesterol lowering therapy. Note that major risk factors can modify treatment goals for LDL &Non

5. Additional testing for Apolipoprotein B, hsCRP,Lp(a) & LP-PLA2 should be considered among patients with moderate risk for ASCVD for risk refinement



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)





# A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

**NAME** : Mr. RAM SINGH

**AGE/ GENDER** : 70 YRS/MALE **PATIENT ID** : 1691245

**COLLECTED BY** : 122412050003 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 05/Dec/2024 08:31 AM BARCODE NO. : 12506000 **COLLECTION DATE** : 05/Dec/2024 03:10PM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 05/Dec/2024 01:20PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**Test Name Value** Unit **Biological Reference interval** 

#### **LIVER FUNCTION TEST (COMPLETE)**

| BILIRUBIN TOTAL: SERUM by DIAZOTIZATION, SPECTROPHOTOMETRY                           | 0.85              | mg/dL | INFANT: 0.20 - 8.00<br>ADULT: 0.00 - 1.20 |
|--|-------------------|-------|---|
| BILIRUBIN DIRECT (CONJUGATED): SERUM by DIAZO MODIFIED, SPECTROPHOTOMETRY            | 0.32              | mg/dL | 0.00 - 0.40                               |
| BILIRUBIN INDIRECT (UNCONJUGATED): SERUM by CALCULATED, SPECTROPHOTOMETRY            | 0.53              | mg/dL | 0.10 - 1.00                               |
| SGOT/AST: SERUM<br>by IFCC, WITHOUT PYRIDOXAL PHOSPHATE                              | 21.56             | U/L   | 7.00 - 45.00                              |
| SGPT/ALT: SERUM<br>by IFCC, WITHOUT PYRIDOXAL PHOSPHATE                              | 18.15             | U/L   | 0.00 - 49.00                              |
| AST/ALT RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY                                | 1.19              | RATIO | 0.00 - 46.00                              |
| ALKALINE PHOSPHATASE: SERUM by PARA NITROPHENYL PHOSPHATASE BY AMINO METHYL PROPANOL | 112.43            | U/L   | 40.0 - 130.0                              |
| GAMMA GLUTAMYL TRANSFERASE (GGT): SERUM by SZASZ, SPECTROPHTOMETRY                   | 20.92             | U/L   | 0.00 - 55.0                               |
| TOTAL PROTEINS: SERUM by BIURET, SPECTROPHOTOMETRY                                   | 6.29              | gm/dL | 6.20 - 8.00                               |
| ALBUMIN: SERUM by BROMOCRESOL GREEN  | 4.21              | gm/dL | 3.50 - 5.50                               |
| GLOBULIN: SERUM by CALCULATED, SPECTROPHOTOMETRY                                     | $2.08^{L}$        | gm/dL | 2.30 - 3.50                               |
| A : G RATIO: SERUM   | 2.02 <sup>H</sup> | RATIO | 1.00 - 2.00                               |

#### INTERPRETATION

by CALCULATED, SPECTROPHOTOMETRY

NOTE:- To be correlated in individuals having SGOT and SGPT values higher than Normal Referance Range.

**USE**:- Differential diagnosis of diseases of hepatobiliary system and pancreas.

#### INCREASED:

| DRUG HEPATOTOXICITY                          | > 2                        |
|--|----------------------------|
| ALCOHOLIC HEPATITIS                          | > 2 (Highly Suggestive)    |
| CIRRHOSIS                                    | 1.4 - 2.0                  |
| INTRAHEPATIC CHOLESTATIS                     | > 1.5                      |
| HEPATOCELLULAR CARCINOMA & CHRONIC HEPATITIS | > 1.3 (Slightly Increased) |



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)







# A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

**NAME** : Mr. RAM SINGH

**AGE/ GENDER** : 70 YRS/MALE **PATIENT ID** : 1691245

**COLLECTED BY** REG. NO./LAB NO. : 122412050003

REFERRED BY **REGISTRATION DATE** : 05/Dec/2024 08:31 AM BARCODE NO. : 12506000 **COLLECTION DATE** : 05/Dec/2024 03:10PM

CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 05/Dec/2024 01:20PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**Test Name Value** Unit **Biological Reference interval** 

#### **DECREASED:**

1. Acute Hepatitis due to virus, drugs, toxins (with AST increased 3 to 10 times upper limit of normal)

2. Extra Hepatic cholestatis: 0.8 (normal or slightly decreased).

#### PROGNOSTIC SIGNIFICANCE:

| NORMAL               | < 0.65    |
|----------------------|-----------|
| GOOD PROGNOSTIC SIGN | 0.3 - 0.6 |
| POOR PROGNOSTIC SIGN | 1.2 - 1.6 |



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)





# PKR JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

### A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

8.50 - 10.60

2.30 - 4.70

**NAME** : Mr. RAM SINGH

**AGE/ GENDER** : 70 YRS/MALE **PATIENT ID** : 1691245

**COLLECTED BY** REG. NO./LAB NO. : 122412050003

REFERRED BY **REGISTRATION DATE** : 05/Dec/2024 08:31 AM BARCODE NO. **COLLECTION DATE** : 05/Dec/2024 03:10PM : 12506000 CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 05/Dec/2024 04:26PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

| Test Name   | Value         | Unit           | Biological Reference interval |
|---|---------------|----------------|-------------------------------|
|   |               |                |                               |
| KIDNE   | EY FUNCTION T | EST (COMPLETE) |                               |
| UREA: SERUM   | 31.77         | mg/dL          | 10.00 - 50.00                 |
| by UREASE - GLUTAMATE DEHYDROGENASE (GLDH)                        |               |                |                               |
| CREATININE: SERUM by ENZYMATIC, SPECTROPHOTOMETERY                | 0.94          | mg/dL          | 0.40 - 1.40                   |
| BLOOD UREA NITROGEN (BUN): SERUM by CALCULATED, SPECTROPHOTOMETRY | 14.85         | mg/dL          | 7.0 - 25.0                    |
| BLOOD UREA NITROGEN (BUN)/CREATININE<br>RATIO: SERUM              | 15.8          | RATIO          | 10.0 - 20.0                   |
| by CALCULATED, SPECTROPHOTOMETRY                                  |               |                |                               |
| UREA/CREATININE RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY     | 33.8          | RATIO          |                               |
| URIC ACID: SERUM by uricase - oxidase peroxidase                  | 3.63          | mg/dL          | 3.60 - 7.70                   |

mg/dL

mg/dL

### by ARSENAZO III, SPECTROPHOTOMETRY PHOSPHOROUS: SERUM

by PHOSPHOMOLYBDATE, SPECTROPHOTOMETRY

#### **ELECTROLYTES**

CALCIUM: SERUM

SODIUM: SERUM 142.6 mmol/L 135.0 - 150.0 by ISE (ION SELECTIVE ELECTRODE) POTASSIUM: SERUM 4.57 mmol/L 3.50 - 5.00by ISE (ION SELECTIVE ELECTRODE) 90.0 - 110.0 CHLORIDE: SERUM 106.95 mmol/L by ISE (ION SELECTIVE ELECTRODE)

9.34

2.52

#### **ESTIMATED GLOMERULAR FILTERATION RATE**

ESTIMATED GLOMERULAR FILTERATION RATE 87.2

(eGFR): SERUM by CALCULATED **INTERPRETATION:** 

To differentiate between pre- and post renal azotemia.

#### INCREASED RATIO (>20:1) WITH NORMAL CREATININE:

- 1. Prerenal azotemia (BUN rises without increase in creatinine) e.g. heart failure, salt depletion, dehydration, blood loss) due to decreased glomerular filtration rate.
- Catabolic states with increased tissue breakdown.
- 3. GI haemorrhage.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)





# A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

: 05/Dec/2024 04:26PM

**NAME** : Mr. RAM SINGH

AGE/ GENDER : 70 YRS/MALE **PATIENT ID** : 1691245

**COLLECTED BY** REG. NO./LAB NO. : 122412050003

REFERRED BY **REGISTRATION DATE** : 05/Dec/2024 08:31 AM BARCODE NO. **COLLECTION DATE** : 05/Dec/2024 03:10PM : 12506000

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

**Test Name Value** Unit **Biological Reference interval** 

REPORTING DATE

4. High protein intake.

CLIENT CODE.

5. Impaired renal function plus

6. Excess protein intake or production or tissue breakdown (e.g. infection, GI bleeding, thyrotoxicosis, Cushing's syndrome, high protein diet, burns, surgery, cachexia, high fever).

7. Urine reabsorption (e.g. ureter colostomy)

8. Reduced muscle mass (subnormal creatinine production)

9. Certain drugs (e.g. tetracycline, glucocorticoids)

#### INCREASED RATIO (>20:1) WITH ELEVATED CREATININE LEVELS:

- 1. Postrenal azotemia (BUN rises disproportionately more than creatinine) (e.g. obstructive uropathy).
- 2. Prerenal azotemia superimposed on renal disease.

#### DECREASED RATIO (<10:1) WITH DECREASED BUN:

- 1. Acute tubular necrosis.
- 2. Low protein diet and starvation.
- 3. Severe liver disease.
- 4. Other causes of decreased urea synthesis.
- 5. Repeated dialysis (urea rather than creatinine diffuses out of extracellular fluid).
- 6. Inherited hyperammonemias (urea is virtually absent in blood).
- 7. SIADH (syndrome of inappropiate antidiuretic harmone) due to tubular secretion of urea.
- 8. Pregnancy.

#### **DECREASED RATIO (<10:1) WITH INCREASED CREATININE:**

- 1. Phenacimide therapy (accelerates conversion of creatine to creatinine).
- 2. Rhabdomyolysis (releases muscle creatinine).
- 3. Muscular patients who develop renal failure.

#### **INAPPROPIATE RATIO:**

- 1. Diabetic ketoacidosis (acetoacetate causes false increase in creatinine with certain methodologies, resulting in normal ratio when dehydration should produce an increased BUN/creatinine ratio).
- 2. Cephalosporin therapy (interferes with creatinine measurement). **ESTIMATED GLOMERULAR FILTERATION RATE**:

| CKD STAGE | DESCRIPTION                           | GFR ( mL/min/1.73m2 ) | ASSOCIATED FINDINGS                               |
|-----------|---------------------------------------|-----------------------|---|
| G1        | Normal kidney function                | >90                   | No proteinuria                                    |
| G2        | Kidney damage with normal or high GFR | >90                   | Presence of Protein ,<br>Albumin or cast in urine |
| G3a       | Mild decrease in GFR                  | 60 -89                |   |
| G3b       | Moderate decrease in GFR              | 30-59                 |   |
| G4        | Severe decrease in GFR                | 15-29                 |   |
| G5        | Kidney failure                        | <15                   |   |



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)







# A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

REPORTING DATE

: 05/Dec/2024 04:26PM

**NAME** : Mr. RAM SINGH

AGE/ GENDER : 70 YRS/MALE **PATIENT ID** : 1691245

**COLLECTED BY** REG. NO./LAB NO. : 122412050003

REFERRED BY **REGISTRATION DATE** : 05/Dec/2024 08:31 AM BARCODE NO. **COLLECTION DATE** : 05/Dec/2024 03:10PM : 12506000

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

**Test Name Value** Unit **Biological Reference interval** 

COMMENTS:

CLIENT CODE.

1. Estimated Glomerular filtration rate (eGFR) is the sum of filtration rates in all functioning nephrons and so an estimation of the GFR provides a measure of functioning nephrons of the kidney.

2. eGFR calculated using the 2009 CKD-EPI creatinine equation and GFR category reported as per KDIGO guideline 2012

3. In patients, with eGFR creatinine between 45-59 ml/min/1.73 m2 (G3) and without any marker of Kidney damage, It is recommended to measure eGFR with Cystatin C for confirmation of CKD

4. eGFR category G1 OR G2 does not fullfill the criteria for CKD, in the absence of evidence of Kidney Damage
5. In a suspected case of Acute Kidney Injury (AKI), measurement of eGFR should be done after 48-96 hours of any Intervention or procedure
6. eGFR calculated by Serum Creatinine may be less accurate due to certain factors like Race, Muscle Mass, Diet, Certain Drugs. In such cases, eGFR should be calculated using Serum Cystatin C
7. A decrease in eGFR implies either progressive renal disease, or a reversible process causing decreased nephron function (eg, severe dehydration).

KDIGO guideline, 2012 recommends Chronic Kidney Disease (CKD) should be classified based on cause, eGFR category and Albuminuria (ACR) category. GFR & ACR category combined together reflect risk of progression and helps Clinician to identify the individual who are progressing at more rapid rate than anticipated



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)





# A PIONEER DIAGNOSTIC CENTRE

**NAME** : Mr. RAM SINGH

AGE/ GENDER : 70 YRS/MALE **PATIENT ID** : 1691245

**COLLECTED BY** REG. NO./LAB NO. : 122412050003

REFERRED BY **REGISTRATION DATE** : 05/Dec/2024 08:31 AM BARCODE NO. : 12506000 **COLLECTION DATE** : 05/Dec/2024 03:10PM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 05/Dec/2024 04:39PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**Value** Unit **Biological Reference interval Test Name** 

# **TUMOUR MARKER** PROSTATE SPECIFIC ANTIGEN (PSA) - TOTAL

PROSTATE SPECIFIC ANTIGEN (PSA) - TOTAL: 0.78 ng/mL 0.0 - 4.0

**SERUM** 

by CLIA (CHEMILUMINESCENCE IMMUNOASSAY)

#### INTERPRETATION:

#### NOTE:

- 1. This is a recommended test for detection of prostate cancer along with Digital Rectal Examination (DRE) in males above 50 years of age.
- 2. False negative / positive results are observed in patients receiving mouse monoclonal antibodies for diagnosis or therapy
- 3. PSA levels may appear consistently elevated / depressed due to the interference by heterophilic antibodies & nonspecific protein binding 4. Immediate PSA testing following digital rectal examination, ejaculation, prostatic massage, indwelling catheterization, ultrasonography and needle biopsy of prostate is not recommended as they falsely elevate levels
- 5. PSA values regardless of levels should not be interpreted as absolute evidence of the presence or absence of disease. All values should be correlated with clinical findings and results of other investigations
- 6. Sites of Non-prostatic PSA production are breast epithelium, salivary glands, peri-urethral & anal glands, cells of male urethra & breast milk
- 7. Physiological decrease in PSA level by 18% has been observed in hospitalized / sedentary patients either due to supine position or suspended sexual activity
- 8. The concentration of PSA in a given specimen, determined with assays from different manufacturers, may not be comparable due to differences in assay methods, calibration, and reagent specificity.

#### RECOMMENDED TESTING INTERVALS

- 1. Preoperatively (Baseline)
- 2. 2-4 Days Post operatively
- 3. Prior to discharge from hospital

Monthly Follow I In if levels are high and showing a rising trend

| 4. Monthly Follow up it levels are flight and showing a risi | ng trend             |
|--|----------------------|
| POST SURGERY   | FREQUENCY OF TESTING |
| 1st Year   | Every 3 Months       |
| 2 <sup>nd</sup> Year   | Every 4 Months       |
| 3 <sup>rd</sup> Year Onwards                                 | Every 6 Months       |

#### CLINICAL USE:

- 1. An aid in the early detection of Prostate cancer when used in conjunction with Digital rectal examination in males more than 50 years of age and in those with two or more affected first degree relatives.
- 2. Followup and management of Prostate cancer patients.
- 3. Detect metastatic or persistent disease in patients following surgical or medical treatment of Prostate cancer

### **INCREASED LEVEL:**

- 1. Prostate cancer
- 2. Benign Prostatic Hyperplasia
- 3. Prostatitis
- 4. Genitourinary infections



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)



CLIENT CODE.



# PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

# A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

: 05/Dec/2024 04:39PM

: Mr. RAM SINGH NAME

**AGE/ GENDER** : 70 YRS/MALE **PATIENT ID** : 1691245

**COLLECTED BY** REG. NO./LAB NO. : 122412050003

REFERRED BY **REGISTRATION DATE** : 05/Dec/2024 08:31 AM BARCODE NO. : 12506000 **COLLECTION DATE** : 05/Dec/2024 03:10PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

**Test Name Value** Unit **Biological Reference interval** 

REPORTING DATE



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS, MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST







# A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

: 05/Dec/2024 01:20PM

**NAME** : Mr. RAM SINGH

**AGE/ GENDER** : 70 YRS/MALE **PATIENT ID** : 1691245

**COLLECTED BY** REG. NO./LAB NO. : 122412050003

REFERRED BY **REGISTRATION DATE** : 05/Dec/2024 08:31 AM BARCODE NO. : 12506000 **COLLECTION DATE** : 05/Dec/2024 03:10PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

**Value** Unit **Biological Reference interval Test Name** 

REPORTING DATE

### **CLINICAL PATHOLOGY URINE ROUTINE & MICROSCOPIC EXAMINATION**

#### PHYSICAL EXAMINATION

CLIENT CODE.

| QUANTITY RECIEVED                          | 30          | ml |               |
|--|-------------|----|---------------|
| by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY |             |    |               |
| COLOUR                                     | PALE YELLOW |    | PALE YELLOW   |
| by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY |             |    |               |
| TRANSPARANCY                               | CLEAR       |    | CLEAR         |
| by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY |             |    |               |
| SPECIFIC GRAVITY                           | 1.01        |    | 1.002 - 1.030 |
|  |             |    |               |

| by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY                |                |       |                |
|---|----------------|-------|----------------|
| CHEMICAL EXAMINATION                                      |                |       |                |
| REACTION by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY       | ACIDIC         |       |                |
| PROTEIN by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY        | NEGATIVE (-ve) |       | NEGATIVE (-ve) |
| SUGAR by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY          | NEGATIVE (-ve) |       | NEGATIVE (-ve) |
| pH by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY             | 6.5            |       | 5.0 - 7.5      |
| BILIRUBIN by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY      | NEGATIVE (-ve) |       | NEGATIVE (-ve) |
| NITRITE by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY.       | NEGATIVE (-ve) |       | NEGATIVE (-ve) |
| UROBILINOGEN by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY   | NOT DETECTED   | EU/dL | 0.2 - 1.0      |
| KETONE BODIES by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY  | NEGATIVE (-ve) |       | NEGATIVE (-ve) |
| BLOOD by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY          | NEGATIVE (-ve) |       | NEGATIVE (-ve) |
| ASCORBIC ACID  by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY | NEGATIVE (-ve) |       | NEGATIVE (-ve) |



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS, MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST

/HPF

0 - 3



NEGATIVE (-ve)

**MICROSCOPIC EXAMINATION** RED BLOOD CELLS (RBCs)





# A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

NAME : Mr. RAM SINGH

**AGE/ GENDER** : 70 YRS/MALE **PATIENT ID** : 1691245

**COLLECTED BY** REG. NO./LAB NO. : 122412050003

REFERRED BY **REGISTRATION DATE** : 05/Dec/2024 08:31 AM BARCODE NO. : 12506000 **COLLECTION DATE** : 05/Dec/2024 03:10PM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 05/Dec/2024 01:20PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

| Test Name  | Value          | Unit | Biological Reference interval |
|--|----------------|------|-------------------------------|
| by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT                                  |                |      |                               |
| PUS CELLS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT                        | 2-3            | /HPF | 0 - 5                         |
| EPITHELIAL CELLS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT                 | 1-2            | /HPF | ABSENT                        |
| CRYSTALS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT                         | NEGATIVE (-ve) |      | NEGATIVE (-ve)                |
| CASTS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT                            | NEGATIVE (-ve) |      | NEGATIVE (-ve)                |
| BACTERIA by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT                         | NEGATIVE (-ve) |      | NEGATIVE (-ve)                |
| OTHERS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT                           | NEGATIVE (-ve) |      | NEGATIVE (-ve)                |
| TRICHOMONAS VAGINALIS (PROTOZOA) by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT | ABSENT         |      | ABSENT                        |

End Of Report



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS, MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST

