

PKR JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

NAME : Mrs. BINDU

AGE/ GENDER : 27 YRS/FEMALE **PATIENT ID** : 1361907

COLLECTED BY REG. NO./LAB NO. : 122412140019

REFERRED BY **REGISTRATION DATE** : 14/Dec/2024 04:25 PM BARCODE NO. : 12506155 **COLLECTION DATE** : 14/Dec/2024 04:34PM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 14/Dec/2024 09:21PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name Value Unit **Biological Reference interval**

HAEMATOLOGY

PROTHROMBIN TIME STUDIES (PT/INR)

PT TEST (PATIENT) by PHOTO OPTICAL CLOT DETECTION	13.8	SECS	11.5 - 14.5
PT (CONTROL) by PHOTO OPTICAL CLOT DETECTION	12	SECS	
ISI by PHOTO OPTICAL CLOT DETECTION	1.1		
INTERNATIONAL NORMALISED RATIO (INR) by PHOTO OPTICAL CLOT DETECTION	1.17		0.80 - 1.20
PT INDEX	86.96	%	

INTERPRETATION:-

- 1.INR is the parameter of choice in monitoring adequacy of oral anti-coagulant therapy. Appropiate therapeutic range varies with the disease and treatment intensity.
- 2. Prolonged INR suggests potential bleeding disorder /bleeding complications
- 3. Results should be clinically correlated.
- 4. Test conducted on Citrated Plasma

RECOMMENDED THERAPEUTIC RANGE FOR ORAL ANTI-COAGULANT THERAPY (INR)				
INDICATION		INTERNATIONAL NORMALIZED RATIO (INR)		
Treatment of venous thrombosis				
Treatment of pulmonary embolism				
Prevention of systemic embolism in tissue heart valves]			
Valvular heart disease	Low Intensity	2.0 - 3.0		
Acute myocardial infarction				
Atrial fibrillation				
Bileaflet mechanical valve in aortic position				
Recurrent embolism				
Mechanical heart valve	High Intensity	2.5 - 3.5		



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)







PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

REPORTING DATE

: 14/Dec/2024 09:21PM

NAME : Mrs. BINDU

AGE/ GENDER : 27 YRS/FEMALE **PATIENT ID** : 1361907

COLLECTED BY REG. NO./LAB NO. : 122412140019

REFERRED BY **REGISTRATION DATE** : 14/Dec/2024 04:25 PM BARCODE NO. : 12506155 **COLLECTION DATE** : 14/Dec/2024 04:34PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

Test Name Value Unit **Biological Reference interval**

Antiphospholipid antibodies+

CLIENT CODE.

The prothrombin time (PT) and its derived measures of prothrombin ratio (PR) and international normalized ratio (INR) are measures of the efficacy of the extrinsic pathway of coagulation. PT test reflects the adequacy of factors I (fibrinogen), II (prothrombin), V, VII, and X. It is used in conjunction with the activated partial thromboplastin time (aPTT) which measures the intrinsic pathway.

The common causes of prolonged prothrombin time are:

1. Oral Anticoagulant therapy.

2.Liver disease.

3. Vit K. deficiency.

4. Disseminated intra vascular coagulation.

5. Factor 5, 7, 10 or Prothrombin dificiency

End Of Report



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)