## NASIRPUR, Hissar Road, AMBALA CITY- (Haryana) A PIONEER DIAGNOSTIC CENTRE

🔽 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

**PKR JAIN HEALTHCARE INSTITUTE** 

NAME	: Mrs. MUSKAN				
AGE/ GENDER : 20 YRS/FEMALE		Р	ATIENT ID	: 1701575	
COLLECTED BY	:	R	EG. NO./LAB NO.	: 122412170015	
REFERRED BY	:	R	EGISTRATION DATE	: 17/Dec/2024 02:14 PM	
BARCODE NO.	: 12506191	С	OLLECTION DATE	: 17/Dec/2024 04:27PM	
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTI	TUTE <b>R</b>	EPORTING DATE	: 17/Dec/2024 05:47PM	
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AME	BALA CITY - HARY	YANA		
Test Name		Value	Unit	Biological Reference interval	
		ENDOCR	INOLOGY		
	THY	ROID FUNCT	ION TEST: TOTAL		
TRIIODOTHYRONINE (T3): SERUM by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)		1.203 AY)	ng/mL	0.35 - 1.93	
THYROXINE (T4): SERUM by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)		8.02 AY)	µgm/dL	4.87 - 12.60	
	TING HORMONE (TSH): SERUM		µIU/mL	0.35 - 5.50	
· · · · · · · · · · · · · · · · · · ·	RASENSITIVE				
3rd GENERATION, ULT INTERPRETATION:	RASENSITIVE	etween 2-4 a.m and a			

day has influence on the measured serum TSH concentrations. TSH stimulates the production and secretion of the metabolically active hormones, thyroxine (T4) and triiodothyronine (T3). Failure at any level of regulation of the hypothalamic-pituitary-thyroid axis will result in either underproduction (hypothyroidism) or overproduction(hyperthyroidism) of T4 and/or T3.

CLINICAL CONDITION	T3	T4	TSH	
Primary Hypothyroidism:	Reduced	Reduced	Increased (Significantly)	
Subclinical Hypothyroidism:	Normal or Low Normal	Normal or Low Normal	High	
Primary Hyperthyroidism:	Increased	Increased	Reduced (at times undetectable)	
Subclinical Hyperthyroidism:	Normal or High Normal	Normal or High Normal	Reduced	

#### LIMITATIONS:-

1. T3 and T4 circulates in reversibly bound form with Thyroid binding globulins (TBG), and to a lesser extent albumin and Thyroid binding Pre Albumin so conditions in which TBG and protein levels alter such as pregnancy, excess estrogens, androgens, anabolic steroids and glucocorticoids may falsely affect the T3 and T4 levels and may cause false thyroid values for thyroid function tests.

2. Normal levels of T4 can also be seen in Hyperthyroid patients with :T3 Thyrotoxicosis, Decreased binding capacity due to hypoproteinemia or ingestion of certain drugs (e.g.: phenytoin , salicylates).

3. Serum T4 levels in neonates and infants are higher than values in the normal adult , due to the increased concentration of TBG in neonate serum.

4. TSH may be normal in central hypothyroidism , recent rapid correction of hyperthyroidism or hypothyroidism , pregnancy , phenytoin therapy.

TRIIODOTHYRONINE (T3)		THYROXINE (T4)		THYROID STIMULATING HORMONE (TSH)		
Age	Refferance Range (ng/mL)	Age	Refferance Range ( µg/dL)	Age	Reference Range ( µIU/mL)	
0-7 Days	0.20 - 2.65	0 - 7 Days	5.90 - 18.58	0 - 7 Days	2.43 - 24.3	
7 Days - 3 Months	0.36 - 2.59	7 Days - 3 Months	6.39 - 17.66	7 Days - 3 Months	0.58 - 11.00	
3 - 6 Months	0.51 - 2.52	3 - 6 Months	6.75 - 17.04	3 Days – 6 Months	0.70 - 8.40	
6 - 12 Months	0.74 - 2.40	6 - 12 Months	7.10 - 16.16	6 – 12 Months	0.70 - 7.00	





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA

CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)

440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)





# PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana) A PIONEER DIAGNOSTIC CENTRE

NAME	: Mrs. MUSKAN			
AGE/ GENDER	: 20 YRS/FEMALE	PATIENT ID	: 1701575	
COLLECTED BY	:	REG. NO./LAB NO.	: 122412170015	
<b>REFERRED BY</b>	:	<b>REGISTRATION DATE</b>	: 17/Dec/2024 02:14 PM	
BARCODE NO.	: 12506191	<b>COLLECTION DATE</b>	: 17/Dec/2024 04:27PM	
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	<b>REPORTING DATE</b>	: 17/Dec/2024 05:47PM	
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA			

Test Name			Value	Unit	t	Biological Reference interval
1 - 10 Years	0.92 - 2.28	1 - 10 Years	6.00 - 13.80	1 – 10 Years	0.60 - 5.50	
11- 19 Years	0.35 - 1.93	11 - 19 Years	4.87-13.20	11 – 19 Years	0.50 - 5.50	
> 20 years (Adults)	0.35 - 1.93	> 20 Years (Adults)	4.87 - 12.60	> 20 Years (Adults)	0.35-5.50	
	RECO	MMENDATIONS OF TSH LE	EVELS DURING PRE	GNANCY ( µIU/mL)		
1st Trimester			0.10 - 2.50			
2nd Trimester			0.20 - 3.00			
	3rd Trimester			0.30 - 4.10		

### **INCREASED TSH LEVELS:**

1. Primary or untreated hypothyroidism may vary from 3 times to more than 100 times normal depending upon degree of hypofunction.

2. Hypothyroid patients receiving insufficient thyroid replacement therapy.

3.Hashimotos thyroiditis

4.DRUGS: Amphetamines, iodine containing agents & dopamine antagonist.

5.Neonatal period, increase in 1st 2-3 days of life due to post-natal surge

### DECREASED TSH LEVELS:

1.Toxic multi-nodular goiter & Thyroiditis.

2. Over replacement of thyroid hormone in treatment of hypothyroidism.

3. Autonomously functioning Thyroid adenoma

4.Secondary pituitary or hypothalamic hypothyroidism

5. Acute psychiatric illness

6.Severe dehydration.

7.DRUGS: Glucocorticoids, Dopamine, Levodopa, T4 replacement therapy, Anti-thyroid drugs for thyrotoxicosis. 8.Pregnancy: 1st and 2nd Trimester

\*\*\* End Of Report \*\*\*





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)

440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600, REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)

