



P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

☎ 0171-2532620, 8222896961 ✉ pkrjainhealthcare@gmail.com

TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT.

NAME	: Mr. BALWINDER KUMAR	PATIENT ID	: 1704126
AGE/ GENDER	: 54 YRS/MALE	REG. NO./LAB NO.	: 122412200013
COLLECTED BY	:	REGISTRATION DATE	: 20/Dec/2024 11:21 AM
REFERRED BY	:	COLLECTION DATE	: 20/Dec/2024 11:29AM
BARCODE NO.	: 12506234	REPORTING DATE	: 20/Dec/2024 07:06PM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE		
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------


CLINICAL CHEMISTRY/BIOCHEMISTRY

KIDNEY FUNCTION TEST (BASIC)

UREA: SERUM <i>by UREASE - GLUTAMATE DEHYDROGENASE (GLDH)</i>	30.66	mg/dL	10.00 - 50.00
CREATININE: SERUM <i>by ENZYMATIC, SPECTROPHOTOMETRY</i>	1.44 ^H	mg/dL	0.40 - 1.40
BLOOD UREA NITROGEN (BUN): SERUM <i>by CALCULATED, SPECTROPHOTOMETRY</i>	14.33	mg/dL	7.0 - 25.0
BLOOD UREA NITROGEN (BUN)/CREATININE RATIO: SERUM <i>by CALCULATED, SPECTROPHOTOMETRY</i>	9.95 ^L	RATIO	10.0 - 20.0
UREA/CREATININE RATIO: SERUM <i>by CALCULATED, SPECTROPHOTOMETRY</i>	21.29	RATIO	
URIC ACID: SERUM <i>by URICASE - OXIDASE PEROXIDASE</i>	4.18	mg/dL	3.60 - 7.70




DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)


DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY)





P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

☎ 0171-2532620, 8222896961 ✉ pkrajainhealthcare@gmail.com

NAME	: Mr. BALWINDER KUMAR	PATIENT ID	: 1704126
AGE/ GENDER	: 54 YRS/MALE	REG. NO./LAB NO.	: 122412200013
COLLECTED BY	:	REGISTRATION DATE	: 20/Dec/2024 11:21 AM
REFERRED BY	:	COLLECTION DATE	: 20/Dec/2024 11:29AM
BARCODE NO.	: 12506234	REPORTING DATE	: 20/Dec/2024 07:06PM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE		
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

INTERPRETATION:

Normal range for a healthy person on normal diet: 12 - 20

To Differentiate between pre- and postrenal azotemia.

INCREASED RATIO (>20:1) WITH NORMAL CREATININE:

- 1.Prerenal azotemia (BUN rises without increase in creatinine) e.g. heart failure, salt depletion,dehydration, blood loss) due to decreased glomerular filtration rate.
- 2.Catabolic states with increased tissue breakdown.
- 3.GI hemorrhage.
- 4.High protein intake.
- 5.Impaired renal function plus .
- 6.Excess protein intake or production or tissue breakdown (e.g. infection, GI bleeding, thyrotoxicosis, Cushings syndrome, high protein diet, burns,surgery, cachexia, high fever).
- 7.Urine reabsorption (e.g. ureterocolostomy)
- 8.Reduced muscle mass (subnormal creatinine production)
- 9.Certain drugs (e.g. tetracycline, glucocorticoids)

INCREASED RATIO (>20:1) WITH ELEVATED CREATININE LEVELS:

- 1.Postrenal azotemia (BUN rises disproportionately more than creatinine) (e.g. obstructive uropathy).
- 2.Prerenal azotemia superimposed on renal disease.

DECREASED RATIO (<10:1) WITH DECREASED BUN :

- 1.Acute tubular necrosis.
- 2.Low protein diet and starvation.
- 3.Severe liver disease.
- 4.Other causes of decreased urea synthesis.
- 5.Repeated dialysis (urea rather than creatinine diffuses out of extracellular fluid).
- 6.Inherited hyperammonemias (urea is virtually absent in blood).
- 7.SIADH (syndrome of inappropriate antidiuretic hormone) due to tubular secretion of urea.
- 8.Pregnancy.

DECREASED RATIO (<10:1) WITH INCREASED CREATININE:


- 1.Phenacimide therapy (accelerates conversion of creatine to creatinine).
- 2.Rhabdomyolysis (releases muscle creatinine).
- 3.Muscular patients who develop renal failure.

INAPPROPRIATE RATIO:

- 1.Diabetic ketoacidosis (acetoacetate causes false increase in creatinine with certain methodologies,resulting in normal ratio when dehydration should produce an increased BUN/creatinine ratio).
- 2.Cephalosporin therapy (interferes with creatinine measurement).




DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)


DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)





P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

☎ 0171-2532620, 8222896961 ✉ pkrjainhealthcare@gmail.com

NAME : Mr. BALWINDER KUMAR
AGE/ GENDER : 54 YRS/MALE
COLLECTED BY :
REFERRED BY :
BARCODE NO. : 12506234
CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE
CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

PATIENT ID : 1704126
REG. NO./LAB NO. : 122412200013
REGISTRATION DATE : 20/Dec/2024 11:21 AM
COLLECTION DATE : 20/Dec/2024 11:29AM
REPORTING DATE : 22/Dec/2024 04:27PM

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

MICROBIOLOGY

CULTURE AEROBIC BACTERIA AND ANTIBIOTIC SENSITIVITY: URINE

CULTURE AND SUSCEPTIBILITY: URINE

DATE OF SAMPLE 20-12-2024
SPECIMEN SOURCE URINE
INCUBATION PERIOD 48 HOURS
by AUTOMATED BROTH CULTURE
CULTURE STERILE
by AUTOMATED BROTH CULTURE
ORGANISM NO AEROBIC PYOGENIC ORGANISM GROWN AFTER 48 HOURS OF
by AUTOMATED BROTH CULTURE INCUBATION AT 37°C

AEROBIC SUSCEPTIBILITY: URINE

INTERPRETATION:

1. In urine culture and sensitivity, presence of more than 100,000 organism per mL in midstream sample of urine is considered clinically significant. However in symptomatic patients, a smaller number of bacteria (100 to 10000/mL) may signify infection.
2. Colony count of 100 to 10000/ mL indicate infection, if isolate from specimen obtained by suprapubic aspiration or "in-and-out" catheterization or from patients with indwelling catheters.

SUSCEPTIBILITY:

1. A test interpreted as **SENSITIVE** implies that infection due to isolate may be appropriately treated with the dosage of an antimicrobial agent recommended for that type of infection and infecting species, unless otherwise indicated..
2. A test interpreted as **INTERMEDIATE** implies that the "infection due to the isolate may be appropriately treated in body sites where the drugs are physiologically concentrated or when a high dosage of drug can be used".
3. A test interpreted as **RESISTANT** implies that the "isolates are not inhibited by the usually achievable concentration of the agents with normal dosage, schedule and/or fall in the range where specific microbial resistance mechanism are likely (e.g. beta-lactamases), and clinical efficacy has not been reliable in treatment studies.

CAUTION:


Conditions which can cause a false Negative culture:

1. Patient is on antibiotics. Please repeat culture post therapy.
2. Anaerobic bacterial infection.
3. Fastidious aerobic bacteria which are not able to grow on routine culture media.
4. Besides all these factors, at least in 25-40 % of cases there is no direct correlation between in vivo clinical picture.
5. Renal tuberculosis to be confirmed by AFB studies.

*** End Of Report ***




DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)


DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY)

