

PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

NAME : Mrs. MINAKSHI

AGE/ GENDER : 36 YRS/FEMALE **PATIENT ID** : 1705952

COLLECTED BY REG. NO./LAB NO. : 122412220006

REFERRED BY **REGISTRATION DATE** : 22/Dec/2024 11:52 AM BARCODE NO. : 12506260 **COLLECTION DATE** : 22/Dec/2024 12:02PM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 22/Dec/2024 06:27PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Value Unit **Biological Reference interval Test Name**

CLINICAL CHEMISTRY/BIOCHEMISTRY **AMYLASE**

AMYLASE - SERUM 41.43 IU/L 0 - 90

by CNPG 3, SPECTROPHOTOMETRY

INTERPRETATION COMMENTS

1. Amylase is produced in the Pancreas and most of the elevation in serum is due to increased rate of Amylase entry into the blood stream / decreased rate of clearance or both.

2.Serum Amylase rises within 6 to 48 hours of onset of Acute pancreatitis in 80% of patients, but is not proportional to the severity of the disease.

3.Activity usually returns to normal in 3-5 days in patients with milder edematous form of the disease.

4.Values persisting longer than this period suggest continuing necrosis of pancreas or Pseudocyst formation.

5.Approximately 20% of patients with Pancreatitis have normal or near normal activity.

6. Hyperlipemic patients with Pancreatitis also show spuriously normal Amylase levels due to suppression of Amylase activity by triglyceride. 7. Low Amylase levels are seen in Chronic Pancreatitis, Congestive Heart failure, 2nd & 3rd trimesters of pregnancy, Gastrointestinal cancer &

bone fractures.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)



PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

NAME : Mrs. MINAKSHI

AGE/ GENDER : 36 YRS/FEMALE **PATIENT ID** : 1705952

COLLECTED BY REG. NO./LAB NO. : 122412220006

REFERRED BY **REGISTRATION DATE** : 22/Dec/2024 11:52 AM BARCODE NO. : 12506260 **COLLECTION DATE** : 22/Dec/2024 12:02PM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 22/Dec/2024 06:27PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Value Unit **Biological Reference interval Test Name**

LIPASE

LIPASE - SERUM 37.46 U/L 0 - 60

by METHYL RESORUFIN, SPECTROPHOTOMETRY

INTERPRETATION

- 1. Pancreas is the major and primary source of serum lipase though lipases are also present in liver, stomach, intestine, WBC, fat cells and milk.
 2. In acute pancreatitis, serum lipase becomes elevated at the same time as amylase and remains high for 7-10 days.
- 3. Increased lipase activity rarely lasts longer than 14 days
- 4. Prolonged increase suggests poor prognosis or presence of a cyst.
- The combined use of serum lipase and serum amylase is effective in ruling out acute pancreatitis.

INCREASED LEVEL:

- 1. Acute & Chronic pancreatitis
- 2. Obstruction of pancreatic duct
- 3. Non pancreatic conditions like renal diseases, acute cholecystitis, intestinal obstruction, duodenal ulcer, alcoholism, diabetic ketoacidosis and following endoscopic retrograde cholangiopancreatography
- 1. Elevations 2 to 50 times the upper reference have been reported. The increase in serum lipase is not necessarily proportional to the severity of the attack. Normalization is not necessarily a sign of resolution.

Concomitant testing of serum amylase and lipase is highly recommended to establish a diagnosis of pancreatic injury

*** End Of Report ***



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)

