



# PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

## A PIONEER DIAGNOSTIC CENTRE

**NAME** : Mrs. INDERJEET KAUR

**AGE/ GENDER** : 29 YRS/FEMALE **PATIENT ID** : 1691525

**COLLECTED BY** : 122501040002 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 04/Jan/2025 10:32 AM BARCODE NO. : 12506385 **COLLECTION DATE** : 04/Jan/2025 11:00AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 04/Jan/2025 01:02PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**Value** Unit **Biological Reference interval Test Name** 

# **ENDOCRINOLOGY** THYROID STIMULATING HORMONE (TSH)

THYROID STIMULATING HORMONE (TSH): SERUM 2.53

μIU/mL 0.35 - 5.50

by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

3rd GENERATION, ULTRASENSITIVE

#### INTERPRETATION:

AGE	REFFERENCE RANGE (μIU/mL)
0 – 5 DAYS	0.70 - 15.20
6 Days – 2 Months	0.70 – 11.00
3 – 11 Months	0.70 - 8.40
1 – 5 Years	0.70 – 7.00
6 – 10 Years	0.60 - 5.50
11 - 15	0.50 - 5.50
> 20 Years (Adults)	0.27 - 5.50
PREGNANCY	
1st Trimester	0.10 - 3.00
2nd Trimester	0.20 - 3.00
3rd Trimester	0.30 - 4.10

NOTE:-TSH levels are subjected to circardian variation, reaching peak levels between 2-4 a.m and at a minimum between 6-10 pm. The variation is of the order of 50 %. Hence time of the day has influence on the measured serum TSH concentration.

USE: TSH controls biosynthesis and release of thyroid harmones T4 & T3. It is a sensitive measure of thyroid function, especially useful in early or subclinical hypothyroidism, before the patient develops any clinical findings or goitre or any other thyroid function abnormality.

#### **INCREASED LEVELS:**

- 1. Primary or untreated hypothyroidism, may vary from 3 times to more than 100 times normal depending on degree of hypofunction.
- 2. Hypothyroid patients receiving insufficient thyroid replacement therapy.
- 3. Hashimotos thyroiditis.
- 4.DRUGS: Amphetamines, Iodine containing agents and dopamine antagonist.
- 5. Neonatal period, increase in 1st 2-3 days of life due to post-natal surge.

### **DECREASED LEVELS:**

- 1. Toxic multi-nodular goitre & Thyroiditis.
- 2. Over replacement of thyroid harmone in treatment of hypothyroidism.
- 3. Autonomously functioning Thyroid adenoma
- 4. Secondary pituatary or hypothalmic hypothyroidism
- 5. Acute psychiatric illness
- 6. Severe dehydration.
- 7.DRUGS: Glucocorticoids, Dopamine, Levodopa, T4 replacement therapy, Anti-thyroid drugs for thyrotoxicosis.

CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)





# PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

## A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

NAME : Mrs. INDERJEET KAUR

**AGE/ GENDER** : 29 YRS/FEMALE **PATIENT ID** : 1691525

**COLLECTED BY** REG. NO./LAB NO. : 122501040002

REFERRED BY **REGISTRATION DATE** : 04/Jan/2025 10:32 AM BARCODE NO. : 12506385 **COLLECTION DATE** : 04/Jan/2025 11:00AM

CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 04/Jan/2025 01:02PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**Test Name Value** Unit **Biological Reference interval** 

8. Pregnancy: 1st and 2nd Trimester LIMITATIONS:

1.TSH may be normal in central hypothyroidism, recent rapid correction of hyperthyroidism or hypothyroidism, pregnancy, phenytoin therapy.

2. Autoimmune disorders may produce spurious results.

DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)



# PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

## A PIONEER DIAGNOSTIC CENTRE

**NAME** : Mrs. INDERJEET KAUR

AGE/ GENDER : 29 YRS/FEMALE **PATIENT ID** : 1691525

**COLLECTED BY** REG. NO./LAB NO. : 122501040002

REFERRED BY **REGISTRATION DATE** : 04/Jan/2025 10:32 AM BARCODE NO. **COLLECTION DATE** : 04/Jan/2025 11:00AM : 12506385 CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 06/Jan/2025 03:50PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**Value** Unit **Biological Reference interval Test Name** 

## MICROBIOLOGY

### CULTURE AEROBIC BACTERIA AND ANTIBIOTIC SENSITIVITY: URINE

### **CULTURE AND SUSCEPTIBILITY: URINE**

DATE OF SAMPLE 04-01-2025 SPECIMEN SOURCE URINE INCUBATION PERIOD 48 HOURS by AUTOMATED BROTH CULTURE

**CULTURE** by AUTOMATED BROTH CULTURE

by AUTOMATED BROTH CULTURE

NO AEROBIC PYOGENIC ORGANISM GROWN AFTER 48 HOURS OF **ORGANISM** 

**STERILE** 

**INCUBATION AT 37\*C** 

## **AEROBIC SUSCEPTIBILITY: URINE**

#### INTERPRETATION:

1. In urine culture and sensitivity, presence of more than 100,000 organism per mL in midstream sample of urine is considered clinically significant. However in symptomatic patients, a smaller number of bacteria (100 to 10000/mL) may signify infection.

2. Colony count of 100 to 10000/ mL indicate infection, if isolate from specimen obtained by suprapubic aspiration or "in-and-out"

catheterization or from patients with indwelling catheters.

#### SUSCEPTIBILITY:

1. A test interpreted as SENSTITIVE implies that infection due to isolate may be appropriately treated with the dosage of an antimicrobial agent recommended for that type of infection and infecting species, unless otherwise indicated..

2. A test interpreted as **INTERMEDIATE** implies that the" Infection due to the isolate may be appropriately treated in body sites where the drugs are

physiologically concentrated or when a high dosage of drug can be used".

3.A test interpreted as **RESISTANT** implies that the "isolates are not inhibited by the usually achievable concentration of the agents with normal dosage, schedule and/or fall in the range where specific microbial resistance mechanism are likely (e.g. beta-lactamases), and clinical efficacy has not been reliable in treatment studies.

## **CAUTION:**

Conditions which can cause a false Negative culture:

- 1. Patient is on antibiotics. Please repeat culture post therapy.
- 2. Anaerobic bacterial infection.
- 3. Fastidious aerobic bacteria which are not able to grow on routine culture media.
- 4. Besides all these factors, at least in 25-40 % of cases there is no direct correlation between in vivo clinical picture.
- 5. Renal tuberculosis to be confirmed by AFB studies.

\*\*\* End Of Report \*\*\*



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)