TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT

PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana) A PIONEER DIAGNOSTIC CENTRE

【 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mr. AJAY GUPTA				
AGE/ GENDER	: 52 YRS/MALE		PATIENT ID	: 1728482	
COLLECTED BY	:		REG. NO./LAB NO.	: 122501200002	
REFERRED BY	:		REGISTRATION DATE	: 20/Jan/2025 08:51 AM	
BARCODE NO.	: 12506584		COLLECTION DATE	: 20/Jan/2025 09:02AM	
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITU	TE	REPORTING DATE	: 20/Jan/2025 01:16PM	
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBAL	A CITY - H	ARYANA		
Test Name		Value	Unit	Biological Reference interv	val
	SWAST	HYA W	ELLNESS PANEL: 1.	.4	
	COMP	LETE B	LOOD COUNT (CBC)		
RED BLOOD CELLS	S (RBCS) COUNT AND INDICES				
HAEMOGLOBIN (H	B)	16.6	gm/dL	12.0 - 17.0	
RED BLOOD CELL ((RBC) COUNT	5.29 ^H	Millions	s/cmm 3.50 - 5.00	
PACKED CELL VOL	UME (PCV) AUTOMATED HEMATOLOGY ANALYZER	45.7	%	40.0 - 54.0	
MEAN CORPUSCUL		86.4	KR fl	80.0 - 100.0	
MEAN CORPUSCUL	AR HAEMOGLOBIN (MCH)	31.5	pg	27.0 - 34.0	
	AR HEMOGLOBIN CONC. (MCHC)	36.4 ^H	g/dL	32.0 - 36.0	
RED CELL DISTRIB	UTION WIDTH (RDW-CV)	13	%	11.00 - 16.00	
RED CELL DISTRIB	UTION WIDTH (RDW-SD) AUTOMATED HEMATOLOGY ANALYZER	43.5	fL	35.0 - 56.0	
MENTZERS INDEX by CALCULATED		16.33	RATIO	BETA THALASSEMIA TRAI 13.0 IRON DEFICIENCY ANEMI	
GREEN & KING INI by calculated	DEX	21.31	RATIO	>13.0 BETA THALASSEMIA TRAI 65.0 IRON DEFICIENCY ANEMIA 65.0	[T:<=
WHITE BLOOD CE					
,	Y BY SF CUBE & MICROSCOPY	6100	/cmm	4000 - 11000	
	<u>UCOCYTE COUNT (DLC)</u>		24	70 70	
NEUTROPHILS by FLOW CYTOMETRY	Y BY SF CUBE & MICROSCOPY	57	%	50 - 70	
LYMPHOCYTES		38	%	20 - 40	



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS , MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST



: Mr. AJAY GUPTA

NAME

PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

【 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mr. AJAY GUPTA			
AGE/ GENDER	: 52 YRS/MALE]	PATIENT ID	: 1728482
COLLECTED BY	:]	REG. NO./LAB NO.	: 122501200002
REFERRED BY	:	1	REGISTRATION DATE	: 20/Jan/2025 08:51 AM
BARCODE NO.	: 12506584		COLLECTION DATE	: 20/Jan/2025 09:02AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTIT	LUTE 1	REPORTING DATE	: 20/Jan/2025 01:16PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMB	ALA CITY - HAF	RYANA	
Test Name		Value	Unit	Biological Reference interval
by FLOW CYTOMET	RY BY SF CUBE & MICROSCOPY			
EOSINOPHILS		0 ^L	%	1 - 6
by FLOW CYTOMETF MONOCYTES	RY BY SF CUBE & MICROSCOPY	5	%	2 - 12
	RY BY SF CUBE & MICROSCOPY	5	70	2 - 12
BASOPHILS		0	%	0 - 1
	RY BY SF CUBE & MICROSCOPY			
	<u>OCYTES (WBC) COUNT</u>			
ABSOLUTE NEUTI by FLOW CYTOMETE	ROPHIL COUNT	3477	/cmm	2000 - 7500
ABSOLUTE LYMPH	HOCYTE COUNT	2318	/cmm	800 - 4900
by FLOW CYTOMETF	RY BY SF CUBE & MICROSCOPY	0 ^L	/cmm	40 - 440
	RY BY SF CUBE & MICROSCOPY	01	/ СШШ	40 - 440
ABSOLUTE MONO	CYTE COUNT	305	/cmm	80 - 880
	RY BY SF CUBE & MICROSCOPY			0.440
ABSOLUTE BASOF	PHIL COUNT RY BY SF CUBE & MICROSCOPY	0	/cmm	0 - 110
-	OTHER PLATELET PREDICTIVE	MARKERS.		
PLATELET COUNT by HYDRO DYNAMIC	" (PLT) FOCUSING, ELECTRICAL IMPEDENCE	256000	/cmm	150000 - 450000
PLATELETCRIT (P	CT) FOCUSING, ELECTRICAL IMPEDENCE	0.24	%	0.10 - 0.36
MEAN PLATELET		9	fL	6.50 - 12.0
by HYDRO DYNAMIC	FOCUSING, ELECTRICAL IMPEDENCE		111	
	CELL COUNT (P-LCC) FOCUSING, ELECTRICAL IMPEDENCE	58000	/cmm	30000 - 90000
PLATELET LARGE by HYDRO DYNAMIC	CELL RATIO (P-LCR) FOCUSING, ELECTRICAL IMPEDENCE	22.7	%	11.0 - 45.0
PLATELET DISTRI	BUTION WIDTH (PDW) FOCUSING, ELECTRICAL IMPEDENCE	15.9	%	15.0 - 17.0
	UCTED ON EDTA WHOLE BLOOD			





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS , MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST



A PIONEER DIAGNOSTIC CENTRE

🔽 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mr. AJAY GUPTA				
AGE/ GENDER	: 52 YRS/MALE		PATIENT ID	: 1728482	
COLLECTED BY	:	:	REG. NO./LAB NO.	: 122501200002	
REFERRED BY	•		REGISTRATION DATE	: 20/Jan/2025 08:51	лм
BARCODE NO.	: 12506584		COLLECTION DATE	: 20/Jan/2025 09:02	AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTI	ГИТЕ	REPORTING DATE	: 20/Jan/2025 04:14	PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMB	IBALA CITY - HARYANA			
Test Name		Value	Unit	Biological	Reference interva
	GLYCOS	SYLATED HA	EMOGLOBIN (HBA1)	C)	
GLYCOSYLATED HAEMOGLOBIN (HbA1c): WHOLE BLOOD by HPLC (HIGH PERFORMANCE LIQUID CHROMATOGRAPHY)		5.8	%	4.0 - 6.4	
ESTIMATED AVERAGE PLASMA GLUCOSE by HPLC (HIGH PERFORMANCE LIQUID CHROMATOGRAPHY) INTERPRETATION:		119.76	mg/dL	60.00 - 140	0.00
	AS PER AMERICAN DI	ABETES ASSOCIA	TION (ADA).		
	REFERENCE GROUP		YCOSYLATED HEMOGLOGIB	(HBAIC) in %	
Non dia	abetic Adults >= 18 years				
A	t Risk (Prediabetes)		5.7 – 6.4		
Diagnosing Diabetes		>= 6.5			
			Age > 19 Years		
Thoropout	is goals for glycomic control		of Therapy:	< 7.0	
rnerapeut	ic goals for glycemic control	Actions	Suggested:	>8.0	
		Cert	Age < 19 Years	7.5	
COMMENTS:		Goal	of therapy:	<7.5	

1.Glycosylated hemoglobin (HbA1c) test is three monthly monitoring done to assess compliace with therapeutic regimen in diabetic patients. 2.Since Hb1c reflects long term fluctuations in blood glucose concentration, a diabetic patient who has recently under good control may still have high concentration of HbAlc. Converse is true for a diabetic previously under good control but now poorly controlled.

3. Target goals of < 7.0 % may be beneficial in patients with short duration of diabetes, long life expectancy and no significant cardiovascular disease. In patients with significant complications of diabetes, limited life expectancy or extensive co-morbid conditions, targetting a goal of < 7.0% may not be appropriate.

4.High HbA1c (>9.0 -9.5 %) is strongly associated with risk of development and rapid progression of microvascular and nerve complications 5.Any condition that shorten RBC life span like acute blood loss, hemolytic anemia falsely lower HbA1c results.

6.HbA1c results from patients with HbSS,HbSC and HbD must be interpreted with caution, given the pathological processes including anemia, increased red cell turnover, and transfusion requirement that adversely impact HbA1c as a marker of long-term gycemic control.

7.Specimens from patients with polycythemia or post-splenctomy may exhibit increse in HbA1c values due to a somewhat longer life span of the red cells.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)



A PIONEER DIAGNOSTIC CENTRE

🔽 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mr. AJAY GUPTA		
AGE/ GENDER	: 52 YRS/MALE	PATIENT ID	: 1728482
COLLECTED BY	:	REG. NO./LAB NO.	: 122501200002
REFERRED BY	:	REGISTRATION DATE	: 20/Jan/2025 08:51 AM
BARCODE NO.	: 12506584	COLLECTION DATE	: 20/Jan/2025 09:02AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	REPORTING DATE	: 20/Jan/2025 03:47PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY	- HARYANA	
Test Name	Value	Unit	Biological Reference interval
	ic test because an elevated result often indic	ates the presence of inflammat	ion associated with infection, cancer and auto
INTERPRETATION:	GATION BY CAPILLARY PHOTOMETRY		
immune disease, but	does not tell the health practitioner exactly v	where the inflammation is in the	e body or what is causing it.
 An ESR can be affe as C-reactive protein 		n. For this reason, the ESR is ty	pically used in conjunction with other test suc
3. This test may also	be used to monitor disease activity and respo	onse to therapy in both of the a	bove diseases as well as some others, such as
systemic lupus erythe	ematosus W FSR		
A low ESR can be see	n with conditions that inhibit the normal sed	imentation of red blood cells, s	uch as a high red blood cell count
(polycythaemia), sigr	hificantly high white blood cell count (leucocy le cell anaemia) also lower the ESR.	/tosis) , and some protein abno	rmalities. Šome changes in red cell shape (suc
NOTE:	also lower the Esk.		
1. ESR and C - reactiv	e protein (C-RP) are both markers of inflamma	ation.	
2. Generally, ESR doe	es not change as rapidly as does CRP, either at by as many other factors as is ESR, making it a	t the start of inflammation or as	s it resolves.
4. If the ESR is elevat	ed, it is typically a result of two types of prote	eins, globulins or fibrinogen.	
5 Women tend to ha	we a higher FSR and menstruation and pregn	ancy can cause temporary eleva	ations

5. Women tend to have a higher ESR, and menstruation and pregnancy can cause temporary elevations. 6. Drugs such as dextran, methyldopa, oral contraceptives, penicillamine procainamide, theophylline, and vitamin A can increase ESR, while

aspirin, cortisone, and quinine may decrease it





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)

440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. **REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)**



TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT

A PIONEER DIAGNOSTIC CENTRE

🔽 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

INSTITUTE	PATIENT ID REG. NO./LAB NO. REGISTRATION DATE COLLECTION DATE REPORTING DATE	: 1728482 : 122501200002 : 20/Jan/2025 08:51 AM : 20/Jan/2025 09:02AM : 20/Jan/2025 10:29AM
	REGISTRATION DATE COLLECTION DATE REPORTING DATE	: 20/Jan/2025 08:51 AM : 20/Jan/2025 09:02AM
	COLLECTION DATE REPORTING DATE	: 20/Jan/2025 09:02AM
	REPORTING DATE	
		: 20/Jan/2025 10:29AM
AMDALA CITY IIA		
), AMBALA CITY - HA	ARTANA	
Value	Unit	Biological Reference interval
NICAL CHEMIS	TRY/BIOCHEMIST	RV
	E FASTING (F)	
103.73 ^H	mg/dL	NORMAL: < 100.0 PREDIABETIC: 100.0 - 125.0 DIABETIC: > 0R = 126.0
	103.73 ^H	103.73^H mg/dL

A fasting plasma glucose level below 100 mg/dl is considered normal.
 A fasting plasma glucose level between 100 - 125 mg/dl is considered as glucose intolerant or prediabetic. A fasting and post-prandial blood

A fasting plasma glucose level of above 125 mg/dl is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for all such patients.
 A fasting plasma glucose level of above 125 mg/dl is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for all such patients.





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



A PIONEER DIAGNOSTIC CENTRE

【 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mr. AJAY GUPTA			
AGE/ GENDER	: 52 YRS/MALE		PATIENT ID	: 1728482
COLLECTED BY	:		REG. NO./LAB NO.	: 122501200002
REFERRED BY	:		REGISTRATION DATE	: 20/Jan/2025 08:51 AM
BARCODE NO.	: 12506584		COLLECTION DATE	: 20/Jan/2025 09:02AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INS	STITUTE	REPORTING DATE	: 20/Jan/2025 10:29AM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HA		ARYANA	
Test Name		Value	Unit	Biological Reference interval
		LIPID PR	OFILE : BASIC	
CHOLESTEROL TO by CHOLESTEROL OX		202.59 ^H	mg/dL	OPTIMAL: < 200.0 BORDERLINE HIGH: 200.0 - 239.0 HIGH CHOLESTEROL: > OR = 240.0
TRIGLYCERIDES: S by GLYCEROL PHOSF	ERUM PHATE OXIDASE (ENZYMATIC)	164.66 ^H	mg/dL	OPTIMAL: < 150.0 BORDERLINE HIGH: 150.0 - 199.0 HIGH: 200.0 - 499.0 VERY HIGH: > OR = 500.0
HDL CHOLESTERO	L (DIRECT): SERUM ion	36.52	mg/dL	LOW HDL: < 30.0 BORDERLINE HIGH HDL: 30.0 60.0 HIGH HDL: > OR = 60.0
LDL CHOLESTEROI by CALCULATED, SPE		133.14 ^H	mg/dL	OPTIMAL: < 100.0 ABOVE OPTIMAL: 100.0 - 129 BORDERLINE HIGH: 130.0 - 159.0 HIGH: 160.0 - 189.0 VERY HIGH: > OR = 190.0
NON HDL CHOLEST by CALCULATED, SPE		166.07 ^H	mg/dL	OPTIMAL: < 130.0 ABOVE OPTIMAL: 130.0 - 159 BORDERLINE HIGH: 160.0 - 189.0 HIGH: 190.0 - 219.0 VERY HIGH: > OR = 220.0
VLDL CHOLESTER(32.93	mg/dL	0.00 - 45.00
TOTAL LIPIDS: SER	RUM	569.84	mg/dL	350.00 - 700.00
CHOLESTEROL/HD by CALCULATED, SPE		5.55 ^H	RATIO	LOW RISK: 3.30 - 4.40 AVERAGE RISK: 4.50 - 7.0 MODERATE RISK: 7.10 - 11.0 HIGH RISK: > 11.0

DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS , MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST



🔽 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mr. AJAY GUPTA		
AGE/ GENDER	: 52 YRS/MALE	PATIENT ID	: 1728482
COLLECTED BY	:	REG. NO./LAB NO.	: 122501200002
REFERRED BY	:	REGISTRATION DATE	: 20/Jan/2025 08:51 AM
BARCODE NO.	: 12506584	COLLECTION DATE	: 20/Jan/2025 09:02AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	REPORTING DATE	: 20/Jan/2025 10:29AM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY -	HARYANA	

Test Name	Value	Unit	Biological Reference interval
LDL/HDL RATIO: SERUM by Calculated, spectrophotometry	3.65 ^H	RATIO	LOW RISK: 0.50 - 3.0 MODERATE RISK: 3.10 - 6.0 HIGH RISK: > 6.0
TRIGLYCERIDES/HDL RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	4.51	RATIO	3.00 - 5.00

INTERPRETATION:

1. Measurements in the same patient can show physiological analytical variations. Three serial samples 1 week apart are recommended for Total Cholesterol, Triglycerides, HDL & LDL Cholesterol.

2. As per NLA-2014 guidelines, all adults above the age of 20 years should be screened for lipid status. Selective screening of children above the age of 2 years with a family history of premature cardiovascular disease or those with at least one parent with high total cholesterol is recommended.

3. Low HDL levels are associated with increased risk for Atherosclerotic Cardiovascular disease (ASCVD) due to insufficient HDL being available

to participate in reverse cholesterol transport, the process by which cholesterol is eliminated from peripheral tissues. 4. NLA-2014 identifies Non HDL Cholesterol (an indicator of all atherogeniclipoproteins such as LDL, VLDL, IDL, Lpa, Chylomicron remnants) along with LDL-cholesterol as co- primary target for cholesterol lowering therapy. Note that major risk factors can modify treatment goals for LDL & Non HDL.

5. Additional testing for Apolipoprotein B, hsCRP,Lp(a) & LP-PLA2 should be considered among patients with moderate risk for ASCVD for risk refinement



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)





A PIONEER DIAGNOSTIC CENTRE

🕻 0171-2532620, 8222896961 🛛 pkrjainhealthcare@gmail.com

NAME	: Mr. AJAY GUPTA			
AGE/ GENDER	: 52 YRS/MALE		PATIENT ID	: 1728482
COLLECTED BY	:		REG. NO./LAB NO.	: 122501200002
REFERRED BY	:		REGISTRATION DATE	: 20/Jan/2025 08:51 AM
BARCODE NO.	: 12506584		COLLECTION DATE	: 20/Jan/2025 09:02AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITU	UTE	REPORTING DATE	: 20/Jan/2025 11:30AM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBA	LA CITY - H	ARYANA	
Test Name		Value	Unit	Biological Reference interval
	LIVER	FUNCTIO	N TEST (COMPLETE)	
BILIRUBIN TOTAL	: SERUM PECTROPHOTOMETRY	0.83	mg/dL	INFANT: 0.20 - 8.00 ADULT: 0.00 - 1.20
	C (CONJUGATED): SERUM	0.3	mg/dL	0.00 - 0.40
BILIRUBIN INDIRE	CT (UNCONJUGATED): SERUM	0.53	mg/dL	0.10 - 1.00
SGOT/AST: SERUM by IFCC, WITHOUT PY	RIDOXAL PHOSPHATE	31.54	U/L	7.00 - 45.00
SGPT/ALT: SERUM		59.24 ^H	U/L	0.00 - 49.00
AST/ALT RATIO: SI by CALCULATED, SPE		0.53	RATIO	0.00 - 46.00
ALKALINE PHOSPH by PARA NITROPHEN PROPANOL	HATASE: SERUM YL PHOSPHATASE BY AMINO METHYL	84.67	U/L	40.0 - 130.0
GAMMA GLUTAMY by SZASZ, SPECTROF	L TRANSFERASE (GGT): SERUM PHTOMETRY	44.36	U/L	0.00 - 55.0
TOTAL PROTEINS: by BIURET, SPECTRO		6.35	gm/dL	6.20 - 8.00
ALBUMIN: SERUM by BROMOCRESOL G		4.3	gm/dL	3.50 - 5.50
GLOBULIN: SERUM	-	2.05 ^L	gm/dL	2.30 - 3.50
A : G RATIO: SERUM		2.1 ^H	RATIO	1.00 - 2.00

INTERPRETATION

NOTE: To be correlated in individuals having SGOT and SGPT values higher than Normal Referance Range. USE: Differential diagnosis of diseases of hepatobiliary system and pancreas.

INCREASED:

DRUG HEPATOTOXICITY	> 2
ALCOHOLIC HEPATITIS	> 2 (Highly Suggestive)
CIRRHOSIS	1.4 - 2.0
INTRAHEPATIC CHOLESTATIS	> 1.5
HEPATOCELLULAR CARCINOMA & CHRONIC HEPATITIS	> 1.3 (Slightly Increased)





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)





【 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mr. AJAY GUPTA		
AGE/ GENDER	: 52 YRS/MALE	PATIENT ID	: 1728482
COLLECTED BY	:	REG. NO./LAB NO.	: 122501200002
REFERRED BY	:	REGISTRATION DATE	: 20/Jan/2025 08:51 AM
BARCODE NO.	: 12506584	COLLECTION DATE	: 20/Jan/2025 09:02AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	REPORTING DATE	: 20/Jan/2025 11:30AM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - H	IARYANA	

DECREASED:

1. Acute Hepatitis due to virus, drugs, toxins (with AST increased 3 to 10 times upper limit of normal)

2. Extra Hepatic cholestatis: 0.8 (normal or slightly decreased).

PROGNOSTIC	SIGNIFICANCE:

NORMAL	< 0.65
GOOD PROGNOSTIC SIGN	0.3 - 0.6
POOR PROGNOSTIC SIGN	1.2 - 1.6



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS , MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST





A PIONEER DIAGNOSTIC CENTRE

🔽 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mr. AJAY GUPTA					
AGE/ GENDER	: 52 YRS/MALE		PATIENT ID	: 1728482		
COLLECTED BY :		REG. NO./LAB NO.		: 122501200002		
REFERRED BY	:		REGISTRATION DATE	: 20/Jan/2025 08:51 AM		
BARCODE NO.	: 12506584		COLLECTION DATE	: 20/Jan/2025 09:02AM		
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTI	TUTE	REPORTING DATE	: 20/Jan/2025 10:29AM		
CLIENT ADDRESS			ARYANA			
Test Name		Value	Unit	Biological Reference interval		
	KIDNE	Y FUNCTI	ON TEST (COMPLETE))		
UREA: SERUM by UREASE - GLUTAM	ATE DEHYDROGENASE (GLDH)	34.16	mg/dL	10.00 - 50.00		
CREATININE: SERU		0.84	mg/dL	0.40 - 1.40		
BLOOD UREA NITR by CALCULATED, SPE	OGEN (BUN): SERUM CTROPHOTOMETRY	15.96	mg/dL	7.0 - 25.0		
BLOOD UREA NITR RATIO: SERUM by CALCULATED, SPE	OGEN (BUN)/CREATININE CTROPHOTOMETRY	19	RATIO	10.0 - 20.0		
UREA/CREATININE		40.67	RATIO			
URIC ACID: SERUM by URICASE - OXIDASI		6.5	mg/dL	3.60 - 7.70		
CALCIUM: SERUM by ARSENAZO III, SPE		9.64	mg/dL	8.50 - 10.60		
	RUM ATE, SPECTROPHOTOMETRY	3.17	mg/dL	2.30 - 4.70		
ELECTROLYTES		100 54	1./7	105.0 150.0		
SODIUM: SERUM by ISE (ION SELECTIVE	E ELECTRODE)	138.54	mmol/L	135.0 - 150.0		
POTASSIUM: SERUN by ISE (ION SELECTIVE	Л	4.19	mmol/L	3.50 - 5.00		
CHLORIDE: SERUM by ISE (ION SELECTIVE	E ELECTRODE)	103.91	mmol/L	90.0 - 110.0		
	ERULAR FILTERATION RATE ERULAR FILTERATION RATE	104.9				

INTERPRETATION:

To differentiate between pre- and post renal azotemia.

INCREASED RATIO (>20:1) WITH NORMAL CREATININE:

1. Prerenal azotemia (BUN rises without increase in creatinine) e.g. heart failure, salt depletion, dehydration, blood loss) due to decreased glomerular filtration rate.

2. Catabolic states with increased tissue breakdown.

3. GI haemorrhage.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS , MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST



A PIONEER DIAGNOSTIC CENTRE

【 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mr. A.	IAY GUPTA					
AGE/ GENDER	: 52 YR	S/MALE	I	PATIENT ID		: 1728482	
COLLECTED BY	:]	REG. NO./LAB NO.		: 122501200002	
REFERRED BY	:]	REGISTRATION D A	ATE	: 20/Jan/2025 08:51	AM
BARCODE NO.	: 12506	584		COLLECTION DATI	Е	: 20/Jan/2025 09:02	AM
CLIENT CODE.		IAIN HEALTHCARE INSTITUT		REPORTING DATE		: 20/Jan/2025 10:29	
CLIENT ADDRESS		PUR, HISSAR ROAD, AMBALA			6	. 20/ Juli/ 2020 10.20	
CLIENT ADDRESS	. INASIN						
Test Name			Value	Uni	it	Biological	Reference interval
9. Certain drugs (e.g. NCREASED RATIO (>2 I. Postrenal azotemia	(e.g. urel ass (subn tetracycl 0:1) WITH (BUN ris superimp 0:1) WITI osis.	ter colostomy) formal creatinine production) ine, glucocorticoids) HELEVATED CREATININE LEVEL es disproportionately more th bosed on renal disease. HDECREASED BUN :		ne) (e.g. obstructive	e uropath	ıy).	
3. Severe liver disease							
4. Other causes of de			t of outros	llulor fluid			
		er than creatinine diffuses ou is (urea is virtually absent in b		enular Huld).			
		piate antidiuretic harmone) d		ar secretion of urea	1.		
8. Pregnancy.							
		H INCREASED CREATININE:	to prophicin				
2. Rhabdomyolysis (r		erates conversion of creatine to buscle creatinine).	to creatinin	e).			
3. Muscular patients							
INAPPROPIATE RATIO	:						
		acetate causes false increase	in creatinir	e with certain meth	hodologi	ies,resulting in norma	I ratio when dehydrati
	apy (inte	BUN/creatinine ratio). rferes with creatinine measure E RATION RATE:	ement).				
CKD STAGE		DESCRIPTION	GFR (m	L/min/1.73m2)	ASSC	DCIATED FINDINGS	
G1		Normal kidney function	· ·	>90	٦	No proteinuria	

CKD STAGE	DESCRIPTION	GFR (mL/min/ 1.73mZ)	ASSOCIATED FINDINGS
G1	Normal kidney function	>90	No proteinuria
G2	Kidney damage with	>90	Presence of Protein,
	normal or high GFR		Albumin or cast in urine
G3a	Mild decrease in GFR	60 -89	
G3b	Moderate decrease in GFR	30-59	
G4	Severe decrease in GFR	15-29	
G5	Kidney failure	<15	





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS , MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST







A PIONEER DIAGNOSTIC CENTRE

0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mr. AJAY GUPTA		
AGE/ GENDER	: 52 YRS/MALE	PATIENT ID	: 1728482
COLLECTED BY	:	REG. NO./LAB NO.	: 122501200002
REFERRED BY	:	REGISTRATION DATE	: 20/Jan/2025 08:51 AM
BARCODE NO.	: 12506584	COLLECTION DATE	: 20/Jan/2025 09:02AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	REPORTING DATE	: 20/Jan/2025 10:29AM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - H	HARYANA	

Test Name	Value	Unit	Biological Reference interval

COMMENTS:

1. Estimated Glomerular filtration rate (eGFR) is the sum of filtration rates in all functioning nephrons and so an estimation of the GFR provides a measure of functioning nephrons of the kidney. 2. eGFR calculated using the 2009 CKD-EPI creatinine equation and GFR category reported as per KDIGO guideline 2012

3. In patients, with eGFR creatinine between 45-59 ml/min/1.73 m2 (G3) and without any marker of Kidney damage, It is recommended to measure eGFR with Cystatin C for confirmation of CKD

4. eGFR category G1 OR G2 does not fullfill the criteria for CKD, in the absence of evidence of Kidney Damage 5. In a suspected case of Acute Kidney Injury (AKI), measurement of eGFR should be done after 48-96 hours of any Intervention or procedure 6. eGFR calculated by Serum Creatinine may be less accurate due to certain factors like Race, Muscle Mass, Diet, Certain Drugs. In such cases, eGFR should be calculated using Serum Cystatin C 7. A decrease in eGFR implies either progressive renal disease, or a reversible process causing decreased nephron function (eg, severe dehydration).

ADVICE:

KDIGO guideline, 2012 recommends Chronic Kidney Disease (CKD) should be classified based on cause, eGFR category and Albuminuria (ACR) category. GFR & ACR category combined together reflect risk of progression and helps Clinician to identify the individual who are progressing at more rapid rate than anticipated



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)





🔽 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mr. AJAY GUPTA			
AGE/ GENDER	: 52 YRS/MALE	PATI	ENT ID	: 1728482
COLLECTED BY	:	REG.	NO./LAB NO.	: 122501200002
REFERRED BY	:	REGI	STRATION DATE	: 20/Jan/2025 08:51 AM
BARCODE NO.	: 12506584	COLL	ECTION DATE	: 20/Jan/2025 09:02AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTIT	UTE REPO	ORTING DATE	: 20/Jan/2025 04:43PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBA	LA CITY - HARYAN	A	
Test Name		Value	Unit	Biological Reference interval
		IRON PRO	FILE	
IRON: SERUM	TROPHOTOMETRY	86.2	µg/dL	59.0 - 158.0
•	ON BINDING CAPACITY (UIBC)	146.19 ^L	µg/dL	150.0 - 336.0
TOTAL IRON BIND	ING CAPACITY (TIBC)	232.39	μg/dL	230 - 430

restrume	Tuiu e	UIIIt	
	IRON	PROFILE	
IRON: SERUM by FERROZINE, SPECTROPHOTOMETRY	86.2	μg/dL	59.0 - 158.0
UNSATURATED IRON BINDING CAPA SERUM by FERROZINE, SPECTROPHOTOMETERY	CITY (UIBC) 146.19^L	µg/dL	150.0 - 336.0
TOTAL IRON BINDING CAPACITY (TI :SERUM by SPECTROPHOTOMETERY	BC) 232.39	µg/dL	230 - 430
%TRANSFERRIN SATURATION: SERI by calculated, spectrophotometery		%	15.0 - 50.0
TRANSFERRIN: SERUM by SPECTROPHOTOMETERY (FERENE)	165 ^L	mg/dL	200.0 - 350.0
INTERPRETATION:-			
VARIABLES A	NEMIA OF CHRONIC DISEASE	IRON DEFICIENCY ANEMIA	THALASSEMIA α/β TRAIT
SERUM IRON:	Normal to Reduced	Reduced	Normal

TOTAL IRON BINDING CAPACITY:

% TRANSFERRIN SATURATION:

SERUM FERRITIN:

IRON:

1.Serum iron studies is recommended for differential diagnosis of microcytic hypochromic anemia.i.e iron deficiency anemia, zinc deficiency anemia, anemia of chronic disease and thalassemia syndromes.

Increased

Decreased < 12-15 %

Decreased

Normal

Normal

Normal or Increased

It is essential to isolate iron deficiency anemia from Beta thalassemia syndromes because during iron replacement which is therapeutic for iron deficiency anemia, is severely contra-indicated in Thalassemia.
 TOTAL IRON BINDING CAPACITY (TIBC):

1.It is a direct measure of protein transferrin which transports iron from the gut to storage sites in the bone marrow. % TRANSFERRIN SATURATION:

Decreased

Decreased

Normal to Increased

1. Occurs in idiopathic hemochromatosis and transfusional hemosiderosis where no unsaturated iron binding capacity is available for iron mobilization. Similar condition is seen in congenital deficiency of transferrin.





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)





NAME	: Mr. AJAY GUPTA			
AGE/ GENDER	: 52 YRS/MALE	PATI	ENT ID	: 1728482
COLLECTED BY	:	REG.	NO./LAB NO.	: 122501200002
REFERRED BY	:	REGI	STRATION DATE	: 20/Jan/2025 08:51 AM
BARCODE NO.	: 12506584	COLL	ECTION DATE	: 20/Jan/2025 09:02AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUT	ГЕ Repo	RTING DATE	: 20/Jan/2025 12:17PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBAL	A CITY - HARYAN	A	
Test Name		Value	Unit	Biological Reference interva
Test Name		Value ENDOCRINO		Biological Reference interva
Test Name	THYRO	ENDOCRINO		Biological Reference interva
TRIIODOTHYRONII		ENDOCRINO	DLOGY	Biological Reference interva 0.35 - 1.93
TRIIODOTHYRONII by CMIA (CHEMILUMIN THYROXINE (T4): S	NE (T3): SERUM ESCENT MICROPARTICLE IMMUNOASSAY)	ENDOCRINO	DLOGY TEST: TOTAL	U
TRIIODOTHYRONII by CMIA (CHEMILUMIN THYROXINE (T4): S by CMIA (CHEMILUMIN THYROID STIMULA	NE (T3): SERUM ESCENT MICROPARTICLE IMMUNOASSAY) ERUM	ENDOCRING DID FUNCTION 0.95	DLOGY I TEST: TOTAL ng/mL	0.35 - 1.93
TRIIODOTHYRONII by CMIA (CHEMILUMIN THYROXINE (T4): S by CMIA (CHEMILUMIN THYROID STIMULA	NE (T3): SERUM ESCENT MICROPARTICLE IMMUNOASSAY) ERUM ESCENT MICROPARTICLE IMMUNOASSAY) TING HORMONE (TSH): SERUM ESCENT MICROPARTICLE IMMUNOASSAY)	ENDOCRING DID FUNCTION 0.95 7.16	DLOGY I TEST: TOTAL ng/mL µgm/dL	0.35 - 1.93 4.87 - 12.60

TSH levels are subject to circadian variation, reaching peak levels between 2-4 a.m and at a minimum between 6-10 pm. The variation is of the order of 50%. Hence time of the day has influence on the measured serum TSH concentrations. TSH stimulates the production and secretion of the metabolically active hormones, thyroxine (T4) and triiodothyronine (T3). Failure at any level of regulation of the hypothalamic-pituitary-thyroid axis will result in either underproduction (hypothyroidism) or overproduction(hyperthyroidism) of T4 and/or T3.

CLINICAL CONDITION	T3	T4	TSH
Primary Hypothyroidism:	Reduced	Reduced	Increased (Significantly)
Subclinical Hypothyroidism:	Normal or Low Normal	Normal or Low Normal	High
Primary Hyperthyroidism:	Increased	Increased	Reduced (at times undetectable)
Subclinical Hyperthyroidism:	Normal or High Normal	Normal or High Normal	Reduced

LIMITATIONS:-

1. T3 and T4 circulates in reversibly bound form with Thyroid binding globulins (TBG), and to a lesser extent albumin and Thyroid binding Pre Albumin so conditions in which TBG and protein levels alter such as pregnancy, excess estrogens, androgens, anabolic steroids and glucocorticoids may falsely affect the T3 and T4 levels and may cause false thyroid values for thyroid function tests.

2. Normal levels of T4 can also be seen in Hyperthyroid patients with :T3 Thyrotoxicosis, Decreased binding capacity due to hypoproteinemia or ingestion of certain drugs (e.g.: phenytoin , salicylates).

3. Serum T4 levels in neonates and infants are higher than values in the normal adult , due to the increased concentration of TBG in neonate serum.

4. TSH may be normal in central hypothyroidism , recent rapid correction of hyperthyroidism or hypothyroidism , pregnancy , phenytoin therapy.

TRIIODOTH	YRONINE (T3)	THYROXINE (T4)		THYROID STIMU	LATING HORMONE (TSH)
Age	Refferance Range (ng/mL)	Age	Refferance Range (µg/dL)	Age	Reference Range (μIU/mL)
0-7 Days	0.20 - 2.65	0 - 7 Days	5.90 - 18.58	0 - 7 Days	2.43 - 24.3
7 Days - 3 Months	0.36 - 2.59	7 Days - 3 Months	6.39 - 17.66	7 Days - 3 Months	0.58 - 11.00
3 - 6 Months	0.51 - 2.52	3 - 6 Months	6.75 - 17.04	3 Days – 6 Months	0.70 - 8.40
6 - 12 Months	0.74 - 2.40	6 - 12 Months	7.10 - 16.16	6 – 12 Months	0.70 - 7.00





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)





A PIONEER DIAGNOSTIC CENTRE

💟 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mr. AJAY GUPTA		
AGE/ GENDER	: 52 YRS/MALE	PATIENT ID	: 1728482
COLLECTED BY	:	REG. NO./LAB NO.	: 122501200002
REFERRED BY	:	REGISTRATION DATE	: 20/Jan/2025 08:51 AM
BARCODE NO.	: 12506584	COLLECTION DATE	: 20/Jan/2025 09:02AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	REPORTING DATE	: 20/Jan/2025 12:17PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - H	IARYANA	

Test Name		Value	Unit		Biological Reference interval	
1 - 10 Years	0.92 - 2.28	1 - 10 Years	6.00 - 13.80	1 – 10 Years	0.60 - 5.50	
11-19 Years	0.35 - 1.93	11 - 19 Years	4.87-13.20	11 – 19 Years	0.50 - 5.50	
> 20 years (Adults)	0.35 - 1.93	> 20 Years (Adults)	4.87 - 12.60	> 20 Years (Adults)	0.35-5.50	
	RECOM	MENDATIONS OF TSH LE	VELS DURING PREC	GNANCY (µIU/mL)		
1st Trimester				0.10 - 2.50		
2nd Trimester			0.20 - 3.00			
	3rd Trimester			0.30 - 4.10		

INCREASED TSH LEVELS:

1. Primary or untreated hypothyroidism may vary from 3 times to more than 100 times normal depending upon degree of hypofunction.

2. Hypothyroid patients receiving insufficient thyroid replacement therapy.

3.Hashimotos thyroiditis

4.DRUGS: Amphetamines, iodine containing agents & dopamine antagonist.

5.Neonatal period, increase in 1st 2-3 days of life due to post-natal surge

DECREASED TSH LEVELS:

1.Toxic multi-nodular goiter & Thyroiditis.

2. Over replacement of thyroid hormone in treatment of hypothyroidism.

3. Autonomously functioning Thyroid adenoma

4.Secondary pituitary or hypothalamic hypothyroidism

5. Acute psychiatric illness

6.Severe dehydration.

7.DRUGS: Glucocorticoids, Dopamine, Levodopa, T4 replacement therapy, Anti-thyroid drugs for thyrotoxicosis.

8. Pregnancy: 1st and 2nd Trimester



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS , MD (PATHOLOGY)



【 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mr. AJAY GUPTA				
AGE/ GENDER	: 52 YRS/MALE	PA	ATIENT ID	: 1728482	
COLLECTED BY	:	RI	EG. NO./LAB NO.	: 122501200002	
REFERRED BY	:	RI	EGISTRATION DATE	: 20/Jan/2025 08:51 AM	
BARCODE NO.	: 12506584	CC	DLLECTION DATE	: 20/Jan/2025 09:02AM	
CLIENT CODE. : P.K.R JAIN HEALTHCARE INST		TITUTE RI	EPORTING DATE	: 20/Jan/2025 01:16PM	
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AM	/IBALA CITY - HARY	ANA		
Test Name		Value	Unit	Biological Reference interval	
		CLINICAL PA	ATHOLOGY		
	URINE RO	UTINE & MICR	OSCOPIC EXAMIN	ATION	
PHYSICAL EXAMIN	NATION				
QUANTITY RECIEV		20	ml		
COLOUR	TANCE SPECTROPHOTOMETRY	PALE YELLO	W	PALE YELLOW	
	TANCE SPECTROPHOTOMETRY				
TRANSPARANCY		CLEAR		CLEAR	
by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY SPECIFIC GRAVITY		ıl PK		1.002 - 1.030	
by DIP STICK/REFLEC	TANCE SPECTROPHOTOMETRY				
<u>CHEMICAL EXAMI</u>	<u>NATION</u>				
REACTION	TANCE SPECTROPHOTOMETRY	ACIDIC			
PROTEIN	TANCE SPECIFICITIONETRY	NEGATIVE (-ve)	NEGATIVE (-ve)	
	TANCE SPECTROPHOTOMETRY				
SUGAR	TANCE SPECTROPHOTOMETRY	NEGATIVE (-ve)	NEGATIVE (-ve)	
pH		6		5.0 - 7.5	
	TANCE SPECTROPHOTOMETRY				
BILIRUBIN by DIP STICK/REFLEC	TANCE SPECTROPHOTOMETRY	NEGATIVE (-ve)	NEGATIVE (-ve)	
NITRITE		NEGATIVE (-ve)	NEGATIVE (-ve)	
by DIP STICK/REFLEC UROBILINOGEN	TANCE SPECTROPHOTOMETRY.	NOT DETEC	TED EU/dL	0.2 - 1.0	
	TANCE SPECTROPHOTOMETRY	NOT DETECT	IED EU/UL	0.2 - 1.0	
KETONE BODIES		NEGATIVE (-ve)	NEGATIVE (-ve)	
BLOOD	TANCE SPECTROPHOTOMETRY	NEGATIVE (-ve)	NEGATIVE (-ve)	
by DIP STICK/REFLEC	TANCE SPECTROPHOTOMETRY	·			
ASCORBIC ACID	TANCE SPECTROPHOTOMETRY	NEGATIVE (-ve)	NEGATIVE (-ve)	
MICROSCOPIC EXA					
RED BLOOD CELLS		NEGATIVE (-ve) /HPF	0 - 3	

DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS , MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST

NOT VALID FOR MEDICO LEGAL PURPOSE



A PIONEER DIAGNOSTIC CENTRE

【 0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

NAME	: Mr. AJAY GUPTA		
AGE/ GENDER	: 52 YRS/MALE	PATIENT ID	: 1728482
COLLECTED BY	:	REG. NO./LAB NO.	: 122501200002
REFERRED BY	:	REGISTRATION DATE	: 20/Jan/2025 08:51 AM
BARCODE NO.	: 12506584	COLLECTION DATE	: 20/Jan/2025 09:02AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	REPORTING DATE	: 20/Jan/2025 01:16PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY -	HARYANA	
Test Name	Value	Unit	Biological Reference interval

PUS CELLS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	3-4	/HPF	0 - 5
EPITHELIAL CELLS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	2-3	/HPF	ABSENT
CRYSTALS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	NEGATIVE (-ve)		NEGATIVE (-ve)
CASTS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	NEGATIVE (-ve)		NEGATIVE (-ve)
BACTERIA by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	NEGATIVE (-ve)		NEGATIVE (-ve)
OTHERS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	NEGATIVE (-ve)		NEGATIVE (-ve)
TRICHOMONAS VAGINALIS (PROTOZOA) by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	ABSENT		ABSENT

*** End Of Report



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS , MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST

