



# P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

**A PIONEER DIAGNOSTIC CENTRE**

☎ 0171-2532620, 8222896961 ✉ pkrjainhealthcare@gmail.com

TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT.

**NAME** : Mrs. SUKHVARSHA SHARMA  
**AGE/ GENDER** : 70 YRS/FEMALE  
**COLLECTED BY** :  
**REFERRED BY** :  
**BARCODE NO.** : 12506680  
**CLIENT CODE.** : P.K.R JAIN HEALTHCARE INSTITUTE  
**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**PATIENT ID** : 1734511  
**REG. NO./LAB NO.** : 122501250008  
**REGISTRATION DATE** : 25/Jan/2025 09:45 AM  
**COLLECTION DATE** : 25/Jan/2025 09:53AM  
**REPORTING DATE** : 25/Jan/2025 04:36PM

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

## CLINICAL CHEMISTRY/BIOCHEMISTRY

### IRON PROFILE

IRON: SERUM by FERROZINE, SPECTROPHOTOMETRY	19.7 <sup>L</sup>	µg/dL	37.0 - 145.0
UNSATURATED IRON BINDING CAPACITY (UIBC):SERUM by FERROZINE, SPECTROPHOTOMETRY	358.06 <sup>H</sup>	µg/dL	150.0 - 336.0
TOTAL IRON BINDING CAPACITY (TIBC):SERUM by SPECTROPHOTOMETRY	377.76	µg/dL	230 - 430
%TRANSFERRIN SATURATION: SERUM by CALCULATED, SPECTROPHOTOMETRY (FERENE)	5.21 <sup>L</sup>	%	15.0 - 50.0
TRANSFERRIN: SERUM by SPECTROPHOTOMETRY (FERENE)	268.21	mg/dL	200.0 - 350.0

#### INTERPRETATION:-

VARIABLES	ANEMIA OF CHRONIC DISEASE	IRON DEFICIENCY ANEMIA	THALASSEMIA α/β TRAIT
SERUM IRON:	Normal to Reduced	Reduced	Normal
TOTAL IRON BINDING CAPACITY:	Decreased	Increased	Normal
% TRANSFERRIN SATURATION:	Decreased	Decreased < 12-15 %	Normal
SERUM FERRITIN:	Normal to Increased	Decreased	Normal or Increased

#### IRON:

1.Serum iron studies is recommended for differential diagnosis of microcytic hypochromic anemia.i.e iron deficiency anemia, zinc deficiency anemia,anemia of chronic disease and thalassemia syndromes.  
2. It is essential to isolate iron deficiency anemia from Beta thalassemia syndromes because during iron replacement which is therapeutic for iron deficiency anemia, is severely contra-indicated in Thalassemia.


#### TOTAL IRON BINDING CAPACITY (TIBC):


1.It is a direct measure of protein transferrin which transports iron from the gut to storage sites in the bone marrow.

#### % TRANSFERRIN SATURATION:

1.Occurs in idiopathic hemochromatosis and transfusional hemosiderosis where no unsaturated iron binding capacity is available for iron mobilization. Similar condition is seen in congenital deficiency of transferrin.



  
DR.VINAY CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

  
DR.YUGAM CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY)





# P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

**A PIONEER DIAGNOSTIC CENTRE**

☎ 0171-2532620, 8222896961 ✉ pkrjainhealthcare@gmail.com

**NAME** : Mrs. SUKHVARSHA SHARMA  
**AGE/ GENDER** : 70 YRS/FEMALE  
**COLLECTED BY** :  
**REFERRED BY** :  
**BARCODE NO.** : 12506680  
**CLIENT CODE.** : P.K.R JAIN HEALTHCARE INSTITUTE  
**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**PATIENT ID** : 1734511  
**REG. NO./LAB NO.** : 122501250008  
**REGISTRATION DATE** : 25/Jan/2025 09:45 AM  
**COLLECTION DATE** : 25/Jan/2025 09:53AM  
**REPORTING DATE** : 25/Jan/2025 04:36PM

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

## VITAMINS

### VITAMIN B12/COBALAMIN


VITAMIN B12/COBALAMIN: SERUM **146<sup>L</sup>** pg/mL 200.0 - 1100.0  
by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)


#### INTERPRETATION:-

INCREASED VITAMIN B12	DECREASED VITAMIN B12
1.Ingestion of Vitamin C	1.Pregnancy
2.Ingestion of Estrogen	2.DRUGS:Aspirin, Anti-convulsants, Colchicine
3.Ingestion of Vitamin A	3.Ethanol lgestion
4.Hepatocellular injury	4. Contraceptive Harmones
5.Myeloproliferative disorder	5.Haemodialysis
6.Uremia	6. Multiple Myeloma

1.Vitamin B12 (cobalamin) is necessary for hematopoiesis and normal neuronal function.  
2.In humans, it is obtained only from animal proteins and requires intrinsic factor (IF) for absorption.  
3.The body uses its vitamin B12 stores very economically, reabsorbing vitamin B12 from the ileum and returning it to the liver; very little is excreted.  
4.Vitamin B12 deficiency may be due to lack of IF secretion by gastric mucosa (eg, gastrectomy, gastric atrophy) or intestinal malabsorption (eg, ileal resection, small intestinal diseases).  
5.Vitamin B12 deficiency frequently causes macrocytic anemia, glossitis, peripheral neuropathy, weakness, hyperreflexia, ataxia, loss of proprioception, poor coordination, and affective behavioral changes. These manifestations may occur in any combination; many patients have the neurologic defects without macrocytic anemia.  
6.Serum methylmalonic acid and homocysteine levels are also elevated in vitamin B12 deficiency states.  
7.Follow-up testing for antibodies to intrinsic factor (IF) is recommended to identify this potential cause of vitamin B12 malabsorption.  
**NOTE:**A normal serum concentration of vitamin B12 does not rule out tissue deficiency of vitamin B12. The most sensitive test for vitamin B12 deficiency at the cellular level is the assay for MMA. If clinical symptoms suggest deficiency, measurement of MMA and homocysteine should be considered, even if serum vitamin B12 concentrations are normal.



  
DR.VINAY CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

  
DR.YUGAM CHOPRA  
CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY)





# P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

**A PIONEER DIAGNOSTIC CENTRE**

☎ 0171-2532620, 8222896961 ✉ pkrajainhealthcare@gmail.com

TEST PERFORMED AT: KOS DIAGNOSTIC LAB, AMBALA CANTT.

**NAME** : Mrs. SUKHVARSHA SHARMA  
**AGE/ GENDER** : 70 YRS/FEMALE  
**COLLECTED BY** :  
**REFERRED BY** :  
**BARCODE NO.** : 12506680  
**CLIENT CODE.** : P.K.R JAIN HEALTHCARE INSTITUTE  
**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**PATIENT ID** : 1734511  
**REG. NO./LAB NO.** : 122501250008  
**REGISTRATION DATE** : 25/Jan/2025 09:45 AM  
**COLLECTION DATE** : 25/Jan/2025 09:53AM  
**REPORTING DATE** : 25/Jan/2025 05:11PM

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

## VITAMIN B9/FOLIC ACID/FOLATE

VITAMIN B9/FOLIC ACID/FOLATE: SERUM  
by CLIA (CHEMILUMINESCENCE IMMUNOASSAY)

11.8 ng/mL

DEFICIENT: < 3.37  
INTERMEDIATE: 3.37 - 5.38  
NORMAL: > 5.38

### INTERPRETATION

RESULT IN ng/mL	REMARKS
0.35 – 3.37	DEFICIENT
3.38 – 5.38	INTERMEDIATE
5.39 – 100.00	NORMAL

### NOTE:

1. Drugs like Methotrexate & Leucovorin interfere with folate measurement
2. To differentiate vitamin B12 & folate deficiency, measurement of Methyl malonic acid in urine & serum Homocysteine level is suggested
3. Risk of toxicity from folic acid is low as it is a water soluble vitamin regularly excreted in urine

### COMMENTS:

1. Folate plays an important role in the synthesis of purine & pyrimidines in the body and is important for the maturation of erythrocytes.
2. It is widely available from plants and to a lesser extent organ meats, but more than half the folate content of food is lost during cooking.
3. Folate deficiency is commonly prevalent in alcoholic liver disease, pregnancy and the elderly. It may result from poor intestinal absorption, nutrition deficiency, excessive demand as in pregnancy or in malignancy and in response to certain drugs like Methotrexate & anticonvulsants.
4. Decreased Levels Megaloblastic anemia, Infantile hyperthyroidism, Alcoholism, Malnutrition, Scurvy, Liver disease, B12 deficiency, dietary amino acid excess, adult Celiac disease, Tropical Sprue, Crohn's disease, Hemolytic anemias, Carcinomas, Myelofibrosis, vitamin B6 deficiency, pregnancy, Whipple's disease, extensive intestinal resection and severe exfoliative dermatitis

\*\*\* End Of Report \*\*\*



  
DR.VINAY CHOPRA

CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

  
DR.YUGAM CHOPRA

CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY)

