

### A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

**NAME** : Mr. CHAMAN LAL

**AGE/ GENDER** : 55 YRS/MALE **PATIENT ID** : 1735998

**COLLECTED BY** REG. NO./LAB NO. : 122501270001

REFERRED BY **REGISTRATION DATE** : 27/Jan/2025 08:33 AM BARCODE NO. : 12506697 **COLLECTION DATE** : 27/Jan/2025 09:53AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 27/Jan/2025 01:03PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**Value** Unit **Biological Reference interval Test Name** 

### **SWASTHYA WELLNESS PANEL: 1.2 COMPLETE BLOOD COUNT (CBC)**

### **RED BLOOD CELLS (RBCS) COUNT AND INDICES**

HAEMOGLOBIN (HB) by CALORIMETRIC	16.7	gm/dL	12.0 - 17.0
RED BLOOD CELL (RBC) COUNT by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	5.03 <sup>H</sup>	Millions/cmm	3.50 - 5.00
PACKED CELL VOLUME (PCV) by calculated by automated hematology analyzer	47.9	%	40.0 - 54.0
MEAN CORPUSCULAR VOLUME (MCV) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	95.1	fL	80.0 - 100.0
MEAN CORPUSCULAR HAEMOGLOBIN (MCH) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	33.1	pg	27.0 - 34.0
MEAN CORPUSCULAR HEMOGLOBIN CONC. (MCHC) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	34.8	g/dL	32.0 - 36.0
RED CELL DISTRIBUTION WIDTH (RDW-CV) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	14.4	%	11.00 - 16.00
RED CELL DISTRIBUTION WIDTH (RDW-SD) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	51.9	fL	35.0 - 56.0
MENTZERS INDEX by CALCULATED	18.91	RATIO	BETA THALASSEMIA TRAIT: < 13.0 IRON DEFICIENCY ANEMIA: >13.0
GREEN & KING INDEX by CALCULATED	27.14	RATIO	BETA THALASSEMIA TRAIT:<= 65.0 IRON DEFICIENCY ANEMIA: > 65.0
WHITE BLOOD CELLS (WBCS)			
TOTAL LEUCOCYTE COUNT (TLC) by flow cytometry by sf cube & microscopy	6890	/cmm	4000 - 11000
<b>DIFFERENTIAL LEUCOCYTE COUNT (DLC)</b>			
NEUTROPHILS by flow cytometry by sf cube & microscopy	64	%	50 - 70
LYMPHOCYTES	28	%	20 - 40



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST







### A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

NAME : Mr. CHAMAN LAL

**AGE/ GENDER** : 55 YRS/MALE **PATIENT ID** : 1735998

**COLLECTED BY** REG. NO./LAB NO. : 122501270001

**REFERRED BY REGISTRATION DATE** : 27/Jan/2025 08:33 AM BARCODE NO. : 12506697 **COLLECTION DATE** : 27/Jan/2025 09:53AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 27/Jan/2025 01:03PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name	Value	Unit	Biological Reference interval
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY			
EOSINOPHILS	1	%	1 - 6
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY			
MONOCYTES	7	%	2 - 12
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY BASOPHILS	0	0/	0 - 1
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	0	%	0-1
ABSOLUTE LEUKOCYTES (WBC) COUNT			
ABSOLUTE NEUTROPHIL COUNT	4410	/cmm	2000 - 7500
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY			
ABSOLUTE LYMPHOCYTE COUNT	1929 <sup>L</sup>	/cmm	800 - 4900
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY			
ABSOLUTE EOSINOPHIL COUNT	69	/cmm	40 - 440
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY  ABSOLUTE MONOCYTE COUNT	482	/amm	80 - 880
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	402	/cmm	80 - 880
ABSOLUTE BASOPHIL COUNT	0	/cmm	0 - 110
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY			
PLATELETS AND OTHER PLATELET PREDICTIVE	MARKERS.		
PLATELET COUNT (PLT)	172000	/cmm	150000 - 450000
by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE			
PLATELETCRIT (PCT)	0.16	%	0.10 - 0.36
by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE		CT.	0.50 10.0
MEAN PLATELET VOLUME (MPV) by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	9	fL	6.50 - 12.0
PLATELET LARGE CELL COUNT (P-LCC)	41000	/cmm	30000 - 90000
by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	41000	/ ciiiii	30000 30000
PLATELET LARGE CELL RATIO (P-LCR)	23.9	%	11.0 - 45.0
by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE			
PLATELET DISTRIBUTION WIDTH (PDW)	16.4	%	15.0 - 17.0
by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE			
NOTE: TEST CONDUCTED ON EDTA WHOLE BLOOD			



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS, MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST







### A PIONEER DIAGNOSTIC CENTRE

: 27/Jan/2025 01:03PM

**NAME** : Mr. CHAMAN LAL

AGE/ GENDER : 55 YRS/MALE **PATIENT ID** : 1735998

**COLLECTED BY** REG. NO./LAB NO. : 122501270001

REFERRED BY **REGISTRATION DATE** : 27/Jan/2025 08:33 AM BARCODE NO. **COLLECTION DATE** : 27/Jan/2025 09:53AM : 12506697

: P.K.R JAIN HEALTHCARE INSTITUTE **CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**Value** Unit **Test Name Biological Reference interval** 

### **ERYTHROCYTE SEDIMENTATION RATE (ESR)**

ERYTHROCYTE SEDIMENTATION RATE (ESR)

mm/1st hr 0 - 20

REPORTING DATE

by RED CELL AGGREGATION BY CAPILLARY PHOTOMETRY

#### INTERPRETATION:

CLIENT CODE.

- 1. ESR is a non-specific test because an elevated result often indicates the presence of inflammation associated with infection, cancer and auto-immune disease, but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it.
- 2. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other test such as C-reactive protein
- 3. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as some others, such as systemic lupus erythematosus

### CONDITION WITH LOW ESR

A low ESR can be seen with conditions that inhibit the normal sedimentation of red blood cells, such as a high red blood cell count (polycythaemia), significantly high white blood cell count (leucocytosis), and some protein abnormalities. Some changes in red cell shape (such as sickle cells in sickle cell anaemia) also lower the ESR.

### NOTE:

- 1. ESR and C reactive protein (C-RP) are both markers of inflammation.
- 2. Generally, ESR does not change as rapidly as does CRP, either at the start of inflammation or as it resolves.
   3. CRP is not affected by as many other factors as is ESR, making it a better marker of inflammation.
   4. If the ESR is elevated, it is typically a result of two types of proteins, globulins or fibringen.
   5. Women tend to average mathyldone and entraceptives professional processing mathyldone and with the opposition of the oppositio

- 6. Drugs such as dextran, methyldopa, oral contraceptives, penicillamine procainamide, theophylline, and vitamin A can increase ESR, while aspirin, cortisone, and quinine may decrease it



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)







### A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

**NAME** : Mr. CHAMAN LAL

AGE/ GENDER : 55 YRS/MALE **PATIENT ID** : 1735998

**COLLECTED BY** REG. NO./LAB NO. : 122501270001

REFERRED BY **REGISTRATION DATE** : 27/Jan/2025 08:33 AM BARCODE NO. : 12506697 **COLLECTION DATE** : 27/Jan/2025 09:53AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 27/Jan/2025 12:42PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**Value** Unit **Biological Reference interval Test Name** 

### CLINICAL CHEMISTRY/BIOCHEMISTRY **GLUCOSE FASTING (F)**

GLUCOSE FASTING (F): PLASMA NORMAL: < 100.0  $111.06^{H}$ mg/dL

by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD) PREDIABETIC: 100.0 - 125.0

DIABETIC: > 0R = 126.0

INTERPRETATION
IN ACCORDANCE WITH AMERICAN DIABETES ASSOCIATION GUIDELINES:

1. A fasting plasma glucose level below 100 mg/dl is considered normal.

2. A fasting plasma glucose level between 100 - 125 mg/dl is considered as glucose intolerant or prediabetic. A fasting and post-prandial blood

test (after consumption of 75 gms of glucose) is recommended for all such patients.

3. A fasting plasma glucose level of above 125 mg/dl is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for all such patients. A fasting plasma glucose level in excess of 125 mg/dl on both occasions is confirmatory for diabetic state.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)





### A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

**NAME** : Mr. CHAMAN LAL

**AGE/ GENDER** : 55 YRS/MALE **PATIENT ID** : 1735998

**COLLECTED BY** REG. NO./LAB NO. : 122501270001

REFERRED BY **REGISTRATION DATE** : 27/Jan/2025 08:33 AM BARCODE NO. : 12506697 **COLLECTION DATE** : 27/Jan/2025 09:53AM

CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 27/Jan/2025 01:03PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name	Value	Unit	Biological Reference interval
	LIPID PROFILE : BA	SIC	
CHOLESTEROL TOTAL: SERUM by CHOLESTEROL OXIDASE PAP	190.23	mg/dL	OPTIMAL: < 200.0 BORDERLINE HIGH: 200.0 - 239.0 HIGH CHOLESTEROL: > OR = 240.0
TRIGLYCERIDES: SERUM by GLYCEROL PHOSPHATE OXIDASE (ENZYMATIC)	406.96 <sup>H</sup>	mg/dL	OPTIMAL: < 150.0 BORDERLINE HIGH: 150.0 - 199.0 HIGH: 200.0 - 499.0 VERY HIGH: > OR = 500.0
HDL CHOLESTEROL (DIRECT): SERUM by SELECTIVE INHIBITION	35.01	mg/dL	LOW HDL: < 30.0 BORDERLINE HIGH HDL: 30.0 - 60.0 HIGH HDL: > OR = 60.0
LDL CHOLESTEROL: SERUM by CALCULATED, SPECTROPHOTOMETRY	NOT CALCULATED	mg/dL	OPTIMAL: < 100.0 ABOVE OPTIMAL: 100.0 - 129.0 BORDERLINE HIGH: 130.0 - 159.0 HIGH: 160.0 - 189.0 VERY HIGH: > OR = 190.0
NON HDL CHOLESTEROL: SERUM by CALCULATED, SPECTROPHOTOMETRY	155.22 <sup>H</sup>	mg/dL	OPTIMAL: < 130.0 ABOVE OPTIMAL: 130.0 - 159.0 BORDERLINE HIGH: 160.0 - 189.0 HIGH: 190.0 - 219.0 VERY HIGH: > OR = 220.0
VLDL CHOLESTEROL: SERUM by CALCULATED, SPECTROPHOTOMETRY	NOT CALCULATED	mg/dL	0.00 - 45.00
TOTAL LIPIDS: SERUM by CALCULATED, SPECTROPHOTOMETRY	NOT CALCULATED	mg/dL	350.00 - 700.00
CHOLESTEROL/HDL RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	5.43 <sup>H</sup>	RATIO	LOW RISK: 3.30 - 4.40 AVERAGE RISK: 4.50 - 7.0 MODERATE RISK: 7.10 - 11.0



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



HIGH RISK: > 11.0





### A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

**NAME** : Mr. CHAMAN LAL

AGE/ GENDER : 55 YRS/MALE **PATIENT ID** : 1735998

**COLLECTED BY** REG. NO./LAB NO. : 122501270001

REFERRED BY **REGISTRATION DATE** : 27/Jan/2025 08:33 AM BARCODE NO. **COLLECTION DATE** : 27/Jan/2025 09:53AM : 12506697 CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 27/Jan/2025 01:03PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name	Value	Unit	Biological Reference interval
LDL/HDL RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	NOT CALCULATED	RATIO	LOW RISK: 0.50 - 3.0 MODERATE RISK: 3.10 - 6.0 HIGH RISK: > 6.0
TRIGLYCERIDES/HDL RATIO: SERUM	11.62 <sup>H</sup>	RATIO	3.00 - 5.00

by CALCULATED, SPECTROPHOTOMETRY

WHEN TRIGLYCERIDES VALUE >400 mg/dL THE CALCULATED VALUES OF LDL AND VLDL ARE NOT RELIABLE

#### **INTERPRETATION:**

NOTE 2

1. Measurements in the same patient can show physiological & analytical variations. Three serial samples 1 week apart are recommended for Total Cholesterol, Triglycerides, HDL & LDL Cholesterol.

2. As per NLA-2014 guidelines, all adults above the age of 20 years should be screened for lipid status. Selective screening of children above the age of 2 years with a family history of premature cardiovascular disease or those with at least one parent with high total cholesterol is recommended.

3. Low HDL levels are associated with increased risk for Atherosclerotic Cardiovascular disease (ASCVD) due to insufficient HDL being available

to participate in reverse cholesterol transport, the process by which cholesterol is eliminated from peripheral tissues.

4. NLA-2014 identifies Non HDL Cholesterol (an indicator of all atherogeniclipoproteins such as LDL, VLDL, IDL, Lpa, Chylomicron remnants) along with LDL-cholesterol as co- primary target for cholesterol lowering therapy. Note that major risk factors can modify treatment goals for LDL &Non

5. Additional testing for Apolipoprotein B, hsCRP,Lp(a) & LP-PLA2 should be considered among patients with moderate risk for ASCVD for risk refinement



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)





### A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

**NAME** : Mr. CHAMAN LAL

**AGE/ GENDER** : 55 YRS/MALE **PATIENT ID** : 1735998

**COLLECTED BY** : 122501270001 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 27/Jan/2025 08:33 AM BARCODE NO. : 12506697 **COLLECTION DATE** : 27/Jan/2025 09:53AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 27/Jan/2025 01:03PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**Test Name Value** Unit **Biological Reference interval** 

### **LIVER FUNCTION TEST (COMPLETE)**

BILIRUBIN TOTAL: SERUM by DIAZOTIZATION, SPECTROPHOTOMETRY	0.85	mg/dL	INFANT: 0.20 - 8.00 ADULT: 0.00 - 1.20
BILIRUBIN DIRECT (CONJUGATED): SERUM by DIAZO MODIFIED, SPECTROPHOTOMETRY	0.25	mg/dL	0.00 - 0.40
BILIRUBIN INDIRECT (UNCONJUGATED): SERUM by CALCULATED, SPECTROPHOTOMETRY	0.6	mg/dL	0.10 - 1.00
SGOT/AST: SERUM by IFCC, WITHOUT PYRIDOXAL PHOSPHATE	19.94	U/L	7.00 - 45.00
SGPT/ALT: SERUM by IFCC, WITHOUT PYRIDOXAL PHOSPHATE	14.48	U/L	0.00 - 49.00
AST/ALT RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	1.38	RATIO	0.00 - 46.00
ALKALINE PHOSPHATASE: SERUM by PARA NITROPHENYL PHOSPHATASE BY AMINO METHYL PROPANOL	82.88	U/L	40.0 - 130.0
GAMMA GLUTAMYL TRANSFERASE (GGT): SERUM by SZASZ, SPECTROPHTOMETRY	21.54	U/L	0.00 - 55.0
TOTAL PROTEINS: SERUM by BIURET, SPECTROPHOTOMETRY	6.14 <sup>L</sup>	gm/dL	6.20 - 8.00
ALBUMIN: SERUM by BROMOCRESOL GREEN	4.43	gm/dL	3.50 - 5.50
GLOBULIN: SERUM by CALCULATED, SPECTROPHOTOMETRY	1.71 <sup>L</sup>	gm/dL	2.30 - 3.50
A : G RATIO: SERUM	2.59 <sup>H</sup>	RATIO	1.00 - 2.00

#### INTERPRETATION

by CALCULATED, SPECTROPHOTOMETRY

NOTE:- To be correlated in individuals having SGOT and SGPT values higher than Normal Referance Range.

**USE**:- Differential diagnosis of diseases of hepatobiliary system and pancreas.

### INCREASED:

DRUG HEPATOTOXICITY	> 2
ALCOHOLIC HEPATITIS	> 2 (Highly Suggestive)
CIRRHOSIS	1.4 - 2.0
INTRAHEPATIC CHOLESTATIS	> 1.5
HEPATOCELLULAR CARCINOMA & CHRONIC HEPATITIS	> 1.3 (Slightly Increased)



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS, MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST







### A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

REPORTING DATE

: 27/Jan/2025 01:03PM

**NAME** : Mr. CHAMAN LAL

**AGE/ GENDER** : 55 YRS/MALE **PATIENT ID** : 1735998

**COLLECTED BY** REG. NO./LAB NO. : 122501270001

REFERRED BY **REGISTRATION DATE** : 27/Jan/2025 08:33 AM BARCODE NO. : 12506697 **COLLECTION DATE** : 27/Jan/2025 09:53AM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

**Test Name Value** Unit **Biological Reference interval** 

#### **DECREASED:**

CLIENT CODE.

1. Acute Hepatitis due to virus, drugs, toxins (with AST increased 3 to 10 times upper limit of normal)

2. Extra Hepatic cholestatis: 0.8 (normal or slightly decreased).

#### PROGNOSTIC SIGNIFICANCE:

NORMAL	< 0.65
GOOD PROGNOSTIC SIGN	0.3 - 0.6
POOR PROGNOSTIC SIGN	1.2 - 1.6



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)





### PKR JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

### A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

2.30 - 4.70

**NAME** : Mr. CHAMAN LAL

**AGE/ GENDER** : 55 YRS/MALE **PATIENT ID** : 1735998

**COLLECTED BY** : 122501270001 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 27/Jan/2025 08:33 AM BARCODE NO. : 12506697 **COLLECTION DATE** : 27/Jan/2025 09:53AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 27/Jan/2025 04:59PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name	Value	Unit	<b>Biological Reference interval</b>
WIDNE	v elikopioki pece	(COMPLETE)	
KIDNE	Y FUNCTION TEST	(COMPLETE)	
UREA: SERUM by UREASE - GLUTAMATE DEHYDROGENASE (GLDH)	21.5	mg/dL	10.00 - 50.00
CREATININE: SERUM by ENZYMATIC, SPECTROPHOTOMETERY	0.91	mg/dL	0.40 - 1.40
BLOOD UREA NITROGEN (BUN): SERUM by CALCULATED, SPECTROPHOTOMETRY	10.05	mg/dL	7.0 - 25.0
BLOOD UREA NITROGEN (BUN)/CREATININE RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	11.04	RATIO	10.0 - 20.0
UREA/CREATININE RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	23.63	RATIO	
URIC ACID: SERUM by URICASE - OXIDASE PEROXIDASE	3.88	mg/dL	3.60 - 7.70
CALCIUM: SERUM by ARSENAZO III, SPECTROPHOTOMETRY	9.27	mg/dL	8.50 - 10.60

PHOSPHOROUS: SERUM

<b>ELECTROLYTES</b>			
SODIUM: SERUM by ISE (ION SELECTIVE ELECTRODE)	141.1	mmol/L	135.0 - 150.0
POTASSIUM: SERUM by ISE (ION SELECTIVE ELECTRODE)	4.43	mmol/L	3.50 - 5.00
CHLORIDE: SERUM by ISE (ION SELECTIVE ELECTRODE)	105.82	mmol/L	90.0 - 110.0

### **ESTIMATED GLOMERULAR FILTERATION RATE**

by PHOSPHOMOLYBDATE, SPECTROPHOTOMETRY

ESTIMATED GLOMERULAR FILTERATION RATE 99.5

(eGFR): SERUM by CALCULATED **INTERPRETATION:** 

To differentiate between pre- and post renal azotemia.

#### INCREASED RATIO (>20:1) WITH NORMAL CREATININE:

- 1. Prerenal azotemia (BUN rises without increase in creatinine) e.g. heart failure, salt depletion, dehydration, blood loss) due to decreased glomerular filtration rate.
- Catabolic states with increased tissue breakdown.
- 3. GI haemorrhage.



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)





### A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

**NAME** : Mr. CHAMAN LAL

AGE/ GENDER : 55 YRS/MALE **PATIENT ID** : 1735998

**COLLECTED BY** REG. NO./LAB NO. : 122501270001

REFERRED BY **REGISTRATION DATE** : 27/Jan/2025 08:33 AM BARCODE NO. **COLLECTION DATE** : 27/Jan/2025 09:53AM : 12506697 CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 27/Jan/2025 04:59PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**Test Name Value** Unit **Biological Reference interval** 

4. High protein intake.

5. Impaired renal function plus

6. Excess protein intake or production or tissue breakdown (e.g. infection, GI bleeding, thyrotoxicosis, Cushing's syndrome, high protein diet, burns, surgery, cachexia, high fever).

7. Urine reabsorption (e.g. ureter colostomy)

8. Reduced muscle mass (subnormal creatinine production)

9. Certain drugs (e.g. tetracycline, glucocorticoids)

### INCREASED RATIO (>20:1) WITH ELEVATED CREATININE LEVELS:

- 1. Postrenal azotemia (BUN rises disproportionately more than creatinine) (e.g. obstructive uropathy).
- 2. Prerenal azotemia superimposed on renal disease.

### DECREASED RATIO (<10:1) WITH DECREASED BUN:

- 1. Acute tubular necrosis.
- 2. Low protein diet and starvation.
- 3. Severe liver disease.
- 4. Other causes of decreased urea synthesis.
- 5. Repeated dialysis (urea rather than creatinine diffuses out of extracellular fluid).
- 6. Inherited hyperammonemias (urea is virtually absent in blood).
- 7. SIADH (syndrome of inappropiate antidiuretic harmone) due to tubular secretion of urea.
- 8. Pregnancy.

### **DECREASED RATIO (<10:1) WITH INCREASED CREATININE:**

- 1. Phenacimide therapy (accelerates conversion of creatine to creatinine).
- 2. Rhabdomyolysis (releases muscle creatinine).
- 3. Muscular patients who develop renal failure.

### **INAPPROPIATE RATIO:**

1. Diabetic ketoacidosis (acetoacetate causes false increase in creatinine with certain methodologies, resulting in normal ratio when dehydration should produce an increased BUN/creatinine ratio).

2. Cephalosporin therapy (interferes with creatinine measurement). **ESTIMATED GLOMERULAR FILTERATION RATE**:

ESTIMINATED SESMENOETH TETERATION INTE.				
CKD STAGE	DESCRIPTION	GFR ( mL/min/1.73m2 )	ASSOCIATED FINDINGS	
G1	Normal kidney function	>90	No proteinuria	
G2	Kidney damage with	>90	Presence of Protein,	
	normal or high GFR		Albumin or cast in urine	
G3a	Mild decrease in GFR	60 -89		
G3b	Moderate decrease in GFR	30-59		
G4	Severe decrease in GFR	15-29		
G5	Kidney failure	<15		



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)







### A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

REPORTING DATE

: 27/Jan/2025 04:59PM

**NAME** : Mr. CHAMAN LAL

AGE/ GENDER : 55 YRS/MALE **PATIENT ID** : 1735998

**COLLECTED BY** REG. NO./LAB NO. : 122501270001

REFERRED BY **REGISTRATION DATE** : 27/Jan/2025 08:33 AM BARCODE NO. **COLLECTION DATE** : 27/Jan/2025 09:53AM : 12506697

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

**Test Name Value** Unit **Biological Reference interval** 

COMMENTS:

CLIENT CODE.

1. Estimated Glomerular filtration rate (eGFR) is the sum of filtration rates in all functioning nephrons and so an estimation of the GFR provides a measure of functioning nephrons of the kidney.

2. eGFR calculated using the 2009 CKD-EPI creatinine equation and GFR category reported as per KDIGO guideline 2012

3. In patients, with eGFR creatinine between 45-59 ml/min/1.73 m2 (G3) and without any marker of Kidney damage, It is recommended to measure eGFR with Cystatin C for confirmation of CKD

4. eGFR category G1 OR G2 does not fullfill the criteria for CKD, in the absence of evidence of Kidney Damage
5. In a suspected case of Acute Kidney Injury (AKI), measurement of eGFR should be done after 48-96 hours of any Intervention or procedure
6. eGFR calculated by Serum Creatinine may be less accurate due to certain factors like Race, Muscle Mass, Diet, Certain Drugs. In such cases, eGFR should be calculated using Serum Cystatin C
7. A decrease in eGFR implies either progressive renal disease, or a reversible process causing decreased nephron function (eg, severe dehydration).

KDIGO guideline, 2012 recommends Chronic Kidney Disease (CKD) should be classified based on cause, eGFR category and Albuminuria (ACR) category. GFR & ACR category combined together reflect risk of progression and helps Clinician to identify the individual who are progressing at more rapid rate than anticipated



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)







### A PIONEER DIAGNOSTIC CENTRE

: 27/Jan/2025 01:03PM

**NAME** : Mr. CHAMAN LAL

AGE/ GENDER : 55 YRS/MALE **PATIENT ID** : 1735998

**COLLECTED BY** : 122501270001 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 27/Jan/2025 08:33 AM BARCODE NO. : 12506697 **COLLECTION DATE** : 27/Jan/2025 09:53AM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

**Value** Unit **Biological Reference interval Test Name** 

REPORTING DATE

### **ENDOCRINOLOGY**

### THYROID FUNCTION TEST: TOTAL

TRIIODOTHYRONINE (T3): SERUM 0.35 - 1.931.21 ng/mL by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

THYROXINE (T4): SERUM 5.78 μgm/dL 4.87 - 12.60

by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY) THYROID STIMULATING HORMONE (TSH): SERUM 2.57 μIU/mL 0.35 - 5.50

by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

3rd GENERATION, ULTRASENSITIVE

#### INTERPRETATION:

CLIENT CODE.

TSH levels are subject to circadian variation, reaching peak levels between 2-4 a.m and at a minimum between 6-10 pm. The variation is of the order of 50%. Hence time of the day has influence on the measured serum TSH concentrations. TSH stimulates the production and secretion of the metabolically active hormones, thyroxine (T4) and triliodothyronine (T3). Failure at any level of regulation of the hypothalamic-pituitary-thyroid axis will result in either underproduction (hypothyroidism) or overproduction(hyperthyroidism) of T4 and/or T3.

CLINICAL CONDITION	Т3	T4	TSH
Primary Hypothyroidism:	Reduced	Reduced	Increased (Significantly)
Subclinical Hypothyroidism:	Normal or Low Normal	Normal or Low Normal	High
Primary Hyperthyroidism:	Increased	Increased	Reduced (at times undetectable)
Subclinical Hyperthyroidism:	Normal or High Normal	Normal or High Normal	Reduced

- 1. T3 and T4 circulates in reversibly bound form with Thyroid binding globulins (TBG), and to a lesser extent albumin and Thyroid binding Pre Albumin so conditions in which TBG and protein levels alter such as pregnancy, excess estrogens, androgens, anabolic steroids and glucocorticoids may falsely affect the T3 and T4 levels and may cause false thyroid values for thyroid function tests
- 2. Normal levels of T4 can also be seen in Hyperthyroid patients with :T3 Thyrotoxicosis, Decreased binding capacity due to hypoproteinemia or ingestion of certain drugs
- 3. Serum T4 levels in neonates and infants are higher than values in the normal adult, due to the increased concentration of TBG in neonate serum.
- 4. TSH may be normal in central hypothyroidism, recent rapid correction of hyperthyroidism or hypothyroidism, pregnancy, phenytoin therapy.

TRIIODOTH	(RONINE (T3)	THYROX	INE (T4)	THYROID STIMUI	ATING HORMONE (TSH)
Age	Refferance Range (ng/mL)	Age	Refferance Range (µg/dL)	Age	Reference Range ( μΙυ/mL)
0 - 7 Days	0.20 - 2.65	0 - 7 Days	5.90 - 18.58	0 - 7 Days	2.43 - 24.3
7 Days - 3 Months	0.36 - 2.59	7 Days - 3 Months	6.39 - 17.66	7 Days - 3 Months	0.58 - 11.00
3 - 6 Months	0.51 - 2.52	3 - 6 Months	6.75 – 17.04	3 Days – 6 Months	0.70 - 8.40
6 - 12 Months	0.74 - 2.40	6 - 12 Months	7.10 – 16.16	6 – 12 Months	0.70 - 7.00



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)







### A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

**NAME** : Mr. CHAMAN LAL

AGE/ GENDER : 55 YRS/MALE **PATIENT ID** : 1735998

**COLLECTED BY** REG. NO./LAB NO. : 122501270001

REFERRED BY **REGISTRATION DATE** : 27/Jan/2025 08:33 AM BARCODE NO. : 12506697 **COLLECTION DATE** : 27/Jan/2025 09:53AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 27/Jan/2025 01:03PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name		Value	Unit	ţ	Biological Reference interval	
1 - 10 Years	0.92 - 2.28	1 - 10 Years	6.00 - 13.80	1 – 10 Years	0.60 - 5.50	
11- 19 Years	0.35 - 1.93	11 - 19 Years	4.87- 13.20	11 – 19 Years	0.50 - 5.50	
> 20 years (Adults)	0.35 - 1.93	> 20 Years (Adults)	4.87 - 12.60	> 20 Years (Adults)	0.35- 5.50	
	RECON	MMENDATIONS OF TSH LI	EVELS DURING PRE	GNANCY ( µIU/mL)		
1st Trimester			0.10 - 2.50			
2nd Trimester			0.20 - 3.00			
3rd Trimester			0.30 - 4.10			

#### **INCREASED TSH LEVELS:**

- 1. Primary or untreated hypothyroidism may vary from 3 times to more than 100 times normal depending upon degree of hypofunction.
- 2. Hypothyroid patients receiving insufficient thyroid replacement therapy.
- 3. Hashimotos thyroiditis
- 4.DRUGS: Amphetamines, iodine containing agents & dopamine antagonist.
- 5. Neonatal period, increase in 1st 2-3 days of life due to post-natal surge

### **DECREASED TSH LEVELS:**

- 1. Toxic multi-nodular goiter & Thyroiditis.
- 2. Over replacement of thyroid hormone in treatment of hypothyroidism.
- 3. Autonomously functioning Thyroid adenoma
- 4. Secondary pituitary or hypothalamic hypothyroidism
- 5. Acute psychiatric illness
- 6. Severe dehydration.
- 7.DRUGS: Glucocorticoids, Dopamine, Levodopa, T4 replacement therapy, Anti-thyroid drugs for thyrotoxicosis.

8. Pregnancy: 1st and 2nd Trimester



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)







### A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 pkrjainhealthcare@gmail.com

**NAME** : Mr. CHAMAN LAL

AGE/ GENDER : 55 YRS/MALE **PATIENT ID** : 1735998

**COLLECTED BY** REG. NO./LAB NO. : 122501270001

REFERRED BY **REGISTRATION DATE** : 27/Jan/2025 08:33 AM BARCODE NO. **COLLECTION DATE** : 27/Jan/2025 09:53AM : 12506697 CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 27/Jan/2025 04:59PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**Value** Unit **Biological Reference interval Test Name** 

### IMMUNOPATHOLOGY/SEROLOGY RHEUMATOID FACTOR (RA): QUANTITATIVE - SERUM

RHEUMATOID (RA) FACTOR QUANTITATIVE: 10.4 IU/mL NEGATIVE: < 18.0

**SERUM** BORDERLINE: 18.0 - 25.0

by NEPHLOMETRY POSITIVE: > 25.0

<u>INTERPRETATION:-</u> RHEUMATOID FACTOR (RA):

- 1. Rheumatoid factors (RF) are antibodies that are directed against the Fc fragment of IgG altered in its tertiary structure.

  2. Over 75% of patients with rheumatoid arthritis (RA) have an IgM antibody to IgG immunoglobulin. This autoantibody (RF) is diagnostically useful although it may not be etiologically related to RA.

  3. Inflammatory Markers such as ESR & C-Reactive protein (CRP) are normal in about 60 % of patients with positive RA.

  4. The titer of RF correlates poorly with disease activity, but those patients with high titers tend to have more severe disease course.

- The test is useful for diagnosis and prognosis of rheumatoid arthritis.

### RHEUMATOID ARTHIRITIS:

- 1. Rheumatoid Arthiritis is a systemic autoimmune disease that is multi-functional in origin and is characterized by chronic inflammation of the membrane lining (synovium) joints which ledas to progressive joint destruction and in most cases to disability and reduction of quality life.
- 2. The disease spredas from small to large joints, with greatest damage in early phase. 3. The diagnosis of RA is primarily based on clinical, radiological & immunological features. The most frequent serological test is the

### measurement of RA factor. CAUTION (FALSE POSTIVE):-

- 1. RA factor is not specific for Rheumatoid arthiritis, as it is often present in healthy individuals with other autoimmune diseases and chronic infections.
  2. Non rheumatoid and rheumatoid arthritis (RA) populations are not clearly separate with regard to the presence of rheumatoid factor (RF) (15% of RA patients have a nonreactive titer and 8% of nonrheumatoid patients have a positive titer).
- 3. Patients with various nonrheumatoid diseases,characterized by chronic inflammation may have positive tests for RF. These diseases include systemic lupus erythematosus, polymyositis, tuberculosis, syphilis, viral hepatitis, infectious mononucleosis, and influenza.
- 4. Anti-CCP have been discovered in joints of patients with RA, but not in other form of joint disease. Anti-CCP2 is HIGHLY SENSITIVE (71%) & more specific (98%) than RA factor.
- 5. Upto 30 % of patients with Seronegative Rheumatoid arthiritis also show Anti-CCP antibodies.
- 6. The positive predictive value of Anti-CCP antibodies for Rheumatoid Arthiritis is far greater than Rheumatoid factor.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)





### A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

REPORTING DATE

: 27/Jan/2025 01:03PM

**NAME** : Mr. CHAMAN LAL

**AGE/ GENDER** : 55 YRS/MALE **PATIENT ID** : 1735998

**COLLECTED BY** REG. NO./LAB NO. : 122501270001

REFERRED BY **REGISTRATION DATE** : 27/Jan/2025 08:33 AM BARCODE NO. : 12506697 **COLLECTION DATE** : 27/Jan/2025 09:53AM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

**Value** Unit **Biological Reference interval Test Name** 

### **CLINICAL PATHOLOGY URINE ROUTINE & MICROSCOPIC EXAMINATION**

### **PHYSICAL EXAMINATION**

CLIENT CODE.

QUANTITY RECIEVED	25 r	nl
by DIP STICK/REFLECTANCE SPECTROPHOTOME	TRY	
COLOUR	PALE YELLOW	PALE YELLOW
by DIP STICK/REFLECTANCE SPECTROPHOTOME	TRY	
TRANSPARANCY	HAZY	CLEAR
by DIP STICK/REFLECTANCE SPECTROPHOTOME	TRY	
SPECIFIC GRAVITY	1.02	1.002 - 1.030
	TDV	

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY			
CHEMICAL EXAMINATION			
REACTION by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY	ACIDIC		
PROTEIN by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY	NEGATIVE (-ve)		NEGATIVE (-ve)
SUGAR by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY	NEGATIVE (-ve)		NEGATIVE (-ve)
pH by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY	5.5		5.0 - 7.5
BILIRUBIN by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY	NEGATIVE (-ve)		NEGATIVE (-ve)
NITRITE by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY.	NEGATIVE (-ve)		NEGATIVE (-ve)
UROBILINOGEN by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY	NOT DETECTED	EU/dL	0.2 - 1.0
KETONE BODIES  by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY	NEGATIVE (-ve)		NEGATIVE (-ve)
BLOOD by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY	NEGATIVE (-ve)		NEGATIVE (-ve)
ASCORBIC ACID by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY	NEGATIVE (-ve)		NEGATIVE (-ve)



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST

/HPF

0 - 3



NEGATIVE (-ve)

**MICROSCOPIC EXAMINATION** RED BLOOD CELLS (RBCs)





### A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

NAME : Mr. CHAMAN LAL

**AGE/ GENDER** : 55 YRS/MALE **PATIENT ID** : 1735998

**COLLECTED BY** REG. NO./LAB NO. : 122501270001

REFERRED BY **REGISTRATION DATE** : 27/Jan/2025 08:33 AM **COLLECTION DATE** BARCODE NO. : 12506697 : 27/Jan/2025 09:53AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 27/Jan/2025 01:03PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name	Value	Unit	Biological Reference interval
by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT			
PUS CELLS	3-4	/HPF	0 - 5
by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT			
EPITHELIAL CELLS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	2-4	/HPF	ABSENT
CRYSTALS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	CALCIUM OXALATE (-	++)	NEGATIVE (-ve)
CASTS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	NEGATIVE (-ve)		NEGATIVE (-ve)
BACTERIA by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	NEGATIVE (-ve)		NEGATIVE (-ve)
OTHERS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	NEGATIVE (-ve)		NEGATIVE (-ve)
TRICHOMONAS VAGINALIS (PROTOZOA) by microscopy on centrifuged urinary sediment	ABSENT		ABSENT

**End Of Report** 



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS, MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST

