

PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

NAME : Mrs. KULDEEP KAUR

AGE/ GENDER : 65 YRS/FEMALE **PATIENT ID** :1732633

COLLECTED BY : 122501300005 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 30/Jan/2025 11:44 AM BARCODE NO. : 12506759 **COLLECTION DATE** : 30/Jan/2025 11:45AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 30/Jan/2025 03:17PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name Value Unit **Biological Reference interval**

HAEMATOLOGY

PROTHROMBIN TIME STUDIES (PT/INR)

PT TEST (PATIENT) by PHOTO OPTICAL CLOT DETECTION	14	SECS	11.5 - 14.5
PT (CONTROL) by PHOTO OPTICAL CLOT DETECTION	12	SECS	
ISI by PHOTO OPTICAL CLOT DETECTION	1.1		
INTERNATIONAL NORMALISED RATIO (INR) by PHOTO OPTICAL CLOT DETECTION	1.18		0.80 - 1.20
PT INDEX	85.71	%	

INTERPRETATION:-

- 1.INR is the parameter of choice in monitoring adequacy of oral anti-coagulant therapy. Appropriate therapeutic range varies with the disease and treatment intensity.
- 2. Prolonged INR suggests potential bleeding disorder /bleeding complications
- 3. Results should be clinically correlated.
- 4. Test conducted on Citrated Plasma

RECOMMENDED THERAPEUTIC RANGE FOR ORAL ANTI-COAGULANT THERAPY (INR)			
INDICATION		INTERNATIONAL NORMALIZED RATIO (INR)	
Treatment of venous thrombosis			
Treatment of pulmonary embolism			
Prevention of systemic embolism in tissue heart valves			
Valvular heart disease	Low Intensity	2.0 - 3.0	
Acute myocardial infarction			
Atrial fibrillation	1		
Bileaflet mechanical valve in aortic position	1		
Recurrent embolism			
Mechanical heart valve	High Intensity	2.5 - 3.5	



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)







PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

NAME : Mrs. KULDEEP KAUR

AGE/ GENDER : 65 YRS/FEMALE **PATIENT ID** :1732633

COLLECTED BY : 122501300005 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 30/Jan/2025 11:44 AM BARCODE NO. : 12506759 **COLLECTION DATE** : 30/Jan/2025 11:45AM

CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 30/Jan/2025 03:17PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name Value Unit **Biological Reference interval**

Antiphospholipid antibodies+

The prothrombin time (PT) and its derived measures of prothrombin ratio (PR) and international normalized ratio (INR) are measures of the efficacy of the extrinsic pathway of coagulation. PT test reflects the adequacy of factors I (fibrinogen), II (prothrombin), V, VII, and X. It is used in conjunction with the activated partial thromboplastin time (aPTT) which measures the intrinsic pathway.

The common causes of prolonged prothrombin time are:

1. Oral Anticoagulant therapy.

2.Liver disease.

3. Vit K. deficiency.

4. Disseminated intra vascular coagulation.

5. Factor 5, 7, 10 or Prothrombin dificiency





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)



PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

NAME : Mrs. KULDEEP KAUR

AGE/ GENDER : 65 YRS/FEMALE **PATIENT ID** :1732633

COLLECTED BY REG. NO./LAB NO. : 122501300005

REFERRED BY **REGISTRATION DATE** : 30/Jan/2025 11:44 AM BARCODE NO. : 12506759 **COLLECTION DATE** : 30/Jan/2025 11:45AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 30/Jan/2025 03:50PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Value Unit **Biological Reference interval Test Name**

CLINICAL CHEMISTRY/BIOCHEMISTRY **GLUCOSE POST PRANDIAL (PP)**

GLUCOSE POST PRANDIAL (PP): PLASMA 104.02 NORMAL: < 140.00 mg/dL

by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD) PREDIABETIC: 140.0 - 200.0 DIABETIC: > 0R = 200.0

INTERPRETATION

IN ACCORDANCE WITH AMERICAN DIABETES ASSOCIATION GUIDELINES:

1. A post-prandial plasma glucose level below 140 mg/dl is considered normal.

2. A post-prandial glucose level between 140 - 200 mg/dl is considered as glucose intolerant or prediabetic. A fasting and post-prandial blood test (after consumption of 75 gms of glucose) is recommended for all such patients.

3. A post-prandial plasma glucose level of above 200 mg/dl is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for all such patients. A fasting plasma glucose level in excess of 125 mg/dl on both occasions is confirmatory for diabetic state.

* * * End Of Report * * *



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)

