P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

0171-2532620, 8222896961 pkrjainhealthcare@gmail.com

NAME	**: Mrs. SHIVANI**		
AGE/ GENDER	: 32 YRS/FEMALE	**PATIENT ID**	: 1740910
COLLECTED BY	:	**REG. NO./LAB NO.**	**: 122501310010**
REFERRED BY	:	**REGISTRATION DATE**	: 31/Jan/2025 10:51 AM
BARCODE NO.	: 12506772	**COLLECTION DATE**	: 31/Jan/2025 11:36AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	**REPORTING DATE**	: 31/Jan/2025 01:02PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA		

HAEMATOLOGY

COMPLETE BLOOD COUNT (CBC)

Test Name	Value	Unit	Biological Reference interval
RED BLOOD CELLS (RBCS) COUNT AND INDICES			
HAEMOGLOBIN (HB) *by CALORIMETRIC*	**10.6**^{L}	gm/dL	12.0 - 16.0
RED BLOOD CELL (RBC) COUNT *by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE*	4.23	Millions/cmm	3.50 - 5.00
PACKED CELL VOLUME (PCV) *by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER*	**31.8**^{L}	%	37.0 - 50.0
MEAN CORPUSCULAR VOLUME (MCV) *by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER*	**75.2**^{L}	fL	80.0 - 100.0
MEAN CORPUSCULAR HAEMOGLOBIN (MCH) *by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER*	**25**^{L}	pg	27.0 - 34.0
MEAN CORPUSCULAR HEMOGLOBIN CONC. (MCHC) *by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER*	33.3	g/dL	32.0 - 36.0
RED CELL DISTRIBUTION WIDTH (RDW-CV) *by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER*	14.3	%	11.00 - 16.00
RED CELL DISTRIBUTION WIDTH (RDW-SD) *by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER*	40.9	fL	35.0 - 56.0
MENTZERS INDEX *by CALCULATED*	17.78	RATIO	BETA THALASSEMIA TRAIT: < 13.0 IRON DEFICIENCY ANEMIA: >13.0
GREEN & KING INDEX *by CALCULATED*	25.36	RATIO	BETA THALASSEMIA TRAIT:<= 65.0 IRON DEFICIENCY ANEMIA: > 65.0
WHITE BLOOD CELLS (WBCS)			
TOTAL LEUCOCYTE COUNT (TLC) *by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY*	10390	/cmm	4000 - 11000
DIFFERENTIAL LEUCOCYTE COUNT (DLC)			
NEUTROPHILS *by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY*	66	%	50 - 70
LYMPHOCYTES	30	%	20 - 40

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)

NAME	**: Mrs. SHIVANI**		
AGE/ GENDER	: 32 YRS/FEMALE	**PATIENT ID**	: 1740910
COLLECTED BY	:	**REG. NO./LAB NO.**	**: 122501310010**
REFERRED BY	:	**REGISTRATION DATE**	: 31/Jan/2025 10:51 AM
BARCODE NO.	: 12506772	**COLLECTION DATE**	: 31/Jan/2025 11:36AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	**REPORTING DATE**	: 31/Jan/2025 01:02PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA		

Test Name	Value	Unit	Biological Reference interval
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY			
EOSINOPHILS	**1**^{L}	%	1 - 6
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY			
MONOCYTES	3	%	2 - 12
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY			
BASOPHILS	0	%	0 - 1
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY			
ABSOLUTE LEUKOCYTES (WBC) COUNT			
ABSOLUTE NEUTROPHIL COUNT	6857	/cmm	2000 - 7500
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY			
ABSOLUTE LYMPHOCYTE COUNT	**3117**^{L}	/cmm	800 - 4900
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY			
ABSOLUTE EOSINOPHIL COUNT	104	/cmm	40 - 440
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY			
ABSOLUTE MONOCYTE COUNT	312	/cmm	80 - 880
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY			
ABSOLUTE BASOPHIL COUNT	0	/cmm	0 - 110
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY			
PLATELETS AND OTHER PLATELET PREDICTIVE MARKERS.			
PLATELET COUNT (PLT)	255000	/cmm	150000 - 450000
by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE			
PLATELETCRIT (PCT)	0.26	%	0.10 - 0.36
by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE			
MEAN PLATELET VOLUME (MPV)	10	fL	6.50 - 12.0
by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE			
PLATELET LARGE CELL COUNT (P-LCC)	75000	/cmm	30000 - 90000
by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE			
PLATELET LARGE CELL RATIO (P-LCR)	29.3	%	11.0 - 45.0
by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE			
PLATELET DISTRIBUTION WIDTH (PDW)	15.8	%	15.0 - 17.0
by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE			

NOTE: TEST CONDUCTED ON EDTA WHOLE BLOOD

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)

NAME	**: Mrs. SHIVANI**		
AGE/ GENDER	: 32 YRS/FEMALE	**PATIENT ID**	: 1740910
COLLECTED BY	:	**REG. NO./LAB NO.**	**: 122501310010**
REFERRED BY	:	**REGISTRATION DATE**	: 31/Jan/2025 10:51 AM
BARCODE NO.	: 12506772	**COLLECTION DATE**	: 31/Jan/2025 11:36AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	**REPORTING DATE**	: 31/Jan/2025 01:02PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA		

Test Name	Value	Unit	Biological Reference interval

ERYTHROCYTE SEDIMENTATION RATE (ESR)

ERYTHROCYTE SEDIMENTATION RATE (ESR)	20	mm/1st hr	0 - 20
by RED CELL AGGREGATION BY CAPILLARY PHOTOMETRY			

INTERPRETATION:

1. ESR is a non-specific test because an elevated result often indicates the presence of inflammation associated with infection, cancer and auto-immune disease, but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it.
2. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other test such as C-reactive protein
3. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as some others, such as systemic lupus erythematosus

CONDITION WITH LOW ESR

A low ESR can be seen with conditions that inhibit the normal sedimentation of red blood cells, such as a high red blood cell count (polycythaemia), significantly high white blood cell count (leucocytosis) , and some protein abnormalities. Some changes in red cell shape (such as sickle cells in sickle cell anaemia) also lower the ESR.

NOTE:

1. ESR and C - reactive protein (C-RP) are both markers of inflammation.
2. Generally, ESR does not change as rapidly as does CRP, either at the start of inflammation or as it resolves.
3. **CRP is not affected by as many other factors as is ESR, making it a better marker of inflammation.**
4. If the ESR is elevated, it is typically a result of two types of proteins, globulins or fibrinogen.
5. Women tend to have a higher ESR, and menstruation and pregnancy can cause temporary elevations.
6. Drugs such as dextran, methyldopa, oral contraceptives, penicillamine procainamide, theophylline, and vitamin A can increase ESR, while aspirin, cortisone, and quinine may decrease it

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)

NAME	**: Mrs. SHIVANI**		
AGE/ GENDER	: 32 YRS/FEMALE	**PATIENT ID**	: 1740910
COLLECTED BY	:	**REG. NO./LAB NO.**	**: 122501310010**
REFERRED BY	:	**REGISTRATION DATE**	: 31/Jan/2025 10:51 AM
BARCODE NO.	: 12506772	**COLLECTION DATE**	: 31/Jan/2025 11:36AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	**REPORTING DATE**	: 31/Jan/2025 03:52PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA		

Test Name	Value	Unit	Biological Reference interval

CLINICAL CHEMISTRY/BIOCHEMISTRY

GLUCOSE RANDOM (R)

GLUCOSE RANDOM (R): PLASMA *by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD)*	101.36	mg/dL	NORMAL: < 140.00 PREDIABETIC: 140.0 - 200.0 DIABETIC: > 0R = 200.0

INTERPRETATION

IN ACCORDANCE WITH AMERICAN DIABETES ASSOCIATION GUIDELINES:

1. A random plasma glucose level below 140 mg/dl is considered normal.
2. A random glucose level between 140 - 200 mg/dl is considered as glucose intolerant or prediabetic. A fasting and post-prnadial blood test (after consumption of 75 gms of glucose) is recommended for all such patients.
3. A random glucose level of above 200 mg/dl is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for all such patients. A fasting plasma glucose level in excess of 125 mg/dl on both occasions is confirmatory for diabetic state.

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)

NAME	**: Mrs. SHIVANI**		
AGE/ GENDER	: 32 YRS/FEMALE	**PATIENT ID**	: 1740910
COLLECTED BY	:	**REG. NO./LAB NO.**	**: 122501310010**
REFERRED BY	:	**REGISTRATION DATE**	: 31/Jan/2025 10:51 AM
BARCODE NO.	: 12506772	**COLLECTION DATE**	: 31/Jan/2025 11:36AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	**REPORTING DATE**	: 31/Jan/2025 03:52PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA		

Test Name	Value	Unit	Biological Reference interval

URIC ACID

URIC ACID: SERUM *by URICASE - OXIDASE PEROXIDASE*	3.5	mg/dL	2.50 - 6.80

INTERPRETATION:-

1.GOUT occurs when high levels of Uric Acid in the blood cause crystals to form & accumulate around a joint.
2.Uric Acid is the end product of purine metabolism . Uric acid is excreted to a large degree by the kidneys and to a smaller degree in the intestinal tract by microbial degradation.

INCREASED:-

(A).DUE TO INCREASED PRODUCTION:-

1.Idiopathic primary gout.
2.Excessive dietary purines (organ meats,legumes,anchovies, etc).
3.Cytolytic treatment of malignancies especially leukemais & lymphomas.
4.Polycythemai vera & myeloid metaplasia.
5.Psoriasis.
6.Sickle cell anaemia etc.

(B).DUE TO DECREASED EXCREATION (BY KIDNEYS)

1.Alcohol ingestion.
2.Thiazide diuretics.
3.Lactic acidosis.
4.Aspirin ingestion (less than 2 grams per day).
5.Diabetic ketoacidosis or starvation.
6.Renal failure due to any cause etc.

DECREASED:-

(A).DUE TO DIETARY DEFICIENCY

1.Dietary deficiency of Zinc, Iron and molybdenum.
2.Fanconi syndrome & Wilsons disease.
3.Multiple sclerosis .
4.Syndrome of inappropriate antidiuretic hormone (SIADH) secretion & low purine diet etc.

(B).DUE TO INCREASED EXCREATION

1.Drugs:-Probenecid , sulphinpyrazone, aspirin doses (more than 4 grams per day), corticosterroids and ACTH, anti-coagulants and estrogens etc.

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)

NAME	**: Mrs. SHIVANI**		
AGE/ GENDER	: 32 YRS/FEMALE	**PATIENT ID**	: 1740910
COLLECTED BY	:	**REG. NO./LAB NO.**	**: 122501310010**
REFERRED BY	:	**REGISTRATION DATE**	: 31/Jan/2025 10:51 AM
BARCODE NO.	: 12506772	**COLLECTION DATE**	: 31/Jan/2025 11:36AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	**REPORTING DATE**	: 31/Jan/2025 01:02PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA		

ENDOCRINOLOGY

THYROID FUNCTION TEST: TOTAL

Test Name	Value	Unit	Biological Reference interval
TRIIODOTHYRONINE (T3): SERUM *by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)*	1.23	ng/mL	0.35 - 1.93
THYROXINE (T4): SERUM *by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)*	6.64	µgm/dL	4.87 - 12.60
THYROID STIMULATING HORMONE (TSH): SERUM *by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)*	3.06	µIU/mL	0.35 - 5.50

3rd GENERATION, ULTRASENSITIVE

INTERPRETATION:

TSH levels are subject to circadian variation, reaching peak levels between 2-4 a.m and at a minimum between 6-10 pm. The variation is of the order of 50%.Hence time of the day has influence on the measured serum TSH concentrations. TSH stimulates the production and secretion of the metabolically active hormones, thyroxine (T4)and triiodothyronine (T3).Failure at any level of regulation of the hypothalamic-pituitary-thyroid axis will result in either underproduction (hypothyroidism) or overproduction(hyperthyroidism) of T4 and/or T3.

CLINICAL CONDITION	T3	T4	TSH
Primary Hypothyroidism:	Reduced	Reduced	Increased (Significantly)
Subclinical Hypothyroidism:	Normal or Low Normal	Normal or Low Normal	High
Primary Hyperthyroidism:	Increased	Increased	Reduced (at times undetectable)
Subclinical Hyperthyroidism:	Normal or High Normal	Normal or High Normal	Reduced

LIMITATIONS:-

1. T3 and T4 circulates in reversibly bound form with Thyroid binding globulins (TBG),and to a lesser extent albumin and Thyroid binding Pre Albumin so conditions in which TBG and protein levels alter such as pregnancy, excess estrogens, androgens, anabolic steroids and glucocorticoids may falsely affect the T3 and T4 levels and may cause false thyroid values for thyroid function tests.
2. Normal levels of T4 can also be seen in Hyperthyroid patients with :T3 Thyrotoxicosis, Decreased binding capacity due to hypoproteinemia or ingestion of certain drugs (e.g.: phenytoin , salicylates).
3. Serum T4 levels in neonates and infants are higher than values in the normal adult , due to the increased concentration of TBG in neonate serum.
4. TSH may be normal in central hypothyroidism , recent rapid correction of hyperthyroidism or hypothyroidism , pregnancy , phenytoin therapy.

TRIIODOTHYRONINE (T3)		THYROXINE (T4)		THYROID STIMULATING HORMONE (TSH)	
Age	Refferance Range (ng/mL)	Age	Refferance Range (µg/dL)	Age	Reference Range (µIU/mL)
0 - 7 Days	0.20 - 2.65	0 - 7 Days	5.90 – 18.58	0 - 7 Days	2.43 - 24.3
7 Days - 3 Months	0.36 - 2.59	7 Days - 3 Months	6.39 - 17.66	7 Days - 3 Months	0.58 - 11.00
3 - 6 Months	0.51 - 2.52	3 - 6 Months	6.75 – 17.04	3 Days – 6 Months	0.70 - 8.40
6 - 12 Months	0.74 - 2.40	6 - 12 Months	7.10 – 16.16	6 – 12 Months	0.70 - 7.00

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)

NAME	**: Mrs. SHIVANI**		
AGE/ GENDER	: 32 YRS/FEMALE	**PATIENT ID**	: 1740910
COLLECTED BY	:	**REG. NO./LAB NO.**	**: 122501310010**
REFERRED BY	:	**REGISTRATION DATE**	: 31/Jan/2025 10:51 AM
BARCODE NO.	: 12506772	**COLLECTION DATE**	: 31/Jan/2025 11:36AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	**REPORTING DATE**	: 31/Jan/2025 01:02PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA		

Test Name	Value	Unit	Biological Reference interval

1 - 10 Years	0.92 - 2.28	1 - 10 Years	6.00 - 13.80	1 – 10 Years	0.60 - 5.50
11- 19 Years	0.35 - 1.93	11 - 19 Years	4.87- 13.20	11 – 19 Years	0.50 – 5.50
> 20 years (Adults)	0.35 - 1.93	> 20 Years (Adults)	4.87 - 12.60	> 20 Years (Adults)	0.35– 5.50
RECOMMENDATIONS OF TSH LEVELS DURING PREGNANCY (µIU/mL)					
1st Trimester			0.10 – 2.50		
2nd Trimester			0.20 – 3.00		
3rd Trimester			0.30 – 4.10		

INCREASED TSH LEVELS:

1.Primary or untreated hypothyroidism may vary from 3 times to more than 100 times normal depending upon degree of hypofunction.
2.Hypothyroid patients receiving insufficient thyroid replacement therapy.
3.Hashimotos thyroiditis
4.DRUGS: Amphetamines, iodine containing agents & dopamine antagonist.
5.Neonatal period, increase in 1st 2-3 days of life due to post-natal surge

DECREASED TSH LEVELS:

1.Toxic multi-nodular goiter & Thyroiditis.
2.Over replacement of thyroid hormone in treatment of hypothyroidism.
3.Autonomously functioning Thyroid adenoma
4.Secondary pituitary or hypothalamic hypothyroidism
5.Acute psychiatric illness
6.Severe dehydration.
7.DRUGS: Glucocorticoids, Dopamine, Levodopa, T4 replacement therapy, Anti-thyroid drugs for thyrotoxicosis.
8.Pregnancy: 1st and 2nd Trimester

*** End Of Report ***

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)