

A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

NAME : Mrs. SAROJ

AGE/ GENDER : 59 YRS/FEMALE **PATIENT ID** : 1743665

COLLECTED BY REG. NO./LAB NO. : 122502030014

REFERRED BY **REGISTRATION DATE** : 03/Feb/2025 11:44 AM BARCODE NO. : 12506816 **COLLECTION DATE** :03/Feb/2025 12:01PM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 03/Feb/2025 01:39PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Value Unit **Biological Reference interval Test Name**

HAEMATOLOGY HAEMOGLOBIN (HB)

HAEMOGLOBIN (HB) 9.4^L 12.0 - 16.0 gm/dL

by CALORIMETRIC

INTERPRETATION:-Hemoglobin is the protein molecule in red blood cells that carries oxygen from the lungs to the bodys tissues and returns carbon dioxide from the tissues back to the lungs.

A low hemoglobin level is referred to as ANEMIA or low red blood count.

ANEMIA (DECRESED HAEMOGLOBIN):

- 1) Loss of blood (traumatic injury, surgery, bleeding, colon cancer or stomach ulcer)
- 2) Nutritional deficiency (iron, vitamin B12, folate)
- 3) Bone marrow problems (replacement of bone marrow by cancer)
- 4) Suppression by red blood cell synthesis by chemotherapy drugs
- 5) Kidney failure
- 6) Abnormal hemoglobin structure (sickle cell anemia or thalassemia).

POLYCYTHEMIA (INCREASED HAEMOGLOBIN):

- 1) People in higher altitudes (Physiological)
- 2) Smoking (Secondary Polycythemia)
- 3) Dehydration produces a falsely rise in hemoglobin due to increased haemoconcentration
- 4) Advanced lung disease (for example, emphysema)
- 5) Certain tumors
- 6) A disorder of the bone marrow known as polycythemia rubra vera,
- 7) Abuse of the drug erythropoetin (Epogen) by athletes for blood doping purposes (increasing the amount of oxygen available to the body by chemically raising the production of red blood cells).

NOTE: TEST CONDUCTED ON EDTA WHOLE BLOOD



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)

CLIENT CODE.



PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

: 03/Feb/2025 05:22PM

NAME : Mrs. SAROJ

AGE/ GENDER : 59 YRS/FEMALE **PATIENT ID** : 1743665

COLLECTED BY REG. NO./LAB NO. : 122502030014

REFERRED BY **REGISTRATION DATE** : 03/Feb/2025 11:44 AM BARCODE NO. : 12506816 **COLLECTION DATE** :03/Feb/2025 12:01PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

Test Name Value Unit **Biological Reference interval**

REPORTING DATE

CLINICAL CHEMISTRY/BIOCHEMISTRY

KIDNEY FUNCTION TEST (BASIC)

UREA: SERUM by UREASE - GLUTAMATE DEHYDROGENASE (GLDH)	37.2	mg/dL	10.00 - 50.00
CREATININE: SERUM by ENZYMATIC, SPECTROPHOTOMETERY	1.39 ^H	mg/dL	0.40 - 1.20
BLOOD UREA NITROGEN (BUN): SERUM by CALCULATED, SPECTROPHOTOMETERY	17.38	mg/dL	7.0 - 25.0
BLOOD UREA NITROGEN (BUN)/CREATININE RATIO: SERUM by CALCULATED, SPECTROPHOTOMETERY	12.5	RATIO	10.0 - 20.0
UREA/CREATININE RATIO: SERUM by CALCULATED, SPECTROPHOTOMETERY	26.76	RATIO	
URIC ACID: SERUM by URICASE - OXIDASE PEROXIDASE	3.52	mg/dL	2.50 - 6.80



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)







A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

NAME : Mrs. SAROJ

AGE/ GENDER : 59 YRS/FEMALE **PATIENT ID** : 1743665

COLLECTED BY : 122502030014 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 03/Feb/2025 11:44 AM BARCODE NO. **COLLECTION DATE** :03/Feb/2025 12:01PM : 12506816 CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 03/Feb/2025 05:22PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name Value Unit **Biological Reference interval**

INTERPRETATION:

Normal range for a healthy person on normal diet: 12 - 20

To Differentiate between pre- and postrenal azotemia. INCREASED RATIO (>20:1) WITH NORMAL CREATININE:

1. Prerenal azotemia (BUN rises without increase in creatinine) e.g. heart failure, salt depletion, dehydration, blood loss) due to decreased glomerular filtration rate.

Ž.Catabolic states with increased tissue breakdown.

3.GI hemorrhage.

4. High protein intake.

5. Impaired renal function plus.

6. Excess protein intake or production or tissue breakdown (e.g. infection, GI bleeding, thyrotoxicosis, Cushings syndrome, high protein diet,

burns, surgery, cachexia, high fever)

7. Urine reabsorption (e.g. ureterocolostomy)
8. Reduced muscle mass (subnormal creatinine production)
9. Certain drugs (e.g. tetracycline, glucocorticoids)
INCREASED RATIO (pia (PLIN rices diegrapartic particular partic

1. Postrenal azotemia (BUN rises disproportionately more than creatinine) (e.g. obstructive uropathy).

2. Prerenal azotemia superimposed on renal disease.

DECREASED RATIO (<10:1) WITH DECREASED BUN:

1.Acute tubular necrosis.

2.Low protein diet and starvation.

3. Severe liver disease.

4. Other causes of decreased urea synthesis.

5. Repeated dialysis (urea rather than creatinine diffuses out of extracellular fluid).

6.Inherited hyperammonemias (urea is virtually absent in blood)

7.SIADH (syndrome of inappropiate antidiuretic harmone) due to tubular secretion of urea.

8. Pregnancy

DECREASED RATIO (<10:1) WITH INCREASED CREATININE:

1. Phenacimide therapy (accelerates conversion of creatine to creatinine).

2. Rhabdomyolysis (releases muscle creatinine).

3. Muscular patients who develop renal failure

INAPPROPIATE RATIO:

1. Diabetic ketoacidosis (acetoacetate causes false increase in creatinine with certain methodologies, resulting in normal ratio when dehydration should produce an increased BUN/creatinine ratio).

2. Cephalosporin therapy (interferes with creatinine measurement).

DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)





A PIONEER DIAGNOSTIC CENTRE

: 03/Feb/2025 01:39PM

NAME : Mrs. SAROJ

AGE/ GENDER : 59 YRS/FEMALE **PATIENT ID** : 1743665

COLLECTED BY REG. NO./LAB NO. : 122502030014

REFERRED BY **REGISTRATION DATE** : 03/Feb/2025 11:44 AM BARCODE NO. : 12506816 **COLLECTION DATE** :03/Feb/2025 12:01PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

Test Name Value Unit **Biological Reference interval**

REPORTING DATE

ELECTROLYTES PROFILE: SODIUM AND POTASSIUM

SODIUM: SERUM 139.3 mmol/L 135.0 - 150.0

by ISE (ION SELECTIVE ELECTRODE)

POTASSIUM: SERUM 4.84 mmol/L 3.50 - 5.00

by ISE (ION SELECTIVE ELECTRODE) **INTERPRETATION:-**

SODIUM:-

CLIENT CODE.

Sodium is the major cation of extra-cellular fluid. Its primary function in the body is to chemically maintain osmotic pressure & acid base balance & to transmit nerve impulse.

HYPONATREMIA (LOW SODIUM LEVEL) CAUSES:-

- 1. Low sodium intake.
- 2. Sodium loss due to diarrhea & vomiting with adequate water and iadequate salt replacement.
- 3. Diuretics abuses.
- 4. Salt loosing nephropathy.
- 5. Metabolic acidosis.
- 6. Adrenocortical issuficiency.
- 7. Hepatic failure.

HYPERNATREMIA (INCREASED SODIUM LEVEL) CAUSES:-

- 1. Hyperapnea (Prolonged)
- 2. Diabetes insipidus
- 3. Diabetic acidosis
- 4. Cushings syndrome
- 5.Dehydration

POTASSIUM:-

Potassium is the major cation in the intracellular fluid. 90% of potassium is concentrated within the cells. When cells are damaged, potassium is released in the blood.

HYPOKALEMIA (LOW POTASSIUM LEVELS):-

- 1. Diarrhoea, vomiting & malabsorption.
- 2. Severe Burns.
- 3.Increased Secretions of Aldosterone

HYPERKALEMIA (INCREASED POTASSIUM LEVELS):-

- 1.Oliguria
- 2. Renal failure or Shock
- 3. Respiratory acidosis
- 4. Hemolysis of blood

DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)





A PIONEER DIAGNOSTIC CENTRE

NAME : Mrs. SAROJ

AGE/ GENDER : 59 YRS/FEMALE **PATIENT ID** : 1743665

COLLECTED BY : 122502030014 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 03/Feb/2025 11:44 AM BARCODE NO. : 12506816 **COLLECTION DATE** : 03/Feb/2025 12:01PM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 03/Feb/2025 01:39PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Value Unit **Biological Reference interval Test Name**

ENDOCRINOLOGY THYROID STIMULATING HORMONE (TSH)

THYROID STIMULATING HORMONE (TSH): SERUM μIU/mL 0.35 - 5.50by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

3rd GENERATION, ULTRASENSITIVE

INTERPRETATION:

AGE	REFFERENCE RANGE (μIU/mL)	
0 – 5 DAYS	0.70 - 15.20	
6 Days – 2 Months	0.70 – 11.00	
3 – 11 Months	0.70 - 8.40	
1 – 5 Years	0.70 – 7.00	
6 – 10 Years	0.60 - 5.50	
11 - 15	0.50 - 5.50	
> 20 Years (Adults)	0.27 - 5.50	
PRE	GNANCY	
1st Trimester	0.10 - 3.00	
2nd Trimester	0.20 - 3.00	
3rd Trimester	0.30 - 4.10	

NOTE:-TSH levels are subjected to circardian variation, reaching peak levels between 2-4 a.m and at a minimum between 6-10 pm. The variation is of the order of 50 %. Hence time of the day has influence on the measured serum TSH concentration.

USE: TSH controls biosynthesis and release of thyroid harmones T4 & T3. It is a sensitive measure of thyroid function, especially useful in early or subclinical hypothyroidism, before the patient develops any clinical findings or goitre or any other thyroid function abnormality.

INCREASED LEVELS:

- 1. Primary or untreated hypothyroidism, may vary from 3 times to more than 100 times normal depending on degree of hypofunction.
- 2. Hypothyroid patients receiving insufficient thyroid replacement therapy.
- 3. Hashimotos thyroiditis.
- 4.DRUGS: Amphetamines, Iodine containing agents and dopamine antagonist.
- 5. Neonatal period, increase in 1st 2-3 days of life due to post-natal surge.

DECREASED LEVELS:

- 1. Toxic multi-nodular goitre & Thyroiditis.
- 2. Over replacement of thyroid harmone in treatment of hypothyroidism.
- 3. Autonomously functioning Thyroid adenoma
- 4. Secondary pituatary or hypothalmic hypothyroidism
- 5. Acute psychiatric illness
- 6. Severe dehydration.
- 7.DRUGS: Glucocorticoids, Dopamine, Levodopa, T4 replacement therapy, Anti-thyroid drugs for thyrotoxicosis.



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)





A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

NAME : Mrs. SAROJ

AGE/ GENDER : 59 YRS/FEMALE **PATIENT ID** : 1743665

COLLECTED BY REG. NO./LAB NO. : 122502030014

REFERRED BY **REGISTRATION DATE** : 03/Feb/2025 11:44 AM BARCODE NO. : 12506816 **COLLECTION DATE** : 03/Feb/2025 12:01PM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 03/Feb/2025 01:39PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name Value Unit **Biological Reference interval**

8. Pregnancy: 1st and 2nd Trimester LIMITATIONS:

1.TSH may be normal in central hypothyroidism, recent rapid correction of hyperthyroidism or hypothyroidism, pregnancy, phenytoin therapy.

2. Autoimmune disorders may produce spurious results.

End Of Report



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)