

## PKR JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

### A PIONEER DIAGNOSTIC CENTRE

gm/dL

**RATIO** 

**■** 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

12.0 - 16.0

**NAME** : Mrs. VAANI

**AGE/ GENDER** : 4 YRS/FEMALE **PATIENT ID** : 1743676

**COLLECTED BY** REG. NO./LAB NO. : 122502030015

REFERRED BY **REGISTRATION DATE** : 03/Feb/2025 11:47 AM BARCODE NO. : 12506817 **COLLECTION DATE** : 03/Feb/2025 12:01PM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 03/Feb/2025 01:39PM

12.5

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**RED BLOOD CELLS (RBCS) COUNT AND INDICES** 

HAEMOGLOBIN (HB)

**GREEN & KING INDEX** 

by CALCULATED

by CALORIMETRIC

**Value** Unit **Biological Reference interval Test Name** 

### **HAEMATOLOGY**

### **COMPLETE BLOOD COUNT (CBC)**

., .			
RED BLOOD CELL (RBC) COUNT by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	4.58	Millions/cmm	3.50 - 5.50
PACKED CELL VOLUME (PCV) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	37.1	%	35.0 - 49.0
MEAN CORPUSCULAR VOLUME (MCV) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	81.1	fL	80.0 - 100.0
MEAN CORPUSCULAR HAEMOGLOBIN (MCH) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	27.3	pg	27.0 - 34.0
MEAN CORPUSCULAR HEMOGLOBIN CONC. (MCHC) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	33.6	g/dL	32.0 - 36.0
RED CELL DISTRIBUTION WIDTH (RDW-CV) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	13.2	%	11.00 - 16.00
RED CELL DISTRIBUTION WIDTH (RDW-SD) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	40.5	fL	35.0 - 56.0
MENTZERS INDEX by CALCULATED	17.71	RATIO	BETA THALASSEMIA TRAIT: < 13.0 IRON DEFICIENCY ANEMIA: >13.0

### WHITE BLOOD CELLS (WBCS) TOTAL LEUCOCYTE COUNT (TLC) 9100 /cmm 5000 - 15000 by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY

23.38

### **DIFFERENTIAL LEUCOCYTE COUNT (DLC)**

**NEUTROPHILS** 51 % 50 - 70 by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY LYMPHOCYTES % 41 20 - 45

DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



BETA THALASSEMIA TRAIT:<=

IRON DEFICIENCY ANEMIA: >

65.0





# PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

### A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

NAME : Mrs. VAANI

**AGE/ GENDER** : 4 YRS/FEMALE **PATIENT ID** : 1743676

**COLLECTED BY** : 122502030015 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 03/Feb/2025 11:47 AM BARCODE NO. : 12506817 **COLLECTION DATE** : 03/Feb/2025 12:01PM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 03/Feb/2025 01:39PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name	Value	Unit	Biological Reference interval		
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY					
EOSINOPHILS	$\mathbf{0^L}$	%	1 - 6		
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY					
MONOCYTES	8	%	3 - 12		
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY BASOPHILS	0	%	0 - 1		
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	U	70	0 - 1		
ABSOLUTE LEUKOCYTES (WBC) COUNT					
ABSOLUTE NEUTROPHIL COUNT	4641	/cmm	2000 - 7500		
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY					
ABSOLUTE LYMPHOCYTE COUNT	3731 <sup>L</sup>	/cmm	800 - 4900		
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	A. PKR				
ABSOLUTE EOSINOPHIL COUNT by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	$\mathbf{0_{\Gamma}}$	/cmm	40 - 440		
ABSOLUTE MONOCYTE COUNT	728	/cmm	80 - 880		
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	120	7 CHIIII	00 000		
ABSOLUTE BASOPHIL COUNT	0	/cmm	0 - 110		
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY					
PLATELETS AND OTHER PLATELET PREDICTIVE MARKERS.					
PLATELET COUNT (PLT)	315000	/cmm	150000 - 450000		
by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE					
PLATELETCRIT (PCT)	0.24	%	0.10 - 0.36		
by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE MEAN PLATELET VOLUME (MPV)	8	fL	6.50 - 12.0		
by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	O	IL	0.30 - 12.0		
PLATELET LARGE CELL COUNT (P-LCC)	39000	/cmm	30000 - 90000		
by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE					
PLATELET LARGE CELL RATIO (P-LCR)	12.5	%	11.0 - 45.0		
by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE					
PLATELET DISTRIBUTION WIDTH (PDW) by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	15.5	%	15.0 - 17.0		
NOTE: TEST CONDUCTED ON EDTA WHOLE BLOOD					
NOTE. TEST CONDUCTED ON EDTA WHOLE BEOOD					



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS, MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST





## PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

### A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

**NAME** : Mrs. VAANI

**AGE/ GENDER** : 4 YRS/FEMALE **PATIENT ID** : 1743676

**COLLECTED BY** REG. NO./LAB NO. : 122502030015

REFERRED BY **REGISTRATION DATE** : 03/Feb/2025 11:47 AM BARCODE NO. **COLLECTION DATE** : 03/Feb/2025 12:01PM : 12506817 CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 03/Feb/2025 04:58PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**Value** Unit **Biological Reference interval Test Name** 

### IMMUNOPATHOLOGY/SEROLOGY **C-REACTIVE PROTEIN (CRP)**

C-REACTIVE PROTEIN (CRP) QUANTITATIVE:  $12.86^{H}$ 0.0 - 6.0

**SERUM** by NEPHLOMETRY

**INTERPRETATION:** 

C-reactive protein (CRP) is one of the most sensitive acute-phase reactants for inflammation.

2. CRP levels can increase dramatically (100-fold or more) after severe trauma, bacterial infection, inflammation, surgery, or neoplastic

3. CRP levels (Quantitative) has been used to assess activity of inflammatory disease, to detect infections after surgery, to detect transplant

rejection, and to monitor these inflammatory processes.

4. As compared to ESR, CRP shows an earlier rise in inflammatory disorders which begins in 4-6 hrs, the intensity of the rise being higher than ESR and the recovery being earlier than ESR. Unlike ESR, CRP levels are not influenced by hematologic conditions like Anemia, Polycythemia etc., 5. Elevated values are consistent with an acute inflammatory process.

NOTE:

1. Elevated C-reactive protein (CRP) values are nonspecific and should not be interpreted without a complete clinical history.

2. Oral contraceptives may increase CRP levels.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)







## PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

### A PIONEER DIAGNOSTIC CENTRE

**NAME** : Mrs. VAANI

AGE/ GENDER : 4 YRS/FEMALE **PATIENT ID** : 1743676

**COLLECTED BY** : 122502030015 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 03/Feb/2025 11:47 AM BARCODE NO. : 12506817 **COLLECTION DATE** :03/Feb/2025 12:01PM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 03/Feb/2025 05:16PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**Value** Unit **Biological Reference interval Test Name** WIDAL SLIDE AGGLUTINATION TEST SALMONELLA TYPHI O 1:80 TITRE 1:80 by SLIDE AGGLUTINATION

SALMONELLA TYPHI H 1:40TITRE 1:160by SLIDE AGGLUTINATION SALMONELLA PARATYPHI AH NIL TITRE 1:160by SLIDE AGGLUTINATION SALMONELLA PARATYPHI BH **TITRE** NIL 1:160by SLIDE AGGLUTINATION

**INTERPRETATION:** 

- 1.Titres of 1:80 or more for "O" agglutinin is considered significant.
- 2. Titres of 1:160 or more for "H" agglutinin is considered significant.

- 1.Agglutinins usually appear by 5th to 6th day of illness of enteric fever, hence a negative result in early stage is inconclusive. The titre then rises till 3rd or 4th week, after which it declines gradually.
- 2.Lower titres may be found in normal individuals.
- 3.A single positive result has less significance than the rising agglutination titre, since demonstration of rising titre four or more in 1st and 3rd week is considered as a definite evidence of infection.
- 4.A simultaneous rise in H agglutinins is suggestive of paratyphoid infection.

- 1. Individuals with prior infection or immunization with TAB vaccine may develop an ANAMNESTIC RESPONSE (False-Positive) during an unrelated fever i.e High titres of antibodies to various antigens. This may be differentiated by repitition of the test after a week.
- 2. The anamnestic response shows only a transient rise, while in enteric fever rise is sustained.
- 3.H agglutinins tend to persist for many months after vaccination but O agglutinins tend to disappear sooner i.e within 6 months. Therefore rise in Oagglutinins indicate recent infection.

\*\*\* End Of Report \*\*\*



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)