



# P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

**A PIONEER DIAGNOSTIC CENTRE**

☎ 0171-2532620, 8222896961 ✉ pkrjainhealthcare@gmail.com

NAME	: Mrs. SUPRIYA	PATIENT ID	: 1745046
AGE/ GENDER	: 35 YRS/FEMALE	REG. NO./LAB NO.	: 122502040005
COLLECTED BY	:	REGISTRATION DATE	: 04/Feb/2025 11:15 AM
REFERRED BY	:	COLLECTION DATE	: 04/Feb/2025 11:16AM
BARCODE NO.	: 12506828	REPORTING DATE	: 04/Feb/2025 06:14PM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE		
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

## IMMUNOPATHOLOGY/SEROLOGY

### ANTI THYROID PEROXIDASE (TPO/AMA) ANTIBODIES

ANTI TPO/AMA ANTIBODIES: SERUM by CLIA (CHEMILUMINESCENCE IMMUNOASSAY)	199.81 <sup>H</sup>	IU/mL	0.00 - 10.0 DIABETES (II): < 25.0
---	---------------------	-------	--------------------------------------

#### INTERPRETATION:

1. Thyroperoxidase (TPO) is an enzyme involved in thyroid hormone synthesis, catalyzing the oxidation of iodide on tyrosine residues in thyroglobulin for the synthesis of triiodothyronine and thyroxine (tetraiodothyronine).
2. TPO is a membrane-associated hemo glycoprotein expressed only in thyrocytes and is one of the most important thyroid gland antigens.
3. Anti-TPO is technically superior and a more specific method for measuring thyroid auto-antibodies, It is especially useful in patients presenting with subclinical hypothyroidism where TSH is elevated but Free T4 levels are normal.

#### INCREASED LEVELS (Autoimmune thyroid disease):

1. Hashimoto thyroiditis.
2. Idiopathic myxedema.
3. Graves disease
4. Post-partum thyroiditis.
5. Primary hypothyroidism due to Hashimoto thyroiditis.

#### NOTE:

1. The highest TPO antibody levels are observed in patients suffering from Hashimoto thyroiditis. In this disease, the prevalence of TPO antibodies is about 90% of cases, confirming the autoimmune origin of the disease.
2. These auto-antibodies also frequently occur (60%-80%) in the course of Graves disease.
3. In patients with subclinical hypothyroidism, the presence of TPO antibodies is associated with an increased risk of developing overt hypothyroidism.

\*\*\* End Of Report \*\*\*



DR.VINAY CHOPRA

CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA

CONSULTANT PATHOLOGIST  
MBBS, MD (PATHOLOGY)

