



P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

☎ 0171-2532620, 8222896961 ✉ pkrjainhealthcare@gmail.com

NAME	: Mr. ROHIT	PATIENT ID	: 1752746
AGE/ GENDER	: 23 YRS/MALE	REG. NO./LAB NO.	: 122502110018
COLLECTED BY	:	REGISTRATION DATE	: 11/Feb/2025 11:24 AM
REFERRED BY	:	COLLECTION DATE	: 11/Feb/2025 11:41AM
BARCODE NO.	: 12506956	REPORTING DATE	: 12/Feb/2025 06:25AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE		
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

CLINICAL CHEMISTRY/BIOCHEMISTRY

CREATININE PHOSPHOKINASE (CPK-NAC) TOTAL

CREATININE PHOSPHO KINASE (CPK-NAC)	556.96 ^H	IU/L	24 - 190
by SPECTROPHOTOMETRY			

Interpretation:-

- 1.Serum creatine kinase (CK) activity is greatly elevated, at some time during the course of the disease, in all types of muscular dystrophy, and especially so in Duchenne type, in which levels up to 50 times the upper limit of normal may be encountered.
- 2.In progressive muscular dystrophy, enzyme activity in serum is highest in infancy and childhood (7-10 years of age) and may be elevated long before the disease is clinically apparent.
- 3.Quite high values of CK are noted in viral myositis, polymyositis, and similar muscle diseases.
- 4.However, in neurogenic Parkinsonism, serum enzyme activity is normal. Very high activity is also encountered in malignant hyperthermia.


Significance:-

- 1.An early rise in CK is also seen after an acute MI, with values peaking at 12 to 24 hours and falling back to normal in 3 to 4 days.
- 2.Although total CK activity has been used as a diagnostic test for MI, it has been replaced by the troponin T and I immunoassays, and is no longer the laboratory test choice for diagnosing and monitoring acute infarctions.
- 3.Serum CK activity may increase in patients with acute cerebrovascular disease or neurosurgical intervention and with cerebral ischemia.
- 4.Serum CK activity also demonstrates an inverse relationship with thyroid activity. About 60% of hypothyroid subjects show an average elevation of CK activity 5-fold over the upper reference limit; elevation of as high as 50-fold may also be found.

Note: Exercise and muscle trauma (contact sports, traffic accidents, intramuscular injections, surgery, convulsions, wasp or bee stings, and burns) can elevate serum creatine kinase values.




DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)


DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY)





P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

☎ 0171-2532620, 8222896961 ✉ pkrjainhealthcare@gmail.com

NAME : Mr. ROHIT
AGE/ GENDER : 23 YRS/MALE
COLLECTED BY :
REFERRED BY :
BARCODE NO. : 12506956
CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE
CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA
PATIENT ID : 1752746
REG. NO./LAB NO. : 122502110018
REGISTRATION DATE : 11/Feb/2025 11:24 AM
COLLECTION DATE : 11/Feb/2025 11:41AM
REPORTING DATE : 11/Feb/2025 05:59PM

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

LACTATE DEHYDROGENASE (LDH): SERUM

LACTATE DEHYDROGENASE (LDH): SERUM	528.5 ^H	U/L	225.0 - 450.0
------------------------------------	--------------------	-----	---------------

by BASED ON SCE, SPECTROPHOTOMETRY

INTERPRETATION:-

- 1.Lactate dehydrogenase (LDH) activity is present in all cells of the body with highest concentrations in heart, liver, muscle, kidney, lung, and erythrocytes.
- 2.The test can be used for monitoring changes in tumor burden after chemotherapy, although, lactate dehydrogenase elevations in patients with cancer are too erratic to be of use in the diagnosis of cancer

INCREASED (MARKED) :-

- 1.Megaloblastic anemia.
- 2.Untreated pernicious anemia.
- 3.Hodgkins disease.
- 4.Abdominal and lung cancers.
- 5.Severe shock.
- 6.Hypoxia.


INCREASED (MODERATE):-


- 1.Myocardial infarction (MI).
- 2.Pulmonary infarction and pulmonary embolism.
- 3.Leukemia.
- 4.Hemolytic anemia.
- 5.Infectious mononucleosis.
- 6.Progressive muscular dystrophy (especially in the early and middle stages of the disease)
- 7.Liver disease and renal disease.

NOTE:-

- 1.In liver disease, elevations of LDH are not as great as the increases in aspartate amino transferase (AST) and alanine aminotransferase (ALT).
- 2.Serum LDH may be falsely elevated in otherwise healthy individuals which can be due to mechanical destruction of RBCs. Therefore, Possibility of mechanical errors (Transportation or vigorous shaking) should always be ruled out.




DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)


DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY)





P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

☎ 0171-2532620, 8222896961 ✉ pkrajainhealthcare@gmail.com

TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT.

NAME : Mr. ROHIT
AGE/ GENDER : 23 YRS/MALE **PATIENT ID** : 1752746
COLLECTED BY : **REG. NO./LAB NO.** : 122502110018
REFERRED BY : **REGISTRATION DATE** : 11/Feb/2025 11:24 AM
BARCODE NO. : 12506956 **COLLECTION DATE** : 11/Feb/2025 11:41AM
CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE **REPORTING DATE** : 11/Feb/2025 04:11PM
CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

ENDOCRINOLOGY

THYROID FUNCTION TEST: TOTAL

TRIIODOTHYRONINE (T3): SERUM by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)	1.28	ng/mL	0.35 - 1.93
THYROXINE (T4): SERUM by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)	6.36	µgm/dL	4.87 - 12.60
THYROID STIMULATING HORMONE (TSH): SERUM by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)	2.42	µIU/mL	0.35 - 5.50

3rd GENERATION, ULTRASENSITIVE

INTERPRETATION:

TSH levels are subject to circadian variation, reaching peak levels between 2-4 a.m and at a minimum between 6-10 pm. The variation is of the order of 50%. Hence time of the day has influence on the measured serum TSH concentrations. TSH stimulates the production and secretion of the metabolically active hormones, thyroxine (T4) and triiodothyronine (T3). Failure at any level of regulation of the hypothalamic-pituitary-thyroid axis will result in either underproduction (hypothyroidism) or overproduction (hyperthyroidism) of T4 and/or T3.

CLINICAL CONDITION	T3	T4	TSH
Primary Hypothyroidism:	Reduced	Reduced	Increased (Significantly)
Subclinical Hypothyroidism:	Normal or Low Normal	Normal or Low Normal	High
Primary Hyperthyroidism:	Increased	Increased	Reduced (at times undetectable)
Subclinical Hyperthyroidism:	Normal or High Normal	Normal or High Normal	Reduced

LIMITATIONS:-

1. T3 and T4 circulates in reversibly bound form with Thyroid binding globulins (TBG), and to a lesser extent albumin and Thyroid binding Pre Albumin so conditions in which TBG and protein levels alter such as pregnancy, excess estrogens, androgens, anabolic steroids and glucocorticoids may falsely affect the T3 and T4 levels and may cause false thyroid values for thyroid function tests.
2. Normal levels of T4 can also be seen in Hyperthyroid patients with : T3 Thyrotoxicosis, Decreased binding capacity due to hypoproteinemia or ingestion of certain drugs (e.g.: phenytoin, salicylates).
3. Serum T4 levels in neonates and infants are higher than values in the normal adult, due to the increased concentration of TBG in neonate serum.
4. TSH may be normal in central hypothyroidism, recent rapid correction of hyperthyroidism or hypothyroidism, pregnancy, phenytoin therapy.

TRIIODOTHYRONINE (T3)		THYROXINE (T4)		THYROID STIMULATING HORMONE (TSH)	
Age	Refferance Range (ng/mL)	Age	Refferance Range (µg/dL)	Age	Reference Range (µIU/mL)
0 - 7 Days	0.20 - 2.65	0 - 7 Days	5.90 - 18.58	0 - 7 Days	2.43 - 24.3
7 Days - 3 Months	0.36 - 2.59	7 Days - 3 Months	6.39 - 17.66	7 Days - 3 Months	0.58 - 11.00
3 - 6 Months	0.51 - 2.52	3 - 6 Months	6.75 - 17.04	3 Days - 6 Months	0.70 - 8.40
6 - 12 Months	0.74 - 2.40	6 - 12 Months	7.10 - 16.16	6 - 12 Months	0.70 - 7.00




DR. VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)


DR. YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY)





P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

☎ 0171-2532620, 8222896961 ✉ pkrajainhealthcare@gmail.com

NAME : Mr. ROHIT
AGE/ GENDER : 23 YRS/MALE **PATIENT ID** : 1752746
COLLECTED BY : **REG. NO./LAB NO.** : 122502110018
REFERRED BY : **REGISTRATION DATE** : 11/Feb/2025 11:24 AM
BARCODE NO. : 12506956 **COLLECTION DATE** : 11/Feb/2025 11:41AM
CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE **REPORTING DATE** : 11/Feb/2025 04:11PM
CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name	Value	Unit	Biological Reference interval
1 - 10 Years	0.92 - 2.28	1 - 10 Years	6.00 - 13.80
11- 19 Years	0.35 - 1.93	11 - 19 Years	4.87- 13.20
> 20 years (Adults)	0.35 - 1.93	> 20 Years (Adults)	4.87 - 12.60
RECOMMENDATIONS OF TSH LEVELS DURING PREGNANCY (μ U/mL)			
1st Trimester			0.10 - 2.50
2nd Trimester			0.20 - 3.00
3rd Trimester			0.30 - 4.10

INCREASED TSH LEVELS:

- 1.Primary or untreated hypothyroidism may vary from 3 times to more than 100 times normal depending upon degree of hypofunction.
- 2.Hypothyroid patients receiving insufficient thyroid replacement therapy.
- 3.Hashimotos thyroiditis
- 4.DRUGS: Amphetamines, iodine containing agents & dopamine antagonist.
- 5.Neonatal period, increase in 1st 2-3 days of life due to post-natal surge

DECREASED TSH LEVELS:

- 1.Toxic multi-nodular goiter & Thyroiditis.
- 2.Over replacement of thyroid hormone in treatment of hypothyroidism.
- 3.Autonomously functioning Thyroid adenoma
- 4.Secondary pituitary or hypothalamic hypothyroidism
- 5.Acute psychiatric illness
- 6.Severe dehydration.
- 7.DRUGS: Glucocorticoids, Dopamine, Levodopa, T4 replacement therapy, Anti-thyroid drugs for thyrotoxicosis.
- 8.Pregnancy: 1st and 2nd Trimester




 DR.VINAY CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY & MICROBIOLOGY)


 DR.YUGAM CHOPRA
 CONSULTANT PATHOLOGIST
 MBBS, MD (PATHOLOGY)





P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

☎ 0171-2532620, 8222896961 ✉ pkrjainhealthcare@gmail.com

NAME	: Mr. ROHIT	PATIENT ID	: 1752746
AGE/ GENDER	: 23 YRS/MALE	REG. NO./LAB NO.	: 122502110018
COLLECTED BY	:	REGISTRATION DATE	: 11/Feb/2025 11:24 AM
REFERRED BY	:	COLLECTION DATE	: 11/Feb/2025 11:41AM
BARCODE NO.	: 12506956	REPORTING DATE	: 11/Feb/2025 05:15PM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE		
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

INTACT PARATHYROID HORMONE (PTH)

INTACT PARATHYROID HORMONE (PTH): SERUM	90.9 ^H	pg/mL	12.0 - 72.0
by CLIA (CHEMILUMINESCENCE IMMUNOASSAY)			

Intrepretation:-

Parathyroid hormone (PTH) is produced and secreted by the parathyroid glands, which are located along the posterior aspect of the thyroid gland. The serum calcium level regulates PTH secretion via negative feedback through the parathyroid calcium sensing receptor (CASR). Decreased calcium levels stimulate PTH release. Secreted PTH interacts with its specific type II G-protein receptor, causing rapid increases in renal tubular reabsorption of calcium and decreased phosphorus reabsorption. It also participates in long-term calciostatic functions by enhancing mobilization of calcium from bone and increasing renal synthesis of 1,25-dihydroxy vitamin D, which, in turn, increases intestinal calcium absorption.

The assay is useful for:

- Differential diagnosis of hypercalcemia
- Diagnosis of primary, secondary, and tertiary hyperparathyroidism
- Diagnosis of hypoparathyroidism
- Monitoring end-stage renal failure patients for possible renal osteodystrophy

Interpretation of results:

- An (appropriately) low PTH level and high phosphorus level in a hypercalcemic patient suggests that the hypercalcemia is not caused by PTH or PTH-like substances.
- An (appropriately) low PTH level with a low phosphorus level in a hypercalcemic patient suggests the diagnosis of paraneoplastic hypercalcemia.
- A low or normal PTH in a patient with hypocalcemia suggests hypoparathyroidism.

Low serum calcium and high PTH levels in a patient with normal renal function suggest resistance to PTH action (pseudohypoparathyroidism type 1a, 1b, 1c, or 2) or, very rarely, bio-ineffective PTH.

Elevated PTH value with a normal serum calcium in many cases in India is due to secondary hyperparathyroidism, primary cause being Vitamin D deficiency.




DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)


DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY)





P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

☎ 0171-2532620, 8222896961 ✉ pkrjainhealthcare@gmail.com

NAME : Mr. ROHIT
AGE/ GENDER : 23 YRS/MALE
COLLECTED BY :
REFERRED BY :
BARCODE NO. : 12506956
CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE
CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

PATIENT ID : 1752746
REG. NO./LAB NO. : 122502110018
REGISTRATION DATE : 11/Feb/2025 11:24 AM
COLLECTION DATE : 11/Feb/2025 11:41AM
REPORTING DATE : 11/Feb/2025 03:59PM

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

VITAMINS

VITAMIN D/25 HYDROXY VITAMIN D3

VITAMIN D (25-HYDROXY VITAMIN D3): SERUM
by CLIA (CHEMILUMINESCENCE IMMUNOASSAY)

12.963^L

ng/mL

DEFICIENCY: < 20.0
INSUFFICIENCY: 20.0 - 30.0
SUFFICIENCY: 30.0 - 100.0
TOXICITY: > 100.0

INTERPRETATION:

DEFICIENT:	< 20	ng/mL
INSUFFICIENT:	21 - 29	ng/mL
PREFERRED RANGE:	30 - 100	ng/mL
INTOXICATION:	> 100	ng/mL

- 1.Vitamin D compounds are derived from dietary ergocalciferol (from plants, Vitamin D2), or cholecalciferol (from animals, Vitamin D3), or by conversion of 7- dihydrocholecalciferol to Vitamin D3 in the skin upon Ultraviolet exposure.
- 2.25-OH--Vitamin D represents the main body resevoir and transport form of Vitamin D and transport form of Vitamin D, being stored in adipose tissue and tightly bound by a transport protein while in circulation.
- 3.Vitamin D plays a primary role in the maintenance of calcium homeostatis. It promotes calcium absorption, renal calcium absorption and phosphate reabsorption, skeletal calcium deposition, calcium mobilization, mainly regulated by parathyroid hormone (PTH).
- 4.Severe deficiency may lead to failure to mineralize newly formed osteoid in bone, resulting in rickets in children and osteomalacia in adults.

DECREASED:

- 1.Lack of sunshine exposure.
- 2.Inadequate intake, malabsorption (celiac disease)
- 3.Depressed Hepatic Vitamin D 25- hydroxylase activity
- 4.Secondary to advanced Liver disease
- 5.Osteoporosis and Secondary Hyperparathyroidism (Mild to Moderate deficiency)
- 6.Enzyme Inducing drugs: anti-epileptic drugs like phenytoin, phenobarbital and carbamazepine, that increases Vitamin D metabolism.

INCREASED:

1. Hypervitaminosis D is Rare, and is seen only after prolonged exposure to extremely high doses of Vitamin D. When it occurs, it can result in severe hypercalcemia and hyperphosphatemia.

CAUTION: Replacement therapy in deficient individuals must be monitored by periodic assessment of Vitamin D levels in order to prevent hypervitaminosis D

NOTE:- Dark coloured individuals as compare to whites, is at higher risk of developing Vitamin D deficiency due to excess of melanin pigment which interfere with Vitamin D absorption.



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY)





P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

☎ 0171-2532620, 8222896961 ✉ pkrjainhealthcare@gmail.com

NAME : Mr. ROHIT
AGE/ GENDER : 23 YRS/MALE
COLLECTED BY :
REFERRED BY :
BARCODE NO. : 12506956
CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE
CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

PATIENT ID : 1752746
REG. NO./LAB NO. : 122502110018
REGISTRATION DATE : 11/Feb/2025 11:24 AM
COLLECTION DATE : 11/Feb/2025 11:41AM
REPORTING DATE : 11/Feb/2025 04:11PM

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

VITAMIN B12/COBALAMIN

VITAMIN B12/COBALAMIN: SERUM
by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

163^L pg/mL 200.0 - 1100.0


INTERPRETATION:-

INCREASED VITAMIN B12	DECREASED VITAMIN B12
1.Ingestion of Vitamin C	1.Pregnancy
2.Ingestion of Estrogen	2.DRUGS:Aspirin, Anti-convulsants, Colchicine
3.Ingestion of Vitamin A	3.Ethanol lgestion
4.Hepatocellular injury	4. Contraceptive Harmones
5.Myeloproliferative disorder	5.Haemodialysis
6.Uremia	6. Multiple Myeloma

- Vitamin B12 (cobalamin) is necessary for hematopoiesis and normal neuronal function.
 - In humans, it is obtained only from animal proteins and requires intrinsic factor (IF) for absorption.
 - The body uses its vitamin B12 stores very economically, reabsorbing vitamin B12 from the ileum and returning it to the liver; very little is excreted.
 - Vitamin B12 deficiency may be due to lack of IF secretion by gastric mucosa (eg, gastrectomy, gastric atrophy) or intestinal malabsorption (eg, ileal resection, small intestinal diseases).
 - Vitamin B12 deficiency frequently causes macrocytic anemia, glossitis, peripheral neuropathy, weakness, hyperreflexia, ataxia, loss of proprioception, poor coordination, and affective behavioral changes. These manifestations may occur in any combination; many patients have the neurologic defects without macrocytic anemia.
 - Serum methylmalonic acid and homocysteine levels are also elevated in vitamin B12 deficiency states.
 - Follow-up testing for antibodies to intrinsic factor (IF) is recommended to identify this potential cause of vitamin B12 malabsorption.
- NOTE:**A normal serum concentration of vitamin B12 does not rule out tissue deficiency of vitamin B12. The most sensitive test for vitamin B12 deficiency at the cellular level is the assay for MMA. If clinical symptoms suggest deficiency, measurement of MMA and homocysteine should be considered, even if serum vitamin B12 concentrations are normal.




DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)


DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY)





P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

☎ 0171-2532620, 8222896961 ✉ pkrijainhealthcare@gmail.com

NAME : Mr. ROHIT
AGE/ GENDER : 23 YRS/MALE
COLLECTED BY :
REFERRED BY :
BARCODE NO. : 12506956
CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE
CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

PATIENT ID : 1752746
REG. NO./LAB NO. : 122502110018
REGISTRATION DATE : 11/Feb/2025 11:24 AM
COLLECTION DATE : 11/Feb/2025 11:41AM
REPORTING DATE : 11/Feb/2025 05:15PM

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

VITAMIN B9/FOLIC ACID/FOLATE

VITAMIN B9/FOLIC ACID/FOLATE: SERUM
by CLIA (CHEMILUMINESCENCE IMMUNOASSAY)

6.2 ng/mL

DEFICIENT: < 3.37
INTERMEDIATE: 3.37 - 5.38
NORMAL: > 5.38

INTERPRETATION

RESULT IN ng/mL	REMARKS
0.35 – 3.37	DEFICIENT
3.38 – 5.38	INTERMEDIATE
5.39 – 100.00	NORMAL

NOTE:

1. Drugs like Methotrexate & Leucovorin interfere with folate measurement
2. To differentiate vitamin B12 & folate deficiency, measurement of Methyl malonic acid in urine & serum Homocysteine level is suggested
3. Risk of toxicity from folic acid is low as it is a water soluble vitamin regularly excreted in urine

COMMENTS:

1. Folate plays an important role in the synthesis of purine & pyrimidines in the body and is important for the maturation of erythrocytes.
2. It is widely available from plants and to a lesser extent organ meats, but more than half the folate content of food is lost during cooking.
3. Folate deficiency is commonly prevalent in alcoholic liver disease, pregnancy and the elderly. It may result from poor intestinal absorption, nutrition deficiency, excessive demand as in pregnancy or in malignancy and in response to certain drugs like Methotrexate & anticonvulsants.
4. Decreased Levels Megaloblastic anemia, Infantile hyperthyroidism, Alcoholism, Malnutrition, Scurvy, Liver disease, B12 deficiency, dietary amino acid excess, adult Celiac disease, Tropical Sprue, Crohn's disease, Hemolytic anemias, Carcinomas, Myelofibrosis, vitamin B6 deficiency, pregnancy, Whipple's disease, extensive intestinal resection and severe exfoliative dermatitis

*** End Of Report ***



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY)

